

2-14-2006

# Atrial (? sinus) tachycardia with block (well concealed)

David H. Spodick

*University of Massachusetts Medical School*

Follow this and additional works at: <http://escholarship.umassmed.edu/oapubs>



Part of the [Cardiology Commons](#)

---

## Repository Citation

Spodick, David H., "Atrial (? sinus) tachycardia with block (well concealed)" (2006). *Open Access Articles*. 16.  
<http://escholarship.umassmed.edu/oapubs/16>

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in Open Access Articles by an authorized administrator of eScholarship@UMMS. For more information, please contact [Lisa.Palmer@umassmed.edu](mailto:Lisa.Palmer@umassmed.edu).

## Atrial (? Sinus) Tachycardia With Block (Well Concealed)

David H. Spodick, MD, DSc

[click an author to search for more articles by that author](#)

A 57-year-old woman with coronary heart disease came to the emergency department because of general malaise. Best seen only in leads II, III, and aVF are apparently ectopic (possibly sinus) P waves (P<sup>-</sup>) at approximately 180/min, mostly in a fixed 2:1 ratio to the QRS (Mobitz II AV block). The first three beats, however, may represent Wenckebach (Mobitz I) block with increasing PR on beats 1 and 2 and complete block of the subsequent P (P<sub>x</sub>) wave; this is repeated in beats 8, 9, and 10. The P, as seen in lead V<sub>1</sub> (P), is wide, consistent with interatrial block, but this cannot be interpreted during an ectopic atrial rhythm. Atrial tachycardia with block is usually a toxic rhythm, frequently due to digitalis excess.

*The American Heart Hospital Journal. 4;1:76 (Winter 2006)*

