

8-1-2016

# Advances in Gene Therapy for Diseases of the Eye

Lolita Petit

*University of Massachusetts Medical School, lolita.petit@umassmed.edu*

Hemant Khanna

*University of Massachusetts Medical School, hemant.khanna@umassmed.edu*

Claudio Punzo

*University of Massachusetts Medical School, claudio.punzo@umassmed.edu*

Follow this and additional works at: [http://escholarship.umassmed.edu/metnet\\_pubs](http://escholarship.umassmed.edu/metnet_pubs)

 Part of the [Cellular and Molecular Physiology Commons](#), [Eye Diseases Commons](#), [Molecular Biology Commons](#), and the [Therapeutics Commons](#)



This work is licensed under a [Creative Commons Attribution-Noncommercial 4.0 License](#)

---

## Recommended Citation

Petit, Lolita; Khanna, Hemant; and Punzo, Claudio, "Advances in Gene Therapy for Diseases of the Eye" (2016). *UMass Metabolic Network Publications*. 36.

[http://escholarship.umassmed.edu/metnet\\_pubs/36](http://escholarship.umassmed.edu/metnet_pubs/36)

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Metabolic Network Publications by an authorized administrator of eScholarship@UMMS. For more information, please contact [Lisa.Palmer@umassmed.edu](mailto:Lisa.Palmer@umassmed.edu).

## Advances in Gene Therapy for Diseases of the Eye

Lolita Petit,<sup>1</sup> Hemant Khanna,<sup>1,2</sup> and Claudio Punzo<sup>1,2,\*</sup>

<sup>1</sup>Department of Ophthalmology and Gene Therapy Center, and <sup>2</sup>Department of Neurobiology, University of Massachusetts Medical School, Worcester, Massachusetts.

Over the last few years, huge progress has been made with regard to the understanding of molecular mechanisms underlying the pathogenesis of neurodegenerative diseases of the eye. Such knowledge has led to the development of gene therapy approaches to treat these devastating disorders. Challenges regarding the efficacy and efficiency of therapeutic gene delivery have driven the development of novel therapeutic approaches, which continue to evolve the field of ocular gene therapy. In this review article, we will discuss the evolution of preclinical and clinical strategies that have improved gene therapy in the eye, showing that treatment of vision loss has a bright future.

### INTRODUCTION

VISION IS CONSIDERED by many to be the most important of our five senses. It is a highly complex process that requires the coordinated activity of numerous components in the eye and the brain. The initial steps are performed by the retina, which is the light-sensitive neuronal tissue situated at the back of the eye. When light reaches the retinal rod and cone photoreceptors, photons are absorbed by a photopigment, which activates a cascade that converts the light signal into an electrochemical signal. This is done in collaboration with the retinal pigment epithelium (RPE), which regenerates the visual chromophore. Electrochemical signals are then transferred through bipolar cells to ganglion cells, where they are converted into action potentials that are sent to the brain.<sup>1</sup> Consistent with the crucial role of the retina in vision, the majority of diseases that lead to blindness are caused by an acquired or inherited degeneration of the retina.

As a gene therapy target, the retina is a particularly well-suited organ for therapeutic interventions. The retina is a small tissue, highly compartmentalized, immune-privileged, and easily accessible. Optical transparency of the eye enables safe evaluation

of reporter gene expression and therapeutic effects by noninvasive methods,<sup>2</sup> such as electroretinography (ERG), funduscopy, and optical coherence tomography (OCT). These favorable factors, along with a thorough knowledge of the molecular pathogenesis of many retinal diseases, the development and characterization of animal models that mimic human diseases, and the advances in gene delivery tools, have fueled a rapid development of multiple gene therapy strategies for several forms of retinopathies. This review is focused on emerging strategies that use gene therapy to combat vision loss, particularly for the treatment of retinal diseases caused by mutations that directly affect the photoreceptors.

### GENE REPLACEMENT THERAPY FOR LCA2: THE FIRST SUCCESS OF OCULAR GENE THERAPY

The most successful example of ocular gene therapy was the gene replacement therapy for *RPE65*, Leber's congenital amaurosis 2 (LCA2), an early onset form of autosomal recessive retinal degeneration caused by mutations in the *RPE65* (RPE-specific 65 kDa protein) gene. *RPE65* encodes an isomerase expressed mainly in the RPE that is critical for

\*Correspondence: Dr. Claudio Punzo, Department of Ophthalmology and Gene Therapy Center, University of Massachusetts Medical School, 386 Plantation Street, Worcester, MA 01605. E-mail: claudio.punzo@umassmed.edu

© Lolita Petit et al., 2016; Published by Mary Ann Liebert, Inc. This Open Access article is distributed under the terms of the Creative Commons Attribution Noncommercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.

recycling the visual chromophore involved in the visual cycle.<sup>3</sup> Mutations in *RPE65* result in defective visual pigment formation (both rhodopsin and cone opsin),<sup>4</sup> hence severely affecting photoreceptor function and vision.<sup>5</sup> Large amounts of opsin apoprotein in photoreceptors, as well as accumulation of toxic retinyl esters in the RPE, are thought to promote the progressive death of photoreceptors. Clinical phenotype analyses revealed that the degenerative component of *RPE65*-LCA2 starts at an early age in patients with a functional loss that is much larger than expected for the amount of cells retained.<sup>6</sup> It is this phenotype that provided a very good starting point for a gene-based intervention for this disorder.

Several murine and canine models of LCA2 have shown marked functional benefits with gene therapy.<sup>6</sup> In particular, results obtained in the *Rpe65*<sup>-/-</sup> Briard dog yielded deep excitement in the field because of its more human-like eye anatomy and immune system. The first study was carried out by subretinal delivery of recombinant adeno-associated virus (AAV) 2 vectors expressing the wild-type canine *Rpe65* cDNA under the control of the ubiquitous chicken  $\beta$ -actin (CBA) promoter.<sup>7</sup> This study revealed a dramatic improvement in photoreceptor function and vision in treated dogs. Subsequent dog studies extended the use of other AAV serotypes, including AAV1, AAV4, and AAV5, and different promoters.<sup>8–18</sup> Improvement of vision persisted for over 11 years after a single injection of the vector.<sup>16</sup> In addition, successful restoration of both cone and rod function was achieved in 20 out of 22 treated eyes at more advanced stages of the disease (dogs over 2 years of age).<sup>12,14,16,18</sup>

Based on these preclinical studies, four separate phase I–II clinical trials were initiated, which yielded promising results after subretinal administration of AAV2-h*RPE65* vectors (NCT00481546, NCT00516377, NCT00643747, NCT00749957; Supplementary Table S1; Supplementary Data are available online at [www.liebertpub.com/hum](http://www.liebertpub.com/hum)). Although the designs of these studies varied with respect to the use of the promoter driving the expression of *RPE65*, the volume of vector injected, and the surgical protocol, the data collectively demonstrated safety of AAV2 delivery to the retina.<sup>16,17,19–24</sup> Remarkably, patients in all trials exhibited several aspects of visual improvements within a few months after treatment, though with varying degrees. These results generated excitement in the field. Subsequent treatment of the second eye of LCA2 patients previously treated with the same vector demonstrated both safety

and efficacy, indicating that subretinal administration of AAV2 is feasible even in the case of pre-existing immunity against the vector capsid.<sup>25</sup> These promising results justified the initiation of a phase III clinical trial (NCT00999609) evaluating the treatment of both eyes in patients over 8 years of age. Thus far the trial is confirming the previous visual improvements (Spark Therapeutics Press Release 10/05/2015). AAV2-h*RPE65* is expected to become the first approved gene therapy product in the United States, marking a pivotal step for the entire gene therapy field.<sup>26</sup> In addition, a phase I/II clinical trial (NCT01496040) evaluating the effects of an alternative vector with increased specificity for the RPE (AAV4-RPE65-h*RPE65*) was recently completed.

Although the pioneering *RPE65* trials went far beyond the primary expectations, they uncovered a number of unforeseen challenges, mainly in the magnitude and the longevity of the therapeutic benefits. Contrary to the remarkable disease rescue obtained in the *Rpe65*<sup>-/-</sup> dog, none of the treated *RPE65*-LCA eyes in any clinical trials have shown improvements in retinal function measurable by full-field ERG.<sup>6</sup> The reason for this species-specific difference remains unknown, but is likely related to the extent of disease progression in patients at the time of treatment. Unlike early-onset degeneration in patients with LCA2, the *RPE65*-mutant dogs exhibit no evidence of photoreceptor degeneration for up to 1.5 years of age and show good preservation of photoreceptors in the peripheral retina at 5–7 years of age (57% of rods and 85% of total cones).<sup>6,14,15</sup> Severe photoreceptor degeneration in patient retinas at the time of treatment may have limited the success of the gene therapy strategies. Nonetheless, a recent study investigating efficiency of AAV2-h*RPE65* delivery in older *Rpe65*-mutant dogs showed no correlation between improvement in ERG, the age of the dogs at the time of treatment, and the number of surviving photoreceptors. These results suggest that additional factors could influence the degree of rescue associated with *RPE65* treatment.<sup>14</sup> Health of the remaining RPE and photoreceptor cells may be a factor to consider. One might also predict that disease progression negatively impacts the mean transduction efficiency (i.e., the percentage/density of total RPE cells transduced and the level of *RPE65* expression in transduced cells), thereby reducing the magnitude of rescue. In the animal studies, the effect of *RPE65* replacement is highly dose dependent,<sup>17</sup> with lower doses associated with improvement in visual-guided behavior without detectable rescue by ERG recordings.<sup>14,17</sup> Inefficient

transduction is in agreement with reports of incomplete restoration of dark adaptation in patients after gene therapy,<sup>17,22,27</sup> indicating that partial restoration of the visual cycle may be insufficient to meet the demand of the surviving photoreceptors. Consistent with that, Bainbridge et al. found that *RPE65* RNA level in the human eye is 2.5 times greater than that in the dog eye, which suggests that the demand for RPE65 by the human retina is higher.<sup>17</sup>

A recent long-term assessment of photoreceptor preservation in treated *RPE65*-LCA2 patients also questioned the longevity of the therapeutic effects. Two studies (NCT00481546 and NCT00643747) demonstrated that, despite visual improvement, photoreceptor death remained unchanged and followed the expected natural history of the disease.<sup>16,17,22</sup> Disappointingly, retinal degeneration was associated with a progressive contraction of the areas of improved vision over a period of 5–6 years after intervention<sup>22</sup> and with a sharp decline of retinal sensitivity by 3 years postinjection in a subset of treated patients,<sup>17</sup> indicating a possible loss of the functionally rescued cells. However, loss of a therapeutic effect at 2–4 years posttreatment has not yet been reported by Spark Therapeutics, potentially reflecting differences in vector design or manufacturing. Nonetheless, the loss of the therapeutic effect observed by Jacobson et al.<sup>22</sup> and Bainbridge et al.<sup>17</sup> has important implications for the evaluation of any retinal gene therapy, and several hypotheses have been formulated to explain it. Cideciyan et al. proposed that gene replacement therapy does not slow the degenerative component of the disease if the intervention is initiated after a threshold of accumulative molecular changes has been reached in RPE and/or photoreceptors.<sup>16</sup> Alternatively, the continuous changes that occur in the tissue not exposed to the vector could have overcome the small number of stably rescued RPE/photoreceptors in these patients.<sup>28</sup> Should results of the phase III clinical trial confirm the lack of long-term efficacy, appreciating the mechanism will be important for managing expectations about the benefits of gene therapy for each patient, as well as for designing strategies to improve the durability of the treatment. If a “point of no return” exists, alleviating stress in cells before treatment may be a solution to extend the efficacy of the gene therapy. Otherwise, it may be necessary to increase the number of transduced cells and also protect rescued photoreceptors against secondary degeneration. Such strategies are being developed for use in other retinal diseases<sup>28,29</sup> and could be adapted to the *RPE65* deficiency.

## RECENT PROGRESS IN CLINICAL APPLICATIONS OF RETINAL GENE REPLACEMENT THERAPY

### Gene replacement therapy for Mer proto-oncogene tyrosine kinase (*MERTK*)- and Rab-escort protein 1 (*REP1*)-associated inherited retinal diseases

Pioneering results of *RPE65*-LCA2 trials laid the groundwork for the initiation of trials for different forms of inherited retinopathies caused by mutations in another RPE-specific gene (*MERTK* retinitis pigmentosa) and in a gene required for both RPE and photoreceptor survival (*REP1* choroideremia).

*MERTK* is involved in the engulfment of outer segment debris by the RPE. Patients with *MERTK* deficiency exhibit severe dysfunction of both rods and cones, associated with an early-onset and progressive degeneration of photoreceptors. In the first clinical safety study for *MERTK* retinitis pigmentosa (NCT01482195),<sup>30</sup> six patients received a submacular injection of AAV2-VMD2-h*MERTK*, in which the *MERTK* cDNA was expressed under the control of the RPE-specific VMD2 (vitelliform macular dystrophy) promoter. No complications attributed to the vector were observed, and three patients displayed improved visual acuity in the treated eye within the first week/month postintervention. However, the improvement declined to baseline 2 years posttreatment in two of the patients. Contrary to the *LCA2* gene therapy studies, this functional decline was not associated with major changes in retinal thickness, as assessed by OCT. However, variability in OCT measurements in these patients with nystagmus was very large. It is possible that the quality of treatment, including the number of cells treated and expression levels of *MERTK*, may have been too low to promote long-term benefits.

Recently, promising results from a multicenter phase I/II study of AAV2-mediated gene replacement therapy for choroideremia (CHM) were reported (NCT01461213),<sup>31,32</sup> strengthening the use of gene therapy for retinal diseases. CHM is a severe X-linked recessive disorder leading to blindness through progressive degeneration of the choroid, RPE, and photoreceptors caused by the loss of function of the Rab escort protein 1 (*REP1*). CHM can be identified in childhood; however, it differs from *RPE65*-LCA2 in that most patients retain 20/20 vision because of fairly well-preserved central retinal cones until the fifth decade of life. The main objective is to preserve this area from further degeneration, which will require long-term follow-up to detect the outcome of the gene therapy. Importantly, no retinal thinning or

loss of visual acuity was observed despite the surgically induced retinal detachment, establishing a favorable safety profile for this gene transfer protocol targeting the fovea. In addition, at 6 months posttreatment, significant improvement in retinal sensitivity was noted in the five patients who received the highest vector dose, as well as large gains in visual acuity in the two patients in whom visual acuity was already reduced at baseline. Preservation or gains in visual acuity were sustained until at least 3.5 years after treatment, while, over the same period, visual acuity in the control noninjected eyes decreased progressively.<sup>32</sup> It will be interesting to see if photoreceptor degeneration is prevented in these patients in a long-term follow-up examination. A 30-patient phase II study (NCT02341807) has started to further assess the functional and anatomical outcomes. Additionally, a phase I/II clinical trial using AAV5, a serotype with increased tropism for photoreceptors as compared with AAV2, is planned (Horama/Nantes Hospital, France). This approach may maximize the efficacy of the therapy as it was demonstrated that rod photoreceptors could degenerate independently from the RPE.<sup>33,34</sup>

#### Additional clinical trials of retinal gene replacement therapy

To date, eight clinical trials testing gene replacement for four other retinal diseases are in progress (Supplementary Table S1). All of them build on the existing AAV2 platform, with the exception of two trials using the equine infectious anemia virus (EIAV)-based lentiviral vectors for the treatment of Stargardt disease (NCT01367444) and Usher type 1B syndrome (NCT01505062). These diseases are caused by mutations in the *ABCA4* and *MYO7A* genes respectively, which are too large to be packaged in an AAV.<sup>35</sup> Lentivirus, which has a higher cargo capacity than AAV, has therefore been chosen as an alternative to AAV.

*ABCA4* is localized in the outer segments of photoreceptors and acts as an important membrane transporter for the recycling of the visual chromophore. *ABCA4* loss of function is associated with accumulation of toxic products in RPE cells, followed by severe RPE and macular photoreceptor death. A proof-of-concept study in the *Abca4*<sup>-/-</sup> mice demonstrated beneficial effects after subretinal injection of EIAV-*Abca4* at postnatal day (PN)4–PN5.<sup>36</sup> *MYO7A* is expressed in cochlear hair cells of the inner ear as well as in retinal photoreceptors and the RPE,<sup>37</sup> where it plays a role in multiple cellular processes, including endocytosis and cellular transport. Although the amount of

*MYO7A* in photoreceptors is lower than that in the RPE, photoreceptors are affected before RPE cells in patients with Usher1B, indicating that photoreceptors are important cells to target in this disease as well. The ability of LV-*MYO7A* to restore RPE abnormalities has been shown in the *Myo7a*<sup>-/-</sup> mouse after subretinal injection of the vector at birth.<sup>38,39</sup> However, a major concern of lentiviral vectors for clinical use is its relative inability to transduce postmitotic photoreceptors. Physical barriers, which are not present in the newborn rodent retina, have been hypothesized to dramatically limit access of the large lentivirus particles to adult photoreceptors.<sup>40</sup> At this point the results of these two clinical trials remain unknown. Nonetheless, these results will provide valuable information not only regarding the efficacy of lentivirus in the retina, but also to determine whether lentivirus may serve as a safe alternative vector for RPE diseases in which high level of transgene expression is required.

#### PRECLINICAL ADVANCES IN GENE-SPECIFIC THERAPY FOR PHOTORECEPTOR DISEASES

One of the major challenges over the next 20 years will be to initiate treatment of many retinal disorders in which the disease-causing mutation is primarily expressed in photoreceptors. Compared with RPE-associated diseases, primary photoreceptor dystrophies have been considered as more difficult to treat. This is because photoreceptors are directly impaired by the genetic mutation and, second, because the efficiency of photoreceptor transduction is relatively lower compared with that of RPE cells.<sup>41</sup> AAV2 and lentiviral vectors, though excellent vectors for transducing the RPE, were found to be inefficient in transducing photoreceptor cells.<sup>42,43</sup> However, several novel AAV serotypes have been identified and characterized. Among these, AAV5, 7, 8, 9, and rh10, as well as vectors with modified capsid proteins, show remarkable improvement in terms of photoreceptor transduction efficiency when compared with AAV2 (for review, see refs.<sup>41,44</sup>). The main question that remains to be answered now is whether the use of these serotypes can offer clinical benefits. To illustrate advances in gene-specific therapies for recessive photoreceptor diseases, we will describe two approaches in more detail: (1) the promising application of gene therapy for the treatment of stationary disorders, in particular cone disorders, and (2) the novel approaches for the treatment of progressive retinal dystrophies initiated by defects that are either in rods or in both rods and cones.

### Gene replacement therapy in animal models of stationary photoreceptor disorders

Despite the fact that stationary photoreceptor diseases are relatively rare, they are ideal translational models for the development of gene replacement therapies targeting photoreceptors. Stationary disorders are associated with congenital retinal dysfunction and can thus be diagnosed early. Their slowly progressive degenerative nature presents therefore a wide window of opportunity for intervention and effects of the therapy can be rapidly assessed through the restoration of retinal function.

Achromatopsia (ACHM) is an autosomal recessive disease associated with severe cone dysfunction, caused by a loss of function of some cone-specific proteins. To date, four small<sup>45–48</sup> and three large animal models<sup>29,49,50</sup> of ACHM have been successfully treated using AAV-mediated gene replacement therapy (Supplementary Table S2). The first report of efficient gene therapy for ACHM was obtained in the *cpfl3* (cone photoreceptor functional loss 3) mouse model of GNAT2 (guanine nucleotide binding G-protein) deficiency.<sup>45</sup> This mouse model retains ~25% of the normal cone-mediated ERG response up to 4 weeks of age with no detectable responses by 9 months of age. In contrast, cone structural integrity is maintained for at least 14 weeks. Subretinal injection of AAV5 encoding the mouse *Gnat2* under the control of the human red green cone opsin promoter (AAV5-PR2.1-*mGnat2*) at PN23-PN29 in *cpfl3* mice restored cone ERG responses and visual-guided behavior to levels indistinguishable from age-matched controls in 80% of treated eyes, for at least 7 months. Interestingly, when the *cpfl3* mice were treated at later stages of the disease (>9 months of age), the degree of rescue was variable, with only one eye showing cone ERG responses within the normal range.<sup>45</sup>

Consistent with this, Carvalho et al. demonstrated that an optimal therapeutic window is present in the *Cngb3*<sup>-/-</sup> (cone-specific cyclic nucleotide gated channel subunit b3) mouse model of ACHM,<sup>48</sup> in which cone degeneration is comparable to that of the *cpfl3* mice. While subretinal injection of AAV8-*mCAR*-*hCNGB3*, expressing *CNGB3* under the control of the mouse cone arrestin promoter at PN15, resulted in long-term restoration of cone-mediated ERG responses up to 90% of wild-type levels, treatment at PN90 and PN180 resulted in functional rescue at 70–80% and 60–70% of wild-type levels, respectively. Restoration of visual acuity was not possible after PN90. Komaromy et al. also showed restoration of vision in two canine

models of *CNGB3*-ACHM using AAV5, marking an important milestone in the photoreceptor gene therapy field.<sup>49</sup> However, the magnitude and longevity of the therapy in dogs was also age dependent.<sup>29,49</sup> Younger dogs (<6 months) treated with AAV5-PR2.1-*cCngb3* displayed the most sustained therapeutic effects, with a restoration of cone function up to 5–10% of wild-type levels, and a restoration of visual-guided behavior in bright light. Therapeutic effects were maintained for at least 33 months postinjection in two dogs. In contrast, only two out of seven dogs treated with less efficient vectors showed any sustained response, whereas the others had either a transient or no functional cone response. At older ages, only 1 dog out of 11 showed any sustained response. The authors hypothesized that the failure might result from the improper reassembly of the components of the phototransduction cascade when retinas were treated at more advanced stages of the disease. Combination of gene therapy with administration of the ciliary neurotrophic factor (CNTF), which promotes outer segment deconstruction and reconstruction, increased the efficacy and durability of gene therapy in older dogs.<sup>29</sup> CNTF alone also transiently restored cone function,<sup>29,51</sup> an effect not reproduced in five patients with *CNGB3*-ACHM.<sup>52</sup>

Taken together these results indicate that the health of photoreceptor cells at the time of intervention and the associated efficacy in transgene expression could be limiting factors for a functional rescue after gene therapy in ACHM. Nonetheless, accurate selection of patients, choice of area for treatment, and selection of an optimal vector system may realistically allow for a favorable outcome in humans. Based on the encouraging results using animal models, Applied Genetic Technologies Corporation is conducting a phase I–II clinical trial to evaluate the potential of an optimized vector system<sup>53</sup> (AAV2tYF-PR1.7-codon-optimized *hCNGB3* [Supplementary Table S1] for the treatment of *CNGB3*-ACHM).

X-linked retinoschisis (XLRS) is characterized by compromised retinal integrity and a subsequent slow loss of central photoreceptors. Juvenile XLRS is caused by loss-of-function mutations in the *RS1* gene (retinoschisin), which encodes for a protein primarily expressed and secreted from photoreceptors and bipolar cells. The function of *RS1* is unknown, but the protein is thought to play a role in cell adhesion and in the organization of the photoreceptor-bipolar cell synapse.<sup>54</sup> Numerous studies have shown that delivery of the *RS1* gene to the photoreceptors, by subretinal (AAV5-*mOP*-*hRS1*; mouse opsin promoter<sup>55</sup>) and intravitreal

injections (AAV2-CMV-*Rs1*,<sup>56</sup> AAV8-RS1-h*RS1*,<sup>57</sup> AAV8-RS/IRBP-h*RS1*; under the control of RS/interphotoreceptor binding protein promoter<sup>58</sup> and AAV7m8-RHO-h*RS1*; rhodopsin promoter<sup>59</sup>) can lead to long-term structural and functional preservation. Nonetheless, gene transfer at advanced stages of the disease (>7 months of age) showed no improvements of ERG responses.<sup>60</sup> Applied Genetic Technologies Corporation recently initiated a phase I/II clinical trial to evaluate the safety and efficacy of intravitreal injection of AAV2tYF-CBA-h*RS1* in patients with XLRS (NCT02599922; Supplementary Table S1). AAV8-RS/IRBP-h*RS1* also entered phase III (NCT02317887; Supplementary Table S1).

### Gene replacement therapy in models of progressive photoreceptor disorders

Progressive photoreceptor degenerations are the most common causes of complete blindness in humans. When inherited, they are primarily caused by mutations in genes expressed in rods only (retinitis pigmentosa [RP]), or in both rods and cones (LCA, RP, and cone-rod dystrophies).<sup>61</sup> The fact that loss of cones is associated with the most devastating aspect in these diseases points to the need to preserve cone function and survival as the primary therapeutic outcome. However, as cone death is invariably linked to the death of rods,<sup>62</sup> the therapy should be able in most cases to also robustly protect rod photoreceptors from degeneration.

Mutations in the *PDE6 $\beta$*  gene, which encodes the  $\beta$  subunit of the rod phosphodiesterase (PDE6) enzyme, are associated with one of the most common and aggressive forms of recessive rod-initiated RP. In the absence of *PDE6 $\beta$* , PDE6 activity is severely impaired and high levels of intracellular cGMP and  $\text{Ca}^{2+}$  accumulate, leading to rod death. Rod dysfunction with early-onset degeneration is collectively seen in all animal models of *PDE6 $\beta$* -RP, including the *rd1*,<sup>63–66</sup> *rd10*,<sup>67,68</sup> and *Pde6 $\beta$* -H620Q<sup>69</sup> mice, as well as the *PDE6 $\beta$* -deficient *rod cone dysplasia (rd1)*<sup>70,71</sup> and *cone rod dystrophy 1 (crd1)* dogs.<sup>72</sup> In all of these models, rod loss is always followed by a mutation-independent loss of cones.

Gene therapy approaches to delay rod death were first employed in the *rd1* mouse, in which the majority of rods are lost by 3 weeks of age. Initial attempts to treat photoreceptor degeneration in these mice were made using adenoviral,<sup>73,74</sup> lentiviral,<sup>69,75</sup> and AAV2 vectors.<sup>76</sup> These gene delivery tools result in overall poor photoreceptor transduction, and these studies revealed minimal structural and functional ERG rescue. Using more efficient gene delivery tools such as AAV8-RHO-h*PDE6 $\beta$*

and AAV9-RHO-h*PDE6 $\beta$* , long-term (13 months) functional and structural improvements were obtained after subretinal injection of *rd1* mice at PN9, after removal of a confounding mutation in the *Gpr179* gene.<sup>77</sup>

Benefits of the levels and kinetics of transgene expression were also evident in the slightly slower degenerating *rd10* mouse model, where injection of an AAV8-m*Pde6 $\beta$*  (but not AAV5-m*Pde6 $\beta$* ) vector at PN14 resulted in a robust functional rescue.<sup>78</sup> In comparison, subretinal injection of AAV5-CMV-h*PDE6 $\beta$*  or AAV8-CMV-h*PDE6 $\beta$*  in the same animal model at PN2, when rod differentiation is incomplete and rod transduction is inefficient (Petit L. and Punzo C., unpublished data), had only minimal therapeutic effects.<sup>79</sup> Another study showed that AAV8(Y773F), a vector that results in high levels of transgene expression within 2–3 days in nearly 100% of rods,<sup>80</sup> successfully altered the course of the disease.<sup>81</sup> In *rd1* dogs, injections of AAV5 and AAV8 vectors expressing *cPde6 $\beta$*  under the control of the rhodopsin kinase (RK) promoter (AAV5-hRK1-*cPde6 $\beta$*  and AAV8-hRK1-*cPde6 $\beta$* ) resulted in similar levels of functional and morphological rescue after subretinal delivery at PN20,<sup>82</sup> for at least 40 months posttreatment.<sup>83</sup> Intervention before the onset of photoreceptor degeneration (PN25) and a relatively slower progression of the disease when compared with the mouse models may have likely allowed for therapeutic transgene expression in a high proportion of photoreceptors in time, independent of the serotype used. Importance of the quality of treatment is in agreement with results obtained by the group of S. Tsang, who tested whether lack of sustained benefit in the *PDE6 $\beta$* -deficient mouse after gene therapy is (1) because of insufficient transduction efficiency and/or (2) because the disease is too advanced at the time of treatment. Using a Cre-inducible Cre-loxP rescue allele, they demonstrated that photoreceptor degeneration in *Pde6 $\beta$* -H620Q/*LoxP* is halted if DNA recombination is initiated at early, mid, and late stages of the disease. In this study, *Pde6 $\beta$*  expression after recombination was considered optimal/not limiting.<sup>84</sup>

The potential of next-generation gene delivery tools was illustrated in animal models of other forms of severe early-onset rod-cone dystrophies (Supplementary Table S2). The *PDE6 $\alpha$* -deficient mouse has a faster rate of photoreceptor degeneration compared with the *rd10* mouse, with the loss of 30% of rods already evident by PN14. Using AAV8(Y788F)-RHO-*Pde6 $\alpha$* , Wert et al. showed that treatment at PN5 and PN21 resulted in a dramatic preservation of retinal structure and

cone function.<sup>85,86</sup> These results were very exciting, although the overall rescue of rod function was too low to result in a detectable difference by ERG. The murine model of CNGB1 retinopathy also benefited from subretinal delivery of AAV8(Y733F)-RHO-m*Cngb1a*.<sup>87</sup> Like *rd1* mice, *Cngb1*<sup>-/-</sup> mice have no recordable rod response by ERG. However, rod degeneration progresses more slowly than in murine models of PDE6-deficiency, as 50–70% of rods are still present at 6 months of age. Treatment of *Cngb1*<sup>-/-</sup> mice at PN14 resulted in dramatic restoration of rod function, accounting for 33% of wild-type levels corresponding to the surface of the retina directly exposed to the vector. Improvement of the functional component of the disease was associated with a preservation of 50–70% of rods in treated eyes at 12 months of age. More recently, Palfi et al. demonstrated that AAVrh10 transduces rods comparably to AAV8 in a degenerating retina.<sup>88</sup> Using this serotype and an optimized murine rhodopsin promoter, they showed great progress in the treatment of the *rhodopsin* knockout mouse, compared with similar doses of AAV5.<sup>89</sup> Nonetheless, despite the fact that therapeutic benefits were observed up to 11 months after treatment with AAVrh10, degeneration was not arrested in treated retinas.<sup>88</sup>

While most of the gene replacement therapies for photoreceptor diseases discussed here target either cones or rods, it is useful to treat the two photoreceptor subtypes simultaneously. This is because many inherited photoreceptor degenerations are caused by mutations in genes expressed in both rods and cones. The short human rhodopsin kinase 1 (hRK1) promoter was the first well-defined promoter able to drive efficient transgene expression in both cell types, when used in conjunction with AAV.<sup>90</sup> While the efficiency of cone transduction remains very low as compared with rod transduction, validation of this promoter in small<sup>91</sup> and large animal models,<sup>91–93</sup> along with the development of next-generation of vectors, led in the past few years to an exponential growth of retinal gene transfer studies targeting both rods and cones.

A case in point is the gene replacement therapy for the retina-specific guanylate cyclase (*GUCY2D*), which is expressed exclusively in rods and cones and constitutes one of the most common causes of LCA.<sup>61</sup> The GC1-deficient mouse undergoes severe cone dysfunction before cone degeneration. Rods retain 30–50% of their function and do not degenerate, because of the presence of a second guanylate cyclase (GC2). Subretinal injection of AAV5-CBA-bovine *Gc1* at PN21 had no effect in the GC1-deficient mouse model.<sup>94</sup> In contrast, AAV5-CBA-m*Gucy2e* and AAV5-hRK1-m*Gucy2e*

at PN14 restored 45% of cone ERG responses and preserved cone survival for at least 9 months post-injection.<sup>95</sup> Subsequently, injection of AAV8-hRK1 vectors carrying the murine *Gucy2e* or human *GUCY2D* cDNA at PN10 provided a 65% rescue of cone ERG responses, cone vision and cone survival for up to 6 months, as well as a 35% rescue of rod function.<sup>96</sup> A more recent study reported a restoration of 54% and 38% of cone ERG responses in mice treated at PN21 with AAV8-CMV-h*GUCY2D* and AAV8-hRK1-h*GUCY2D*, respectively.<sup>91</sup>

Comparison of AAV5-hRK1-m*Gucy2e* and AAV8(Y733F)-hRK1-m*Gucy2e*/h*GUCY2D* confirmed the superiority of AAV8(Y733F) in restoring cone function in *Gucy2e*<sup>-/-</sup> mice<sup>97</sup> and *Nrl*<sup>-/-</sup> *Gucy2e*<sup>-/-</sup> mice.<sup>98</sup> This difference likely reflects the ability of AAV8(Y733F) to drive faster transgene expression than AAV5, and indicates that this temporal difference is important in terms of the ability to restore function. However, it is unclear whether there is a higher overall number of preserved cones and/or a better functional rescue of transduced cones. Regarding rod function, there were no differences between the AAV5 and the AAV8(Y733F) vectors.

The ability of AAV8(Y733F)-hRK1-m*Gucy2e* to rescue both cones and rods was confirmed in the GC1/GC2 double-deficient mouse, which exhibits complete loss of both cone and rod ERG responses, as well as slow degeneration of rods. Both cone and rod ERG responses were restored to 42–44% of wild-type levels after treatment at PN18, and to 26–29% of wild-type levels after treatment at PN108. The rescue remained stable for at least 1 year posttreatment. However, only intervention before PN108 slowed photoreceptor degeneration.<sup>99</sup>

Interestingly, benefits of AAV8-based vectors over AAV5 have also been observed in the *RPGRiP1* (retinitis pigmentosa GTPase regulator interacting protein 1)-deficient<sup>100,101</sup> and *AiPL1* (aryl hydrocarbon receptor interacting protein-like 1)-deficient<sup>102–104</sup> mouse models of early-onset severe photoreceptor dystrophies, but not in the *RPGRIP1*-deficient dog, in which AAV5 and AAV8 gave similar therapeutic effects,<sup>93</sup> probably reflecting differences in the kinetics of photoreceptor loss and timing of therapeutic transgene expression between these species.

The first evidence of the efficacy of gene replacement therapy in a large animal model of severe photoreceptor dystrophies was obtained in the XLPRA1 (X-linked progressive retinal atrophy) and XLPRA2 canine models of *RPGR*-X-linked RP.<sup>105,106</sup> In these dogs, Beltran et al. evaluated the efficacy of AAV5-hRK1-h*RPGR* and AAV5-IRBP-h*RPGR* to mitigate retinal degeneration.<sup>105</sup>



The XLPRA1 dogs were treated at 28 weeks of age, before any apparent signs of photoreceptor degeneration. In contrast, XLPRA2 dogs were treated at the onset of the disease, which is at 5 weeks of age. In both cases, treatment resulted in convincing preservation of retinal structure in the vector-exposed area for a 2-year period.<sup>105</sup> However, the benefits seen in XLPRA2 dogs treated with AAV5-hRK1-hRPGR or with low dose of AAV5-IRBP-hRPGR were smaller likely because these animals were treated at the onset of the disease.<sup>106</sup> Early evaluation of the effects of gene transfer on retinal function was difficult, because of the long-term preservation of functional photoreceptors in areas not exposed by the vector.<sup>106</sup> Nonetheless, retention of rod and cone function was clearly seen in XLPRA2 dogs up to 3 years of age with improved long-term vision.<sup>105</sup>

Interestingly, XLPRA2 dogs treated at 12 weeks of age (loss of ~40% of the total photoreceptors) showed an initial decline in outer nuclear layer (ONL) thickness, but ONL loss in the vector-exposed area was halted from 31 weeks to 2.7 years of age. XLPRA2 dogs treated at 26 weeks of age (loss of 50–60% of total photoreceptors) showed a similar profile, but with a decrease and stabilization of ONL thickness after 52 weeks of age. A higher proportion of nontreated photoreceptors could explain this delay in ONL stabilization after late intervention. Remarkably, ERG analysis showed preserved rod function in all 3 dogs treated at this late stage, and preserved cone function in 2 out of 3 dogs, accounting for 8% of wild-type ERG levels, providing very exciting proof-of-concept to support a future clinical trial.<sup>105</sup>

#### **Additional improvements**

Extensive efforts have also been made to expand the applicability of gene-specific therapy in the eye, including the treatment of photoreceptor dystrophies caused by mutations in large genes, the treatment of autosomal dominant retinal diseases, and the delivery of transgene to a large retinal surface by intravitreal injection. The reader is referred to recent reviews for further information about the strategies that are being devised to overcome these limitations.<sup>107–109</sup>

### **BROADENING THE SPECTRUM OF TREATABLE PATIENTS BY THE DEVELOPMENT OF GENE-INDEPENDENT THERAPIES**

Over the last two decades, proof-of-concept studies of corrective gene therapy have been established in many different animal models of inherited re-


tinopathies, strongly supporting further translation from animal models to human. However, evidence of the potency and efficacy of gene transfer to the retina at late stages of the disease is less robust. Early physiological alteration and loss of photoreceptor cells remain important factors limiting the therapeutic window for corrective gene therapy. In addition, with over 240 genes associated with retinal degeneration in humans, targeting individually each group of patients will likely not be possible. Mutation-independent therapies that allow treating entire families of retinal degenerative diseases would represent a more cost-effective approach (Fig. 1). Current strategies that utilize gene therapy and might be the more broadly applicable are detailed below.

#### **Preventing secondary cone death in retinitis pigmentosa**

Diseases that benefit the most from a mutation-independent approach are those associated with photoreceptor loss, in particular RP. The reason why this disease context is attractive for a mutation-independent approach is that, in the vast majority of cases, the genetic defects are specific to rod photoreceptors; however, cones die as well. It is this secondary loss of cones that causes the severe visual disabilities and complete blindness in humans. Consequently, gene therapies that prolong cone function and survival for such patients would maintain the temporal window of useful vision in a large number of patients.

A first point of intervention that is being developed inhibits rod cell death, which should prevent or delay the secondary loss of cones. Because the onset of cone death always follows the major rod death phase,<sup>62</sup> even a small delay in the death of rods could translate into a high preservation of cones.

One of the most straightforward ways to inhibit cell death is to inhibit the execution of cell death itself. Photoreceptor cell death has been found under various stress conditions to be predominantly apoptotic<sup>110</sup> and executed by caspases.<sup>111</sup> The X-linked inhibitor of apoptosis (XIAP) is one of a series of proteins that can inhibit cell death by binding to the two executioner caspases 3 and 7 and to the initiator caspase 9.<sup>112,113</sup> Misexpression of XIAP by adenoviral- or AAV2-mediated gene transfer has been shown to delay ganglion cell death in various rodent models of optic nerve injury<sup>114–116</sup> and protect neurons from retinal ischemia<sup>117</sup> and from the mutagen N-nitroso-N-methylurea.<sup>118</sup> Most recently, subretinal injection of AAV5-CBA-hXIAP has been shown to prevent photoreceptor death in the P23H rat model of dominant retinitis pigmentosa for at



**Figure 1.** Gene therapy strategies for inherited retinal diseases. **(A)** Schematic of subretinal gene therapy delivery in human patients. **(B)** Possible therapeutic interventions during the progression of inherited retinal dystrophies. **(B, top)** Time course of photoreceptor and/or RPE cell death. Therapeutic strategies at each stage of the disease are indicated. At early stages of the disease, affected photoreceptors and/or RPE cells can be targeted using mutation-specific gene therapy (gene replacement or gene suppression therapy). Photoreceptor death can also be prevented or delayed by mutation-independent strategies, for example, the delivery of growth factors. Therapies initiated at a more advanced stage of the disease, when most of the primary affected cells are gone, aim at targeting the common mechanism of secondary cone death. This can be done by the delivery of antioxidant genes (in cones or in other cells of the retina if the encoded factor is secreted, e.g., RdCVF) or by the delivery of genes that boost cell metabolism. After cones have lost their outer segments and become unresponsive to light stimulation, they can be reactivated by the introduction of NpHR. Nonphotoreceptor cells (bipolar and ganglion cells) can be reactivated by the introduction of various optogenes. **(B, bottom)** Images of retinal sections of the retinal degeneration 1 (*rd1*) mouse model of retinitis pigmentosa. Arrows indicate the target cells at each stage of photoreceptor death. GCL, ganglion cell layer; INL, inner nuclear layer; IS, inner segments; ONL, outer nuclear layer; OS, outer segments; RPE, retinal pigmented epithelium.

least 30 weeks<sup>119</sup> and confer photoreceptor protection 2 months after sodium hyaluronate-induced retinal detachment.<sup>120</sup>

Inhibition of photoreceptor degeneration using growth factors has also been tested. The ciliary neurotrophic factor (CNTF) is the most studied. CNTF has been shown to delay ganglion cell death through direct and AAV delivery<sup>121–125</sup> and to delay photoreceptor death in various animal models of RP.<sup>126–129</sup> A recent study in *Rho*<sup>-/-</sup> mice showed that intraocular delivery of AAV2-hCNTF can confer long-term rod and cone photoreceptors protection and significantly delay vision loss even when rod death is well advanced.<sup>130</sup> However, the mode of action of CNTF is not fully understood, and even though CNTF is protective in animals with photoreceptor degeneration, high doses are associated with an acute deconstruction of photoreceptor outer segments<sup>131,132</sup> and gene expression changes that are similar to those seen in light-induced photoreceptor plasticity.<sup>133</sup> Interestingly, the temporary deconstruction of photoreceptor outer segments appears to enhance gene transfer by AAV5 in a canine model of CNBG3-ACMH, making CNTF a potential therapeutic candidate to pretreat the eye before receiving the actual gene therapy.<sup>29</sup> Clinical evaluation of CNTF for the treatment of retinal degeneration has been already conducted using encapsulated cell implants in patients with advanced stages of RP (NCT00063765). In this phase 1 study, 10 patients received the implants in one eye for 6 months without major side effects.<sup>134</sup> Subsequently, three phase 2 clinical trials for early RP (NCT00447980, 68 patients), late RP (NCT00447993, 65 patients), and geographic atrophy (NCT00447954, 39 patients) were initiated. Data at 2 years post-implantation demonstrated that expression of CNTF was maintained over 24 months, with no serious adverse effects.<sup>135,136</sup> In patients with early-stage RP, no protection of rods has been demonstrated. Macular cone photoreceptors remained stable over 12–35 months, when sham-treated eyes experience a 9–24% decrease in cone number; however, this protective effect was not associated with detectable changes in visual acuity and ERG responses.<sup>137</sup> Determining final outcomes of the therapy on cone survival will require a longer follow-up because of the slow progressive nature of RP. If successful, AAV-mediated CNTF gene therapies may follow.

Besides the classical neuroprotective factors, both AAV-mediated delivery of erythropoietin (EPO), a cytokine that is upregulated during hypoxia, and AAV-mediated delivery of proinsulin have been shown to have neuroprotective proper-

ties in several models of RP.<sup>138–140</sup> Interestingly, however, only AAV-mediated systemic administration of EPO but not intraocular administration appears to be protective.<sup>141</sup> To translate this into a human gene therapy, EPO derivatives have been used that were protective after subretinal delivery by AAV in various models of retinal degeneration.<sup>139</sup> However, how applicable EPO derivatives and proinsulin are for a human therapy remains to be determined.

A second point of intervention targets cones directly to protect them from degeneration once the majority of rods have died. This requires an understanding of the mechanism of cone death, for which several models have been proposed. Some of these models seem to converge around energy availability and the associated redox potential (for review see ref.<sup>142</sup>).

Originally, it was believed that rods produce a trophic factor that is required for cone survival simply because cone death always follows rod death, regardless of the circumstances that lead to rod death. Identification of such factor would inevitably provide a unifying therapy for all forms of RP that are caused by mutations in rod-specific genes. In 2004, the rod-derived cone viability factor (RdCVF),<sup>143</sup> encoded by the nucleoredoxin-like 1 (*NXNL1*) gene, was identified and since has moved from the proof-of-concept as an injectable trophic factor that can delay cone death<sup>143,144</sup> into an AAV-mediated gene therapeutic approach.<sup>145</sup> RdCVF is a thioredoxin-like protein with two isoforms<sup>146</sup> and was initially believed to reduce oxidative stress, a disease condition that accompanies the degeneration of rods and cones because of the massive loss of rods.<sup>142</sup> Meanwhile, the protein has been shown to indirectly interact with the glucose transporter-1, thereby promoting glucose uptake in cones during the period of cone degeneration.<sup>147</sup> This finding is in line with one of the proposed mechanisms of secondary cone death, which postulates that cones are nutrient deprived; in particular, they are short of glucose.<sup>62,148</sup> Systemic intravenous delivery to *rd10* mice of AAV92YF-CAG-*Nxn11* at PN1 resulted in improved cone-mediated ERG responses and cone survival at 1 month of age. In the same mouse model, intravitreal injection at PN14 of a novel AAV2 variant with increased photoreceptor cells tropism, 7m8-CAG-*Nxn11*, also resulted in good preservation of cone function and survival 1 month posttreatment.<sup>145</sup> Clinical trials will undoubtedly determine if RdCVF becomes a therapeutic agent with broad applicability in humans.

The realization that oxidative stress is a contributing factor to secondary cone death<sup>149</sup> led to

extensive research in antioxidant therapies, many of which have shown promising results in mouse.<sup>150,151</sup> In humans, various combinations of vitamins and omega-3 fatty acids have been tested, showing a slight effect in delaying the disease progression.<sup>152–154</sup> However, the problem with orally supplemented antioxidants is that they may never reach critical concentrations in the tissue of interest. This circumstance, as well as the finding that antioxidant enzymes may need to be present in the right combination<sup>155</sup> directly in sick cones to reduce oxidative stress,<sup>156</sup> led to the first AAV-mediated approaches in which various enzymes and transcription factors that regulate the expression of detoxifying enzymes were tested.<sup>155,157</sup> The most promising of these candidates, nuclear factor erythroid-derived 2 like 2 (NRF2), showed a remarkable delay in cone death in two mouse models of RP, opening the door for a new mutation-independent approach for secondary cone death.

By investigating the molecular mechanisms of secondary cone death, we recently revealed that cones are nutrient deprived in RP,<sup>62</sup> a finding that is in line with the increase in oxidative stress seen in cones during degeneration,<sup>149</sup> since lack of glucose reduces the redox potential of a cell.<sup>142</sup> Because rods account for over 95% of all photoreceptors, we proposed that once a critical threshold of rod death is breached, cone death initiates a cell autonomous event caused by the collapse of the relatively few cone–RPE interactions reducing nutrient flow to cones.<sup>142</sup> This model explains why cones may survive for extended periods in the cone-rich central retina of patients and large animal models of RP, despite the total loss of rods, and why cone-specific diseases do not lead to rod degeneration. Genetic hyperactivation of a key kinase that promotes cell metabolism, the kinase mechanistic target of rapamycin complex 1 (mTORC1), has led to a remarkable delay of cone death for at least 8 months of age in two mouse models of RP.<sup>148</sup> The data represent the most profound and long-lasting effect of cone survival seen thus far, strongly suggesting that boosting cell metabolism in cones is a viable strategy to prolong cone survival in RP.<sup>158</sup> A targeted gene therapy approach that augments mTORC1 target genes in cones is currently under development. Interestingly, extending cell autonomous cone survival may go beyond RP and hold promise for many other degenerative diseases, such as age-related macular degeneration (AMD).<sup>159</sup> In these diseases, once a critical number of photoreceptors have died in a specific region, the remaining healthy photoreceptors in that region may be affected by reduced nutrient flow, as are cones in RP. In addition, the

degeneration of rescued photoreceptors observed in RPE65-treated patients may involve a similar mechanism of cone death to the one in RP.<sup>28</sup>

### Restoration of visual sensitivity by optogenetics

Restoration of vision through optogenetics provides an interesting strategy to restore visual perception in blind retinas (see reviews<sup>160,161</sup>). This therapeutic approach relies on delivering a gene encoding a light-sensitive channel protein (microbial opsins, endogenous opsins, or synthetic light-sensitive ion channels) to either reactivate dormant cones or activate other retinal neurons at late stages of degeneration.

Opsin-like microbial proteins have the advantage to reversibly isomerize a vitamin A derivative as the chromophore.<sup>162</sup> In response to light, microbial opsins can thus independently induce changes in the membrane potential of a cell allowing the electrical signal to propagate to the brain. Channelrhodopsin (CHR2) was the first microbial opsin to be used in the retina. CHR2 pumps cations upon excitation by light and produces excitatory currents. AAV-mediated expression of CHR2 at the level of ON bipolar cells<sup>163–167</sup> and ganglion cells<sup>168</sup> in mouse models of RP successfully restored light-evoked potentials in treated retinas. While transduction of ganglion and bipolar cells is still difficult in large animal models, novel vectors and promoters with better affinity for the inner retina have been recently reported.<sup>166,169,170</sup> Moreover, good preservation of these cells has been demonstrated in patients with advanced RP and LCA.<sup>171</sup> Another approach is to reactivate nonfunctional “dormant” cone photoreceptors using halorhodopsins, which produce inhibitory currents.<sup>172</sup> This strategy may restore visual processing in all layers of the retina. However, it remains to be determined who is an appropriate patient for this procedure as many RP patients keep functional cones for decades and the remodeling of the retinal network upon loss of photoreceptors could make this approach challenging in humans.<sup>173</sup> In addition, long-term effects of membrane potential depolarization/repolarization on nutrient-deprived cones remain unexplored.

Despite these advances, one problem microbial optogenetics faces is the light sensitivity and dynamic range of these various channels. Recently, a new generation of synthetic light-acting channels has been described that use azobenzene-based light switches fused to ion channels or receptors.<sup>174–176</sup> One such approach remodeled the ionotropic glutamate receptor<sup>177</sup> restoring visual function at the level of ganglion cells<sup>178</sup> and bipolar cells.<sup>179</sup> Initially,

optogenetics was introduced in 2002 by simultaneous coexpression of the *Drosophila* rhodopsin, arrestin-2, and the alpha subunit of the G-protein to activate culture hippocampal neurons by light.<sup>180</sup> The approach never gained much favorability over the microbial opsins because it requires a triple transduction, something that remains difficult to achieve in the retina. However, the idea of using endogenous G-protein-coupled receptors (GPCR) such as melanopsin,<sup>181</sup> which is involved in circadian rhythm and the pupillary reflex<sup>182</sup> and is naturally expressed in a subset of ganglion cells, has been revisited on the premise that other cells may express some of the components needed to activate the cascade with a GPCR. Recent experiments have shown that, when rhodopsin is delivered to bipolar cells, light sensitivity is restored by 2–3 orders of magnitude lower than what can be achieved by the microbial channels, making this a feasible approach to restore vision in late stages of RP.<sup>183</sup>

#### **Antiangiogenic gene therapy for acquired retinopathies**

The most successful mutation-independent approach thus far has been applied to two diseases, neither of which is caused by a specific mutation but rather by a combination of genetic and environmental risk factors. Wet AMD<sup>184</sup> and diabetic retinopathy (DR)<sup>185</sup> are both characterized by formation of new blood vessels of the choroidal and retinal vasculature, respectively. The subsequent leakage of fluid from these newly formed blood vessels into the retinal proper is what generally causes massive neuronal loss. The realization that vascular endothelial growth factor (VEGF) is one of the key culprits in promoting the disease pathology has led to the development of anti-VEGF therapies with first-generation therapies utilizing anti-VEGF antibodies (Lucentis [approved in 2006 for wet AMD] and Avastin [Genetech], Avastin being the parent molecule of Lucentis) that are administered intravitreally. However, the therapy requires continuous administration on an interval of 4–8 weeks, is costly, is associated with potential side effects from the repeated injections, and is a burden for patients because of the many clinical visits. Hence, gene therapeutic approaches have been established to either target VEGF itself, by viral-mediated overexpression of the soluble ligand binding part of the vascular endothelial growth factor receptor 1 gene (*FLT1*),<sup>186–192</sup> or by inhibiting its action through use of angiogenesis inhibitors such as pigment epithelium-derived factor (PEDF)<sup>193–195</sup> or a combination of two angiostatic factors, angiostatin and endostatin.<sup>196</sup> All three gene therapy ap-

proaches have since moved to the clinic. Because of its vast experimental evidence in animal models<sup>186–192</sup> and preclinical trials,<sup>197</sup> expression of *sFLT1* by AAV2-mediated gene transfer is the most advanced approach with two ongoing clinical trials. Avalanche-Lions Institute is evaluating an alternatively spliced form of *FLT1*, using the subretinal route (NCT01494805). Published results at 1 year posttreatment indicate that 4 out of 6 patients did not require any rescue injections with anti-VEGF antibodies.<sup>198</sup> Genzyme uses intravitreal delivery (NCT01024998). In parallel, Oxford Biomedica is evaluating the efficacy of simultaneous expression of angiostatin and endostatin by subretinal injection of EIAV-LV (RetinoStat, NCT01301443).

#### **CONCLUSION AND FUTURE PROSPECTS**

Despite existing limitations and questions that still need to be addressed and resolved, gene therapy for ocular diseases continues to show great promise for the future. Current clinical studies of gene replacement therapy, although not all are convincing in terms of longevity of therapeutic effects, have demonstrated a good safety record. They provide evidence that it is possible to obtain clinically relevant visual improvements after gene replacement. With the recent improvements in AAV vectors and the increase in therapeutic benefits observed in preclinical studies, new clinical trials should translate into even more encouraging results in patients. During the next decade, a better understanding of the mechanisms that commonly limit the efficacy of gene replacement therapy over the long-term will be a key aspect that needs to be solved to move forward with retinal gene replacement therapies. Simultaneously, lessons learned from the biology of retinopathies should enable the development of viable therapeutic strategies to prolong vision independently of the disease-causing mutations, further encouraging transfer to clinical trials. With the advent of genome editing technology for gene and cell therapy, future studies should also be aimed at safe and long-term AAV-mediated delivery of genome-modifying components to the target cell types in the eye, as well as other organs.

#### **ACKNOWLEDGMENTS**

The following funding support is gratefully acknowledged: the “Information Recherche Retinite Pigmentaire” research grant, the Fulbright/Fondation Mohanan Fellowship, the Fondation de France/Fondation Berthe Fouassier Fellowship, and the “Association Francaise contre les Myopathies” Fellowship. Work in the H.K. lab is sup-

ported by grants from the National Institutes of Health (EY022372), University of Massachusetts Center for Clinical and Translational Sciences (UMCCTS), Massachusetts Lions Eye Research Fund, and Foundation Fighting Blindness (FFB). Work in the C.P. lab is supported by the NIH

(EY023570) and the Massachusetts Lions Eye Research Fund.

## AUTHOR DISCLOSURE

No competing financial interests exist.

## REFERENCES

- Arshavsky VY, Lamb TD, Pugh EN Jr. G proteins and phototransduction. *Annu Rev Physiol* 2002; 64:153–187.
- Fishman GA, Jacobson SG, Alexander KR, et al. Outcome measures and their application in clinical trials for retinal degenerative diseases: Outline, review, and perspective. *Retina* 2005; 25:772–777.
- Moiseyev G, Chen Y, Takahashi Y, et al. RPE65 is the isomerohydrolase in the retinoid visual cycle. *Proc Natl Acad Sci U S A* 2005;102: 12413–12418.
- Redmond TM, Poliakov E, Yu S, et al. Mutation of key residues of RPE65 abolishes its enzymatic role as isomerohydrolase in the visual cycle. *Proc Natl Acad Sci U S A* 2005;102:13658–13663.
- Zhang T, Enemchukwu NO, Jones A, et al. Genetic deletion of S-opsin prevents rapid cone degeneration in a mouse model of Leber congenital amaurosis. *Hum Mol Genet* 2015;24: 1755–1763.
- Cideciyan AV. Leber congenital amaurosis due to RPE65 mutations and its treatment with gene therapy. *Prog Retin Eye Res* 2010;29:398–427.
- Acland GM, Aguirre GD, Ray J, et al. Gene therapy restores vision in a canine model of childhood blindness. *Nat Genet* 2001;28:92–95.
- Narfstrom K, Katz ML, Bragadottir R, et al. Functional and structural recovery of the retina after gene therapy in the RPE65 null mutation dog. *Invest Ophthalmol Vis Sci* 2003;44:1663–1672.
- Narfstrom K, Katz ML, Ford M, et al. *In vivo* gene therapy in young and adult RPE65<sup>-/-</sup> dogs produces long-term visual improvement. *J Hered* 2003;94:31–37.
- Acland GM, Aguirre GD, Bennett J, et al. Long-term restoration of rod and cone vision by single dose rAAV-mediated gene transfer to the retina in a canine model of childhood blindness. *Mol Ther* 2005;12:1072–1082.
- Narfstrom K, Vaegan, Katz M, et al. Assessment of structure and function over a 3-year period after gene transfer in RPE65<sup>-/-</sup> dogs. *Doc Ophthalmol* 2005;111:39–48.
- Le Meur G, Stieger K, Smith AJ, et al. Restoration of vision in RPE65-deficient Briard dogs using an AAV serotype 4 vector that specifically targets the retinal pigmented epithelium. *Gene Ther* 2007;14:292–303.
- Bennicelli J, Wright JF, Komaromy A, et al. Reversal of blindness in animal models of leber congenital amaurosis using optimized AAV2-mediated gene transfer. *Mol Ther* 2008;16: 458–465.
- Annear MJ, Mowat FM, Bartoe JT, et al. Successful gene therapy in older Rpe65-deficient dogs following subretinal injection of an adeno-associated vector expressing RPE65. *Hum Gene Ther* 2013;24:883–893.
- Mowat FM, Breuwer AR, Bartoe JT, et al. RPE65 gene therapy slows cone loss in Rpe65-deficient dogs. *Gene Ther* 2013;20:545–555.
- Cideciyan AV, Jacobson SG, Beltran WA, et al. Human retinal gene therapy for Leber congenital amaurosis shows advancing retinal degeneration despite enduring visual improvement. *Proc Natl Acad Sci U S A* 2013;110:E517–E525.
- Bainbridge JW, Mehat MS, Sundaram V, et al. Long-term effect of gene therapy on Leber's congenital amaurosis. *N Engl J Med* 2015;372: 1887–1897.
- Annear MJ, Bartoe JT, Barker SE, et al. Gene therapy in the second eye of RPE65-deficient dogs improves retinal function. *Gene Ther* 2011; 18:53–61.
- Maguire AM, Simonelli F, Pierce EA, et al. Safety and efficacy of gene transfer for Leber's congenital amaurosis. *N Engl J Med* 2008;358: 2240–2248.
- Bainbridge JW, Smith AJ, Barker SS, et al. Effect of gene therapy on visual function in Leber's congenital amaurosis. *N Engl J Med* 2008;358: 2231–2239.
- Hauswirth WW, Aleman TS, Kaushal S, et al. Treatment of leber congenital amaurosis due to RPE65 mutations by ocular subretinal injection of adeno-associated virus gene vector: Short-term results of a phase I trial. *Hum Gene Ther* 2008; 19:979–990.
- Jacobson SG, Cideciyan AV, Roman AJ, et al. Improvement and decline in vision with gene therapy in childhood blindness. *N Engl J Med* 2015;372:1920–1926.
- Pierce EA, Bennett J. The status of RPE65 gene therapy trials: Safety and efficacy. *Cold Spring Harb Perspect Med* 2015;5:a017285.
- Weleber RG, Pennesi ME, Wilson DJ, et al. Results at 2 years after gene therapy for RPE65-deficient Leber congenital amaurosis and severe early-childhood-onset retinal dystrophy. *Ophthalmology* 2016. [Epub ahead of print]
- Bennett J, Ashtari M, Wellman J, et al. AAV2 gene therapy readministration in three adults with congenital blindness. *Sci Transl Med* 2012; 4:120ra115.
- Schimmer J, Breazzano S. Investor outlook: Significance of the positive LCA2 gene therapy phase III results. *Hum Gene Ther Clin Dev* 2015; 26:208–210.
- Cideciyan AV, Aleman TS, Boye SL, et al. Human gene therapy for RPE65 isomerase deficiency activates the retinoid cycle of vision but with slow rod kinetics. *Proc Natl Acad Sci U S A* 2008;105:15112–15117.
- Cepko CL, Vandenbergh LH. Retinal gene therapy coming of age. *Hum Gene Ther* 2013;24: 242–244.
- Komaromy AM, Rowlan JS, Corr AT, et al. Transient photoreceptor deconstruction by CNTF enhances rAAV-mediated cone functional rescue in late stage CNGB3-achromatopsia. *Mol Ther* 2013;21:1131–1141.
- Ghazi NG, Abboud EB, Nowilaty SR, et al. Treatment of retinitis pigmentosa due to MERTK mutations by ocular subretinal injection of adeno-associated virus gene vector: Results of a phase I trial. *Hum Genet* 2016;135:327–343.
- MacLaren RE, Groppe M, Barnard AR, et al. Retinal gene therapy in patients with choroideremia: Initial findings from a phase 1/2 clinical trial. *Lancet* 2014;383:1129–1137.
- Edwards TL, Jolly JK, Groppe M, et al. Visual acuity after retinal gene therapy for choroideremia. *N Engl J Med* 2016;374:1996–1998.
- Syed N, Smith JE, John SK, et al. Evaluation of retinal photoreceptors and pigment epithelium in a female carrier of choroideremia. *Ophthalmology* 2001;108:711–720.
- Tolmachova T, Anders R, Abrink M, et al. Independent degeneration of photoreceptors and retinal pigment epithelium in conditional knockout mouse models of choroideremia. *J Clin Invest* 2006;116:386–394.
- Lai Y, Yue Y, Duan D. Evidence for the failure of adeno-associated virus serotype 5 to package a viral genome > or = 8.2 kb. *Mol Ther* 2010;18: 75–79.

36. Kong J, Kim SR, Binley K, et al. Correction of the disease phenotype in the mouse model of Stargardt disease by lentiviral gene therapy. *Gene Ther* 2008;15:1311–1320.
37. Hasson T, Heintzelman MB, Santos-Sacchi J, et al. Expression in cochlea and retina of myosin VIIa, the gene product defective in Usher syndrome type 1B. *Proc Natl Acad Sci U S A* 1995;92:9815–9819.
38. Zallocchi M, Binley K, Lad Y, et al. ElAV-based retinal gene therapy in the shaker1 mouse model for usher syndrome type 1B: Development of UshStat. *PLoS One* 2014;9:e94272.
39. Hashimoto T, Gibbs D, Lillo C, et al. Lentiviral gene replacement therapy of retinas in a mouse model for Usher syndrome type 1B. *Gene Ther* 2007;14:584–594.
40. Gruter O, Kostic C, Crippa SV, et al. Lentiviral vector-mediated gene transfer in adult mouse photoreceptors is impaired by the presence of a physical barrier. *Gene Ther* 2005;12:942–947.
41. Vandenberghe LH, Auricchio A. Novel adeno-associated viral vectors for retinal gene therapy. *Gene Ther* 2012;19:162–168.
42. Allocca M, Mussolino C, Garcia-Hoyos M, et al. Novel adeno-associated virus serotypes efficiently transduce murine photoreceptors. *J Virol* 2007;81:11372–11380.
43. Vandenberghe LH, Bell P, Maguire AM, et al. Dosage thresholds for AAV2 and AAV8 photoreceptor gene therapy in monkey. *Sci Transl Med* 2011;3:88ra54.
44. Surace EM, Auricchio A. Versatility of AAV vectors for retinal gene transfer. *Vision Res* 2008;48:353–359.
45. Alexander JJ, Umino Y, Everhart D, et al. Restoration of cone vision in a mouse model of achromatopsia. *Nat Med* 2007;13:685–687.
46. Michalakos S, Muhlfriedel R, Tanimoto N, et al. Restoration of cone vision in the CNGA3<sup>-/-</sup> mouse model of congenital complete lack of cone photoreceptor function. *Mol Ther* 2010;18:2057–2063.
47. Pang JJ, Deng WT, Dai X, et al. AAV-mediated cone rescue in a naturally occurring mouse model of CNGA3-achromatopsia. *PLoS One* 2012;7:e35250.
48. Carvalho LS, Xu J, Pearson RA, et al. Long-term and age-dependent restoration of visual function in a mouse model of CNGB3-associated achromatopsia following gene therapy. *Hum Mol Genet* 2011;20:3161–3175.
49. Komaromy AM, Alexander JJ, Rowlan JS, et al. Gene therapy rescues cone function in congenital achromatopsia. *Hum Mol Genet* 2010;19:2581–2593.
50. Banin E, Gootwine E, Obolensky A, et al. Gene augmentation therapy restores retinal function and visual behavior in a sheep model of CNGA3 achromatopsia. *Mol Ther* 2015;23:1423–1433.
51. Marangoni D, Vijayasarathy C, Bush RA, et al. Intravitreal ciliary neurotrophic factor transiently improves cone-mediated function in a CNGB3<sup>-/-</sup> mouse model of achromatopsia. *Invest Ophthalmol Vis Sci* 2015;56:6810–6822.
52. Langlo C, Dubis A, Michaelides M, et al. CNGB3-achromatopsia clinical trial with CNTF: Diminished rod pathway responses with no evidence of improvement in cone function. *Invest Ophthalmol Vis Sci* 2015;56:1505.
53. Ye GJ, Budzynski E, Sonnentag P, et al. Cone-specific promoters for gene therapy of achromatopsia and other retinal diseases. *Hum Gene Ther* 2016;27:72–82.
54. Weber BH, Schrewe H, Molday LL, et al. Inactivation of the murine X-linked juvenile retinoschisis gene, Rs1h, suggests a role of retinoschisin in retinal cell layer organization and synaptic structure. *Proc Natl Acad Sci U S A* 2002;99:6222–6227.
55. Min SH, Molday LL, Seeliger MW, et al. Prolonged recovery of retinal structure/function after gene therapy in an Rs1h-deficient mouse model of x-linked juvenile retinoschisis. *Mol Ther* 2005;12:644–651.
56. Zeng Y, Takada Y, Kjellstrom S, et al. RS-1 gene delivery to an adult Rs1h knockout mouse model restores ERG b-wave with reversal of the electronegative waveform of X-linked retinoschisis. *Invest Ophthalmol Vis Sci* 2004;45:3279–3285.
57. Park TK, Wu Z, Kjellstrom S, et al. Intravitreal delivery of AAV8 retinoschisin results in cell type-specific gene expression and retinal rescue in the Rs1-KO mouse. *Gene Ther* 2009;16:916–926.
58. Ou J, Vijayasarathy C, Ziccardi L, et al. Synaptic pathology and therapeutic repair in adult retinoschisis mouse by AAV-RS1 transfer. *J Clin Invest* 2015;125:2891–2903.
59. Dalkara D, Byrne LC, Klimczak RR, et al. *In vivo*-directed evolution of a new adeno-associated virus for therapeutic outer retinal gene delivery from the vitreous. *Sci Transl Med* 2013;5:189ra176.
60. Janssen A, Min SH, Molday LL, et al. Effect of late-stage therapy on disease progression in AAV-mediated rescue of photoreceptor cells in the retinoschisin-deficient mouse. *Mol Ther* 2008;16:1010–1017.
61. den Hollander AI, Black A, Bennett J, et al. Lighting a candle in the dark: Advances in genetics and gene therapy of recessive retinal dystrophies. *J Clin Invest* 2010;120:3042–3053.
62. Punzo C, Kornacker K, Cepko CL. Stimulation of the insulin/mTOR pathway delays cone death in a mouse model of retinitis pigmentosa. *Nat Neurosci* 2009;12:44–52.
63. Keeler CE. The inheritance of a retinal abnormality in white mice. *Proc Natl Acad Sci U S A* 1924;10:329–333.
64. Blanks JC, Adinolfi AM, Lolley RN. Photoreceptor degeneration and synaptogenesis in retinal-degenerative (rd) mice. *J Comp Neurol* 1974;156:95–106.
65. Carter-Dawson LD, LaVail MM, Sidman RL. Differential effect of the rd mutation on rods and cones in the mouse retina. *Invest Ophthalmol Vis Sci* 1978;17:489–498.
66. Jimenez AJ, Garcia-Fernandez JM, Gonzalez B, et al. The spatio-temporal pattern of photoreceptor degeneration in the aged rd/rd mouse retina. *Cell Tissue Res* 1996;284:193–202.
67. Chang B, Hawes NL, Pardue MT, et al. Two mouse retinal degenerations caused by missense mutations in the beta-subunit of rod cGMP phosphodiesterase gene. *Vis Res* 2007;47:624–633.
68. Gargini C, Terzibasi E, Mazzoni F, et al. Retinal organization in the retinal degeneration 10 (rd10) mutant mouse: A morphological and ERG study. *J Comp Neurol* 2007;500:222–238.
69. Davis RJ, Tosi J, Janisch KM, et al. Functional rescue of degenerating photoreceptors in mice homozygous for a hypomorphic cGMP phosphodiesterase 6 b allele (Pde6bH620Q). *Invest Ophthalmol Vis Sci* 2008;49:5067–5076.
70. Aguirre G, Farber D, Lolley R, et al. Retinal degeneration in the dog. III. Abnormal cyclic nucleotide metabolism in rod-cone dysplasia. *Exp Eye Res* 1982;35:625–642.
71. Aguirre GD, Rubin LF. Rod-cone dysplasia (progressive retinal atrophy) in Irish setters. *J Am Vet Med Assoc* 1975;166:157–164.
72. Goldstein O, Mezey JG, Schweitzer PA, et al. IQCB1 and PDE6B mutations cause similar early onset retinal degenerations in two closely related terrier dog breeds. *Invest Ophthalmol Vis Sci* 2013;54:7005–7019.
73. Bennett J, Tanabe T, Sun D, et al. Photoreceptor cell rescue in retinal degeneration (rd) mice by *in vivo* gene therapy. *Nat Med* 1996;2:649–654.
74. Kumar-Singh R, Farber DB. Encapsidated adenovirus mini-chromosome-mediated delivery of genes to the retina: Application to the rescue of photoreceptor degeneration. *Hum Mol Genet* 1998;7:1893–1900.
75. Takahashi M, Miyoshi H, Verma IM, et al. Rescue from photoreceptor degeneration in the rd mouse by human immunodeficiency virus vector-mediated gene transfer. *J Virol* 1999;73:7812–7816.
76. Jomary C, Vincent KA, Grist J, et al. Rescue of photoreceptor function by AAV-mediated gene transfer in a mouse model of inherited retinal degeneration. *Gene Ther* 1997;4:683–690.
77. Nishiguchi KM, Carvalho LS, Rizzi M, et al. Gene therapy restores vision in rd1 mice after removal of a confounding mutation in Gpr179. *Nat Commun* 2015;6:6006.
78. Pang JJ, Boye SL, Kumar A, et al. AAV-mediated gene therapy for retinal degeneration in the rd10 mouse containing a recessive PDEbeta mutation. *Invest Ophthalmol Vis Sci* 2008;49:4278–4283.
79. Allocca M, Manfredi A, Iodice C, et al. AAV-mediated gene replacement, either alone or in combination with physical and pharmacological agents, results in partial and transient protection

- from photoreceptor degeneration associated with betaPDE deficiency. *Invest Ophthalmol Vis Sci* 2011;52:5713–5719.
80. Mowat FM, Gornik KR, Dinculescu A, et al. Tyrosine capsid-mutant AAV vectors for gene delivery to the canine retina from a subretinal or intravitreal approach. *Gene Ther* 2014;21:96–105.
  81. Pang JJ, Dai X, Boye SE, et al. Long-term retinal function and structure rescue using capsid mutant AAV8 vector in the rd10 mouse, a model of recessive retinitis pigmentosa. *Mol Ther* 2011;19:234–242.
  82. Petit L, Lheriteau E, Weber M, et al. Restoration of vision in the pde6beta-deficient dog, a large animal model of rod-cone dystrophy. *Mol Ther* 2012;20:2019–2030.
  83. Pichard V, Provost N, Mendes-Madeira A, et al. AAV-mediated gene therapy halts retinal degeneration in PDE6beta-deficient dogs. *Mol Ther* 2016.
  84. Koch SF, Tsai YT, Duong JK, et al. Halting progressive neurodegeneration in advanced retinitis pigmentosa. *J Clin Invest* 2015;125:3704–3713.
  85. Wert KJ, Davis RJ, Sancho-Pelluz J, et al. Gene therapy provides long-term visual function in a pre-clinical model of retinitis pigmentosa. *Hum Mol Genet* 2013;22:558–567.
  86. Wert KJ, Sancho-Pelluz J, Tsang SH. Mid-stage intervention achieves similar efficacy as conventional early-stage treatment using gene therapy in a pre-clinical model of retinitis pigmentosa. *Hum Mol Genet* 2014;23:514–523.
  87. Michalakos S, Koch S, Sothilingam V, et al. Gene therapy restores vision and delays degeneration in the CNGB1(-/-) mouse model of retinitis pigmentosa. *Adv Exp Med Biol* 2014;801:733–739.
  88. Palfi A, Chadderton N, O'Reilly M, et al. Efficient gene delivery to photoreceptors using AAV2/rh10 and rescue of the Rho(-/-) mouse. *Mol Ther Methods Clin Dev* 2015;2:15016.
  89. Palfi A, Millington-Ward S, Chadderton N, et al. Adeno-associated virus-mediated rhodopsin replacement provides therapeutic benefit in mice with a targeted disruption of the rhodopsin gene. *Hum Gene Ther* 2010;21:311–323.
  90. Khani SC, Pawlyk BS, Bulgakov OV, et al. AAV-mediated expression targeting of rod and cone photoreceptors with a human rhodopsin kinase promoter. *Invest Ophthalmol Vis Sci* 2007;48:3954–3961.
  91. Manfredi A, Marrocco E, Puppo A, et al. Combined rod and cone transduction by adeno-associated virus 2/8. *Hum Gene Ther* 2013;24:982–992.
  92. Boye SE, Alexander JJ, Boye SL, et al. The human rhodopsin kinase promoter in an AAV5 vector confers rod- and cone-specific expression in the primate retina. *Hum Gene Ther* 2012;23:1101–1115.
  93. Lheriteau E, Petit L, Weber M, et al. Successful gene therapy in the RPGRIP1-deficient dog: A large model of cone-rod dystrophy. *Mol Ther* 2014;22:265–277.
  94. Haire SE, Pang J, Boye SL, et al. Light-driven cone arrestin translocation in cones of postnatal guanylate cyclase-1 knockout mouse retina treated with AAV-GC1. *Invest Ophthalmol Vis Sci* 2006;47:3745–3753.
  95. Boye SE, Boye SL, Pang J, et al. Functional and behavioral restoration of vision by gene therapy in the guanylate cyclase-1 (GC1) knockout mouse. *PLoS One* 2010;5:e11306.
  96. Mihelec M, Pearson RA, Robbie SJ, et al. Long-term preservation of cones and improvement in visual function following gene therapy in a mouse model of leber congenital amaurosis caused by guanylate cyclase-1 deficiency. *Hum Gene Ther* 2011;22:1179–1190.
  97. Boye SL, Conlon T, Erger K, et al. Long-term preservation of cone photoreceptors and restoration of cone function by gene therapy in the guanylate cyclase-1 knockout (GC1KO) mouse. *Invest Ophthalmol Vis Sci* 2011;52:7098–7108.
  98. Boye SL, Peterson JJ, Choudhury S, et al. Gene therapy fully restores vision to the all-cone Nrl(-/-) Gucy2e(-/-) mouse model of leber congenital amaurosis-1. *Hum Gene Ther* 2015;26:575–592.
  99. Boye SL, Peshenko IV, Huang WC, et al. AAV-mediated gene therapy in the guanylate cyclase (RetGC1/RetGC2) double knockout mouse model of Leber congenital amaurosis. *Hum Gene Ther* 2013;24:189–202.
  100. Pawlyk BS, Smith AJ, Buch PK, et al. Gene replacement therapy rescues photoreceptor degeneration in a murine model of Leber congenital amaurosis lacking RPGRIP. *Invest Ophthalmol Vis Sci* 2005;46:3039–3045.
  101. Pawlyk BS, Bulgakov OV, Liu X, et al. Replacement gene therapy with a human RPGRIP1 sequence slows photoreceptor degeneration in a murine model of Leber congenital amaurosis. *Hum Gene Ther* 2010;21:993–1004.
  102. Tan MH, Smith AJ, Pawlyk B, et al. Gene therapy for retinitis pigmentosa and Leber congenital amaurosis caused by defects in AIPL1: Effective rescue of mouse models of partial and complete Aipl1 deficiency using AAV2/2 and AAV2/8 vectors. *Hum Mol Genet* 2009;18:2099–2114.
  103. Sun X, Pawlyk B, Xu X, et al. Gene therapy with a promoter targeting both rods and cones rescues retinal degeneration caused by AIPL1 mutations. *Gene Ther* 2010;17:117–131.
  104. Ku CA, Chiodo VA, Boye SL, et al. Gene therapy using self-complementary Y733F capsid mutant AAV2/8 restores vision in a model of early onset Leber congenital amaurosis. *Hum Mol Genet* 2011;20:4569–4581.
  105. Beltran WA, Cideciyan AV, Iwabe S, et al. Successful arrest of photoreceptor and vision loss expands the therapeutic window of retinal gene therapy to later stages of disease. *Proc Natl Acad Sci U S A* 2015;112:E5844–5853.
  106. Beltran WA, Cideciyan AV, Lewin AS, et al. Gene therapy rescues photoreceptor blindness in dogs and paves the way for treating human X-linked retinitis pigmentosa. *Proc Natl Acad Sci U S A* 2012;109:2132–2137.
  107. Trapani I, Banfi S, Simonelli F, et al. Gene therapy of inherited retinal degenerations: Prospects and challenges. *Hum Gene Ther* 2015;26:193–200.
  108. Boye SE, Boye SL, Lewin AS, et al. A comprehensive review of retinal gene therapy. *Mol Ther* 2013;21:509–519.
  109. Carvalho LS, Vandenbergh LH. Promising and delivering gene therapies for vision loss. *Vision Res* 2015;111:124–133.
  110. Chinskey ND, Besirli CG, Zacks DN. Retinal cell death and current strategies in retinal neuroprotection. *Curr Opin Ophthalmol* 2014;25:228–233.
  111. Boatright KM, Renatus M, Scott FL, et al. A unified model for apical caspase activation. *Mol Cell* 2003;11:529–541.
  112. Roy N, Deveraux QL, Takahashi R, et al. The c-IAP-1 and c-IAP-2 proteins are direct inhibitors of specific caspases. *EMBO J* 1997;16:6914–6925.
  113. Deveraux QL, Takahashi R, Salvesen GS, et al. X-linked IAP is a direct inhibitor of cell-death proteases. *Nature* 1997;388:300–304.
  114. Kugler S, Straten G, Kreppel F, et al. The X-linked inhibitor of apoptosis (XIAP) prevents cell death in axotomized CNS neurons *in vivo*. *Cell Death Differ* 2000;7:815–824.
  115. Straten G, Schmeer C, Kretz A, et al. Potential synergistic protection of retinal ganglion cells from axotomy-induced apoptosis by adenoviral administration of glial cell line-derived neurotrophic factor and X-chromosome-linked inhibitor of apoptosis. *Neurobiol Dis* 2002;11:123–133.
  116. McKinnon SJ, Lehman DM, Tahzib NG, et al. Baculoviral IAP repeat-containing-4 protects optic nerve axons in a rat glaucoma model. *Mol Ther* 2002;5:780–787.
  117. Renwick J, Narang MA, Coupland SG, et al. XIAP-mediated neuroprotection in retinal ischemia. *Gene Ther* 2006;13:339–347.
  118. Petrin D, Baker A, Coupland SG, et al. Structural and functional protection of photoreceptors from MNU-induced retinal degeneration by the X-linked inhibitor of apoptosis. *Invest Ophthalmol Vis Sci* 2003;44:2757–2763.
  119. Leonard KC, Petrin D, Coupland SG, et al. XIAP protection of photoreceptors in animal models of retinitis pigmentosa. *PLoS One* 2007;2:e314.
  120. Zadro-Lamoureux LA, Zacks DN, Baker AN, et al. XIAP effects on retinal detachment-induced photoreceptor apoptosis [corrected]. *Invest Ophthalmol Vis Sci* 2009;50:1448–1453.
  121. Ji JZ, Elyaman W, Yip HK, et al. CNTF promotes survival of retinal ganglion cells after induction of ocular hypertension in rats: The possible involvement of STAT3 pathway. *Eur J Neurosci* 2004;19:265–272.



122. Leaver SG, Cui Q, Plant GW, et al. AAV-mediated expression of CNTF promotes long-term survival and regeneration of adult rat retinal ganglion cells. *Gene Ther* 2006;13:1328–1341.
123. MacLaren RE, Buch PK, Smith AJ, et al. CNTF gene transfer protects ganglion cells in rat retinae undergoing focal injury and branch vessel occlusion. *Exp Eye Res* 2006;83:1118–1127.
124. Maier K, Rau CR, Storch MK, et al. Ciliary neurotrophic factor protects retinal ganglion cells from secondary cell death during acute autoimmune optic neuritis in rats. *Brain Pathol* 2004;14:378–387.
125. Pease ME, Zack DJ, Berlinicke C, et al. Effect of CNTF on retinal ganglion cell survival in experimental glaucoma. *Invest Ophthalmol Vis Sci* 2009;50:2194–2200.
126. Tao W, Wen R, Goddard MB, et al. Encapsulated cell-based delivery of CNTF reduces photoreceptor degeneration in animal models of retinitis pigmentosa. *Invest Ophthalmol Vis Sci* 2002;43:3292–3298.
127. LaVail MM, Yasumura D, Matthes MT, et al. Protection of mouse photoreceptors by survival factors in retinal degenerations. *Invest Ophthalmol Vis Sci* 1998;39:592–602.
128. Chong NH, Alexander RA, Waters L, et al. Repeated injections of a ciliary neurotrophic factor analogue leading to long-term photoreceptor survival in hereditary retinal degeneration. *Invest Ophthalmol Vis Sci* 1999;40:1298–1305.
129. Liang FQ, Dejneka NS, Cohen DR, et al. AAV-mediated delivery of ciliary neurotrophic factor prolongs photoreceptor survival in the rhodopsin knockout mouse. *Mol Ther* 2001;3:241–248.
130. Lipinski DM, Barnard AR, Singh MS, et al. CNTF gene therapy confers lifelong neuroprotection in a mouse model of human retinitis pigmentosa. *Mol Ther* 2015;23:1308–1319.
131. Schlichtenbrede FC, MacNeil A, Bainbridge JW, et al. Intraocular gene delivery of ciliary neurotrophic factor results in significant loss of retinal function in normal mice and in the Prph2Rd2/Rd2 model of retinal degeneration. *Gene Ther* 2003;10:523–527.
132. Bok D, Yasumura D, Matthes MT, et al. Effects of adeno-associated virus-vectored ciliary neurotrophic factor on retinal structure and function in mice with a P216L rds/peripherin mutation. *Exp Eye Res* 2002;74:719–735.
133. Wen R, Song Y, Kjellstrom S, et al. Regulation of rod phototransduction machinery by ciliary neurotrophic factor. *J Neurosci* 2006;26:13523–13530.
134. Sieving PA, Caruso RC, Tao W, et al. Ciliary neurotrophic factor (CNTF) for human retinal degeneration: Phase I trial of CNTF delivered by encapsulated cell intraocular implants. *Proc Natl Acad Sci U S A* 2006;103:3896–3901.
135. Birch DG, Weleber RG, Duncan JL, et al. Randomized trial of ciliary neurotrophic factor delivered by encapsulated cell intraocular implants for retinitis pigmentosa. *Am J Ophthalmol* 2013;156:283–292 e281.
136. Zhang K, Hopkins JJ, Heier JS, et al. Ciliary neurotrophic factor delivered by encapsulated cell intraocular implants for treatment of geographic atrophy in age-related macular degeneration. *Proc Natl Acad Sci U S A* 2011;108:6241–6245.
137. Talcott KE, Ratnam K, Sundquist SM, et al. Longitudinal study of cone photoreceptors during retinal degeneration and in response to ciliary neurotrophic factor treatment. *Invest Ophthalmol Vis Sci* 2011;52:2219–2226.
138. Sullivan R, Rex TS. Systemic gene delivery protects the photoreceptors in the retinal degeneration slow mouse. *Neurochem Res* 2011;36:613–618.
139. Colella P, Iodice C, Di Vicino U, et al. Non-erythropoietic erythropoietin derivatives protect from light-induced and genetic photoreceptor degeneration. *Hum Mol Genet* 2011;20:2251–2262.
140. Fernandez-Sanchez L, Lax P, Isiegas C, et al. Proinsulin slows retinal degeneration and vision loss in the P23H rat model of retinitis pigmentosa. *Hum Gene Ther* 2012;23:1290–1300.
141. Rex TS, Allocca M, Domenici L, et al. Systemic but not intraocular Epo gene transfer protects the retina from light-and genetic-induced degeneration. *Mol Ther* 2004;10:855–861.
142. Punzo C, Xiong W, Cepko CL. Loss of daylight vision in retinal degeneration: Are oxidative stress and metabolic dysregulation to blame? *J Biol Chem* 2012;287:1642–1648.
143. Leveillard T, Mohand-Said S, Lorentz O, et al. Identification and characterization of rod-derived cone viability factor. *Nat Genet* 2004;36:755–759.
144. Yang Y, Mohand-Said S, Danan A, et al. Functional cone rescue by RdCVF protein in a dominant model of retinitis pigmentosa. *Mol Ther* 2009;17:787–795.
145. Byrne LC, Dalkara D, Luna G, et al. Viral-mediated RdCVF and RdCVFL expression protects cone and rod photoreceptors in retinal degeneration. *J Clin Invest* 2015;125:105–116.
146. Fridlich R, Delalande F, Jaillard C, et al. The thioredoxin-like protein rod-derived cone viability factor (RdCVFL) interacts with TAU and inhibits its phosphorylation in the retina. *Mol Cell Proteomics* 2009;8:1206–1218.
147. Ait-Ali N, Fridlich R, Millet-Puel G, et al. Rod-derived cone viability factor promotes cone survival by stimulating aerobic glycolysis. *Cell* 2015;161:817–832.
148. Venkatesh A, Ma S, Le YZ, et al. Activated mTORC1 promotes long-term cone survival in retinitis pigmentosa mice. *J Clin Invest* 2015;125:1446–1458.
149. Shen J, Yang X, Dong A, et al. Oxidative damage is a potential cause of cone cell death in retinitis pigmentosa. *J Cell Physiol* 2005;203:457–464.
150. Komeima K, Rogers BS, Campochiaro PA. Antioxidants slow photoreceptor cell death in mouse models of retinitis pigmentosa. *J Cell Physiol* 2007;213:809–815.
151. Komeima K, Rogers BS, Lu L, et al. Antioxidants reduce cone cell death in a model of retinitis pigmentosa. *Proc Natl Acad Sci U S A* 2006;103:11300–11305.
152. Berson EL, Rosner B, Sandberg MA, et al. omega-3 intake and visual acuity in patients with retinitis pigmentosa receiving vitamin A. *Arch Ophthalmol* 2012;130:707–711.
153. Berson EL, Rosner B, Sandberg MA, et al. Clinical trial of lutein in patients with retinitis pigmentosa receiving vitamin A. *Arch Ophthalmol* 2010;128:403–411.
154. Berson EL, Rosner B, Sandberg MA, et al. A randomized trial of vitamin A and vitamin E supplementation for retinitis pigmentosa. *Arch Ophthalmol* 1993;111:761–772.
155. Usui S, Oveson BC, Iwase T, et al. Overexpression of SOD in retina: Need for increase in H2O2-detoxifying enzyme in same cellular compartment. *Free Radic Biol Med* 2011.
156. Usui S, Komeima K, Lee SY, et al. Increased expression of catalase and superoxide dismutase 2 reduces cone cell death in retinitis pigmentosa. *Mol Ther* 2009;17:778–786.
157. Xiong W, MacColl Garfinkel AE, Li Y, et al. NRF2 promotes neuronal survival in neurodegeneration and acute nerve damage. *J Clin Invest* 2015;125:1433–1445.
158. Petit L, Punzo C. mTORC1 sustains vision in retinitis pigmentosa. *Oncotarget* 2015;6:16786–16787.
159. Zieger M, Punzo C. Improved cell metabolism prolongs photoreceptor survival upon retinal-pigmented epithelium loss in the sodium iodate induced model of geographic atrophy. *Oncotarget* 2016;7.
160. Busskamp V, Picaud S, Sahel JA, et al. Optogenetic therapy for retinitis pigmentosa. *Gene Ther* 2012;19:169–175.
161. Sahel JA, Roska B. Gene therapy for blindness. *Annu Rev Neurosci* 2013;36:467–488.
162. Deisseroth K. Optogenetics: 10 years of microbial opsins in neuroscience. *Nat Neurosci* 2015;18:1213–1225.
163. Lagali PS, Balya D, Awatramani GB, et al. Light-activated channels targeted to ON bipolar cells restore visual function in retinal degeneration. *Nat Neurosci* 2008;11:667–675.
164. Doroudchi MM, Greenberg KP, Liu J, et al. Virally delivered channelrhodopsin-2 safely and effectively restores visual function in multiple mouse models of blindness. *Mol Ther* 2011;19:1220–1229.
165. Bi A, Cui J, Ma YP, et al. Ectopic expression of a microbial-type rhodopsin restores visual responses in mice with photoreceptor degeneration. *Neuron* 2006;50:23–33.
166. Cronin T, Vandenberghe LH, Hantz P, et al. Efficient transduction and optogenetic stimulation of

- retinal bipolar cells by a synthetic adeno-associated virus capsid and promoter. *EMBO Mol Med* 2014;6:1175–1190.
167. Mace E, Caplette R, Marre O, et al. Targeting channelrhodopsin-2 to ON-bipolar cells with vitreally administered AAV Restores ON and OFF visual responses in blind mice. *Mol Ther* 2015; 23:7–16.
  168. Thyagarajan S, van Wyk M, Lehmann K, et al. Visual function in mice with photoreceptor degeneration and transgenic expression of channelrhodopsin 2 in ganglion cells. *J Neurosci* 2010; 30:8745–8758.
  169. Scalabrino ML, Boye SL, Fransen KM, et al. Intravitreal delivery of a novel AAV vector targets ON bipolar cells and restores visual function in a mouse model of complete congenital stationary night blindness. *Hum Mol Genet* 2015;24: 6229–6239.
  170. Lu Q, Ganjawala TH, Ivanova E, et al. AAV-mediated transduction and targeting of retinal bipolar cells with improved mGluR6 promoters in rodents and primates. *Gene Ther* 2016.
  171. Jacobson SG, Cideciyan AV, Ratnakaram R, et al. Gene therapy for leber congenital amaurosis caused by RPE65 mutations: Safety and efficacy in 15 children and adults followed up to 3 years. *Arch Ophthalmol* 2012;130:9–24.
  172. Busskamp V, Duebel J, Balya D, et al. Genetic reactivation of cone photoreceptors restores visual responses in retinitis pigmentosa. *Science* 2010;329:413–417.
  173. Jacobson SG, Sumaroka A, Luo X, et al. Retinal optogenetic therapies: Clinical criteria for candidacy. *Clin Genet* 2013;84:175–182.
  174. Fehrentz T, Schonberger M, Trauner D. Optochemical genetics. *Angew Chem Int Ed Engl* 2011; 50:12156–12182.
  175. Kramer RH, Fortin DL, Trauner D. New photochemical tools for controlling neuronal activity. *Curr Opin Neurobiol* 2009;19:544–552.
  176. Specht A, Bolze F, Omran Z, et al. Photochemical tools to study dynamic biological processes. *HFSP J* 2009;3:255–264.
  177. Volgraf M, Gorostiza P, Numano R, et al. Allosteric control of an ionotropic glutamate receptor with an optical switch. *Nat Chem Biol* 2006;2: 47–52.
  178. Caporale N, Kolstad KD, Lee T, et al. LiGluR restores visual responses in rodent models of inherited blindness. *Mol Ther* 2011;19:1212–1219.
  179. Gaub BM, Berry MH, Holt AE, et al. Restoration of visual function by expression of a light-gated mammalian ion channel in retinal ganglion cells or ON-bipolar cells. *Proc Natl Acad Sci U S A* 2014;111:E5574–5583.
  180. Zemelman BV, Lee GA, Ng M, et al. Selective photostimulation of genetically chARGed neurons. *Neuron* 2002;33:15–22.
  181. Lin B, Koizumi A, Tanaka N, et al. Restoration of visual function in retinal degeneration mice by ectopic expression of melanopsin. *Proc Natl Acad Sci U S A* 2008;105:16009–16014.
  182. Benarroch EE. The melanopsin system: Phototransduction, projections, functions, and clinical implications. *Neurology* 2011;76:1422–1427.
  183. Gaub BM, Berry MH, Holt AE, et al. Optogenetic vision restoration using rhodopsin for enhanced sensitivity. *Mol Ther* 2015;23:1562–1571.
  184. Yonekawa Y, Miller JW, Kim IK. Age-related macular degeneration: Advances in management and diagnosis. *J Clin Med* 2015;4:343–359.
  185. Park YG, Roh YJ. New Diagnostic and therapeutic approaches for preventing the progression of diabetic retinopathy. *J Diabetes Res* 2016; 2016:1753584.
  186. Kendall RL, Thomas KA. Inhibition of vascular endothelial cell growth factor activity by an endogenously encoded soluble receptor. *Proc Natl Acad Sci U S A* 1993;90:10705–10709.
  187. Lai YK, Shen WY, Brankov M, et al. Potential long-term inhibition of ocular neovascularisation by recombinant adeno-associated virus-mediated secretion gene therapy. *Gene Ther* 2002;9:804–813.
  188. Gehlbach P, Demetriades AM, Yamamoto S, et al. Periocular gene transfer of sFlt-1 suppresses ocular neovascularization and vascular endothelial growth factor-induced breakdown of the blood-retinal barrier. *Hum Gene Ther* 2003;14:129–141.
  189. Ideno J, Mizukami H, Kakehashi A, et al. Prevention of diabetic retinopathy by intraocular soluble flt-1 gene transfer in a spontaneously diabetic rat model. *Int J Mol Med* 2007;19:75–79.
  190. Igarashi T, Miyake K, Masuda I, et al. Adeno-associated vector (type 8)-mediated expression of soluble Flt-1 efficiently inhibits neovascularization in a murine choroidal neovascularization model. *Hum Gene Ther* 2010;21: 631–637.
  191. Rota R, Riccioni T, Zaccarini M, et al. Marked inhibition of retinal neovascularization in rats following soluble-flt-1 gene transfer. *J Gene Med* 2004;6:992–1002.
  192. Tuo J, Pang JJ, Cao X, et al. AAV5-mediated sFLT01 gene therapy arrests retinal lesions in Ccl2(-)/Cx3cr1(-) mice. *Neurobiol Aging* 2012; 33:433.e1–e10.
  193. Dawson DW, Volpert OV, Gillis P, et al. Pigment epithelium-derived factor: A potent inhibitor of angiogenesis. *Science* 1999;285:245–248.
  194. Auricchio A, Behling KC, Maguire AM, et al. Inhibition of retinal neovascularization by intraocular viral-mediated delivery of anti-angiogenic agents. *Mol Ther* 2002;6:490–494.
  195. Mori K, Duh E, Gehlbach P, et al. Pigment epithelium-derived factor inhibits retinal and choroidal neovascularization. *J Cell Physiol* 2001; 188:253–263.
  196. Scappaticci FA, Smith R, Pathak A, et al. Combination angiostatin and endostatin gene transfer induces synergistic antiangiogenic activity *in vitro* and antitumor efficacy in leukemia and solid tumors in mice. *Mol Ther* 2001;3:186–196.
  197. MacLachlan TK, Lukason M, Collins M, et al. Preclinical safety evaluation of AAV2-sFLT01- a gene therapy for age-related macular degeneration. *Mol Ther* 2011;19:326–334.
  198. Rakoczy EP, Lai CM, Magno AL, et al. Gene therapy with recombinant adeno-associated vectors for neovascular age-related macular degeneration: 1 year follow-up of a phase 1 randomised clinical trial. *Lancet* 2015;386:2395–2403.

Received for publication March 28, 2016;  
accepted after revision May 10, 2016.

Published online: May 13, 2016.