

6-1-2007

ED Diagnosis of Acute Coronary Syndromes: No Gender-Related Difference of 'Chest Discomfort'

Garreth C. Biegun

University of Massachusetts Medical School

Follow this and additional works at: <http://escholarship.umassmed.edu/ssp>

 Part of the [Cardiology Commons](#), [Emergency Medicine Commons](#), and the [Health Services Research Commons](#)

Repository Citation

Biegun, Garreth C., "ED Diagnosis of Acute Coronary Syndromes: No Gender-Related Difference of 'Chest Discomfort'" (2007).
University of Massachusetts Medical School. *Senior Scholars Program*. Paper 42.
<http://escholarship.umassmed.edu/ssp/42>

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in Senior Scholars Program by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.

ED Diagnosis of Acute Coronary Syndromes: No Gender-Related Difference in the Incidence of 'Chest Discomfort'

Garreth C Biegun, Craig L Longo MD, Karin Przyklenk PhD, Chad E Darling MD
Emergency Medicine, University of Massachusetts Medical School, Worcester MA



Introduction

- There is evidence for gender and diabetes-related differences in symptoms of ACS upon presentation to the ED: i.e., non-diabetic men typically report 'chest pain,' whereas women and diabetics may report atypical complaints.
- This may reflect differences in either ACS-related chest pain or differences in the perception of pain.

Objective

- Our aim was to compare the frequency of broadly defined 'chest associated discomfort' rather than 'chest pain' reported by men vs. women and diabetics vs. non-diabetics with MI.

Methods

- This prospective, ongoing study enrolls patients presenting to an urban academic medical center with the subsequent diagnosis of NSTEMI or STEMI.
- After admission, patients were interviewed using a focused, semi-structured format and queried as to the presence (yes/no), severity, and quality of chest discomfort—defined as any symptom referred to the thorax—upon ED presentation.
- Severity was scored on a scale of 1 to 10.
- Patients were excluded if they were unstable or otherwise unable to give a history.

	Males	Females
Total	47	34
Number with chest discomfort	44 (94%)	31 (91%)
Number without chest discomfort	3 (6%)	3 (9%)
Average severity (p=0.76 by t-test)	7.2 ± 2.4	7.4 ± 2.8

Figure 1: Data: males vs. females

	Diabetics	Non-diabetics
Total	21	60
Number with chest discomfort	18 (86%)	57 (95%)
Number without chest discomfort	3 (14%)	3 (5%)
Average severity (p=0.34 by t-test)	6.8 ± 2.9	7.4 ± 2.4

Figure 2: Data: diabetics vs. non-diabetics

Results

- Incidence of chest discomfort was 91% in women and 94% in men. (p=0.69 by Fischer's exact test)
- Incidence of chest discomfort was 86% in diabetics and 95% in non-diabetics. (p=0.18)

Number of patients reporting severity 1-10




Figure 3: Number of patients reporting each level of severity

Conclusion

- These preliminary results suggest that, while there may be gender or diabetes-related differences in the perception of 'chest pain', there is an equivalent incidence and severity of 'chest discomfort' in all groups.
- This reinforces the importance of pursuing broad complaints of chest discomfort in the ED.