

May 22nd, 4:30 PM - 6:00 PM

Disrupted Resting-State Functional Connectivity in Medication-free Women with Postpartum Depression

Kristina M. Deligiannidis

University of Massachusetts Medical School, kristina.deligiannidis@umassmemorial.org

Elif M. Sikoglu

University of Massachusetts Medical School, elif.sikoglu@umassmed.edu

Anthony J. Rothschild

University of Massachusetts Medical School, anthony.rothschild@umassmemorial.org

See next page for additional authors

Follow this and additional works at: http://escholarship.umassmed.edu/cts_retreat

 Part of the [Maternal and Child Health Commons](#), [Neuroscience and Neurobiology Commons](#),
and the [Psychiatry Commons](#)

Kristina M. Deligiannidis, Elif M. Sikoglu, Anthony J. Rothschild, and Constance M. Moore, "Disrupted Resting-State Functional Connectivity in Medication-free Women with Postpartum Depression" (May 22, 2012). *UMass Center for Clinical and Translational Science Research Retreat*. Paper 14.
http://escholarship.umassmed.edu/cts_retreat/2012/posters/14

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.

Presenter Information

Kristina M. Deligiannidis, Elif M. Sikoglu, Anthony J. Rothschild, and Constance M. Moore

Creative Commons License

This work is licensed under a [Creative Commons Attribution-Noncommercial-Share Alike 3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/).

DISRUPTED RESTING-STATE FUNCTIONAL CONNECTIVITY IN MEDICATION-FREE WOMEN WITH POSTPARTUM DEPRESSION

Kristina M. Deligiannidis, MD¹, Elif M. Sikoglu, PhD², Anthony J. Rothschild, MD^{1*}, Constance M. Moore, PhD^{2*}

¹Center for Psychopharmacologic Research & Treatment, University of Massachusetts Medical School and UMass Memorial Health Care, Worcester, MA; ²Center for Comparative Neuroimaging, University of Massachusetts Medical School, Worcester, MA; * *equal contribution*

Contact: kristina.deligiannidis@umassmemorial.org

Background: Women are at increased risk of developing depression in the postpartum, a time of gonadal steroid flux. Abnormalities of gonadal steroids have been identified in some depressed women at times of reproductive flux. Gonadal steroids modulate corticocortical and corticolimbic functional connectivity (FC) in healthy, non-puerperal subjects; however there are no published studies of FC in postpartum depression (PPD).

Methods: Healthy comparison (HCS) (n=9) and medication-free subjects with unipolar PPD (n=8) were scanned at 3-9 weeks postpartum (using 3T Philips MRI) while 'at rest' with eyes open. Data analysis was carried out using SPM-8 and Data Processing Assistant for Resting-State fMRI (DPARSF) to perform seed based resting-state functional connectivity (rs-FC) analysis. Seeds were placed at bilateral anterior cingulate (ACC) and dorsolateral prefrontal cortices (DLPFC), hippocampi (HIP) and amygdalae (AMYG). Correlation coefficients obtained from individual subject analysis were used in performing two-sample t-test to compare the two cohorts.

Results: Functional connectivity maps revealed a more distributed pattern of connectivity with each seed (i.e. ACC, DLPFX, HIP, and AMYG) for HCS than PPD. In PPD subjects as compared to HCS, there was attenuation of rs-FC between corticocortical and corticolimbic areas.

Conclusions: In the early postpartum period, rs-FC patterns are disrupted in women with PPD in brain regions important for cognition, affect and the stress response. Larger studies are necessary to elucidate the role of disrupted neural connections in the pathophysiology of PPD and the potential modulatory role of gonadal steroids in women.

Supported by Worcester Foundation for Biomedical Research