### **University of Massachusetts Medical School** eScholarship@UMMS

Research Retreat

UMass Center for Clinical and Translational Science 2012 UMass Center for Clinical and Translational Science Research Retreat

May 22nd, 4:30 PM - 6:00 PM

# Importance of clinically-refined medical and musculoskeletal co-mórbidities in registries that evaluate patient-reported outcomes following TKR

Patricia D. Franklin University of Massachusetts Medical School, patricia.franklin@umassmed.edu

Wenjun Li University of Massachusetts Medical School, wenjun.li@umassmed.edu

Carol Otis Arcadia University

See next page for additional authors

Follow this and additional works at: http://escholarship.umassmed.edu/cts retreat Part of the Community Health and Preventive Medicine Commons, Health Services Research Commons, and the Orthopedics Commons

Patricia D. Franklin, Wenjun Li, Carol Otis, Benjamin Snyder, Milagros C. Rosal, and David C. Ayers, "Importance of clinically-refined medical and musculoskeletal co-morbidities in registries that evaluate patient-reported outcomes following TKR" (May 22, 2012). UMass Center for Clinical and Translational Science Research Retreat. Paper 27. http://escholarship.umassmed.edu/cts\_retreat/2012/posters/27

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.

#### **Presenter Information**

Patricia D. Franklin, Wenjun Li, Carol Otis, Benjamin Snyder, Milagros C. Rosal, and David C. Ayers

#### **Creative Commons License**



This work is licensed under a Creative Commons Attribution-Noncommercial-Share Alike 3.0 License.

## IMPORTANCE OF CLINICALLY-REFINED MEDICAL AND MUSCULOSKELETAL CO-MORBIDITIES IN REGISTRIES THAT EVALUATE PATIENT-REPORTED OUTCOMES FOLLOWING TKR

Patricia D. Franklin<sup>1</sup>; Wenjun Li<sup>3</sup>, PhD; Carol Otis, PhD; Benjamin Snyder<sup>1</sup>, MD MS; Milagros Rosal<sup>3</sup>, PhD; David C. Ayers<sup>1</sup>, MD

UMass Medical School departments of: <sup>1</sup>Orthopedics and Physical Rehabilitation, <sup>2</sup>Preventive and Behavioral Medicine, <sup>3</sup>Arcadia University

Contact: Patricia D. Franklin, MD MBA MPH, Email: patricia.franklin@umassmed.edu

**BACKGROUND:** As national joint registries broaden their focus to include patient-reported outcomes, such as pain relief and functional gain, the role of confounding peri-operative complications and comorbidities must be considered. We hypothesized that emotional, medical, and musculoskeletal comorbidities influence post-surgical functional gain following primary total knee (TKR) replacement surgery.

**METHODS:** We performed secondary analyses of comprehensive data from 180 primary TKR patients to evaluate the association of age, sex, body mass index (BMI), pre-operative emotional health (SF36 MCS/metal component score and CES-D depression screen), medical comorbidities (modified Charlson index), and musculoskeletal comorbidities (pain in low back, hips, and knees) on change in pre-to-6 month post-TKR physical function (SF36 PCS/physical component score).

**RESULTS:** Patients were 68% female with mean age of 65 years, mean BMI of 32. Mean pre-TKR PCS was 32.0 (SE=0.65), mean MCS was 52.0 (SE= 0.46) and 32% reported pre-TKR CES-D score at the mild to moderate depression level. Musculoskeletal comorbidity was scored as percent of patients with moderate or severe pain; 12% had hip pain, 46% contra-lateral knee pain, 27% low back pain, and 12% foot pain. Six percent reported COPD, 7% cardiac conditions, and 3% renal conditions. A multivariate linear regression model showed BMI>30, lower pre-MCS, lower pre-PCS, moderate or severe low back pain, and higher Charlson co-morbidity score, to be significantly (p<0.05) associated with poorer 6 month post-TKR PCS (physical function).

**CONCLUSION:** Pre-operative BMI, emotional health, and medical and musculoskeletal co-morbidities are required to interpret pre-to-post-operative change in physical function. Further understanding of the role of these factors is critical before national registries can analyze and report valid comparisons of patient-reported outcomes.