

May 22nd, 4:30 PM - 6:00 PM

Molecular imaging of the bioeffects of β -amyloid and metal ions on live human neuroblastoma cells: internalization, subcellular localization and induction of ROS

Young Jiao

University of Massachusetts - Dartmouth

Yi Zhang

University of Massachusetts - Dartmouth

Maolin Guo

University of Massachusetts - Dartmouth, mguo@umassd.edu

Follow this and additional works at: http://escholarship.umassmed.edu/cts_retreat



Part of the [Biochemistry, Biophysics, and Structural Biology Commons](#)



This work is licensed under a [Creative Commons Attribution-Noncommercial-Share Alike 3.0 License](#).

Young Jiao, Yi Zhang, and Maolin Guo, "Molecular imaging of the bioeffects of β -amyloid and metal ions on live human neuroblastoma cells: internalization, subcellular localization and induction of ROS" (May 22, 2012). *UMass Center for Clinical and Translational Science Research Retreat*. Paper 33.

http://escholarship.umassmed.edu/cts_retreat/2012/posters/33

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.

MOLECULAR IMAGING OF THE BIOEFFECTS OF β -AMYLOID AND METAL IONS ON LIVE HUMAN NEUROBLASTOMA CELLS: INTERNALIZATION, SUBCELLULAR LOCALIZATION AND INDUCTION OF ROS

Yong Jiao, Yi Zhang, Maolin Guo

Department of Chemistry and Biochemistry, University of Massachusetts, Dartmouth, MA

Contact: Maolin Guo, mguo@umassd.edu

Alzheimer's disease (AD) is a progressive neurodegenerative disease characterized by the deposition of extracellular amyloid- β (A β) plaques that are rich in metal ions such as zinc, copper and iron. The neurotoxic role of A β has been well established but the mechanism of action is still poorly understood. Recent in vitro evidence suggest that A β can interact with metal ions such as Zn(II), Cu(II) or Fe(II/III) which promote its aggregation and/or ROS production. However, it is unclear whether this is the case in cells and whether/how extracellular A β can get into cells. Our recently developed molecular imaging probes for iron, copper and ROS enable us to look at this in real time in live cells at subcellular resolution. First, we tagged the A β covalently with a fluorescent dye which allows its interactions with cells to be monitored under a microscope. Our laser confocal imaging experiments with human neuroblastoma cells revealed that A β accumulated at the cell surface first and subsequently entered the cells via endocytosis pathway over a period of a few hours and finally deposited into endosomes/lysosomes in the cells. The deposition of A β induced a marked production of oxygen free radicals in the mitochondria of the cells, as revealed by our oxygen free radical probe and colocalization experiments. Incubation of metal ions such as copper(II) increased the production of oxygen radicals significantly while zinc(II) appears to be protective against ROS production. Our data provided compelling and direct evidence on how amyloid- β (A β) entering the cells and its induction of oxygen free radicals as well as the effects of metal ions on the radical production at subcellular level.