

9-2005

# The effect of IFN-gamma and TNF-alpha on the NADPH oxidase system of human colostrum macrophages, blood monocytes, and THP-1 cells

Ana Carolina Almeida  
*State University of Campinas Medical School*

Jussara Rehder  
*Universidade Estadual de Campinas*

Silvana Dalge Severino  
*State University of Campinas Medical School*

*See next page for additional authors*

Follow this and additional works at: [http://escholarship.umassmed.edu/peds\\_hematology](http://escholarship.umassmed.edu/peds_hematology)

 Part of the [Hematology Commons](#), [Oncology Commons](#), and the [Pediatrics Commons](#)

---

## Repository Citation

Almeida, Ana Carolina; Rehder, Jussara; Severino, Silvana Dalge; Martins-Filho, Jose; Newburger, Peter E.; and Condino-Neto, Antonio, "The effect of IFN-gamma and TNF-alpha on the NADPH oxidase system of human colostrum macrophages, blood monocytes, and THP-1 cells" (2005). *Hematology/Oncology*. 73.  
[http://escholarship.umassmed.edu/peds\\_hematology/73](http://escholarship.umassmed.edu/peds_hematology/73)

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in Hematology/Oncology by an authorized administrator of eScholarship@UMMS. For more information, please contact [Lisa.Palmer@umassmed.edu](mailto:Lisa.Palmer@umassmed.edu).

---

# The effect of IFN-gamma and TNF-alpha on the NADPH oxidase system of human colostrum macrophages, blood monocytes, and THP-1 cells

## **Authors**

Ana Carolina Almeida, Jussara Rehder, Silvana Dalge Severino, Jose Martins-Filho, Peter E. Newburger, and Antonio Condino-Neto

## **Comments**

Citation: J Interferon Cytokine Res. 2005 Sep;25(9):540-6. doi: 10.1089/jir.2005.25.540. [Link to article on publisher's website](#)

This is a copy of an article published in the Journal of Interferon & Cytokine Research © 2005 copyright Mary Ann Liebert, Inc. and available online at: <http://online.liebertpub.com>.

# The Effect of IFN- $\gamma$ and TNF- $\alpha$ on the NADPH Oxidase System of Human Colostrum Macrophages, Blood Monocytes, and THP-1 Cells

ANA CAROLINA ALMEIDA,<sup>1</sup> JUSSARA REHDER,<sup>1</sup> SILVANA DALGÈ SEVERINO,<sup>1</sup>  
JOSÉ MARTINS-FILHO,<sup>1</sup> PETER E. NEWBURGER,<sup>2</sup> and ANTONIO CONDINO-NETO<sup>1,3</sup>

## ABSTRACT

The aim of this work was to analyze the effect of Interferon- $\gamma$  (IFN- $\gamma$ ) and tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) on NADPH oxidase activity and gp91-*phox* gene expression in human colostrum macrophages (CM), peripheral blood monocytes (PBM), and myelomonocytic THP-1 cells. We also investigated the effect of IFN- $\gamma$  on the release of TNF- $\alpha$  by these cells. Our results show that under basal culture conditions, CM release more superoxide than PBM and THP-1 cells ( $p < 0.05$ ). The addition of IFN- $\gamma$ , alone or in combination with TNF- $\alpha$ , increased spontaneous superoxide release by PBM and THP-1 cells ( $p < 0.05$ ) and increased phorbol myristate acetate (PMA)-stimulated superoxide release by CM, PBM, and THP-1 cells ( $p < 0.05$ ). The NADPH oxidase activity of THP-1 cells consistently remained lower than that of CM or PBM, despite a dramatic response to IFN- $\gamma$  and TNF- $\alpha$ . Under basal conditions, gp91-*phox* gene expression was significantly higher in CM and PBM compared with THP-1 cells ( $p < 0.05$ ). The addition of IFN- $\gamma$  alone or in combination with TNF- $\alpha$  caused a dramatic increase in gp91-*phox* gene expression in THP-1 cells ( $p < 0.05$ ) but not in CM or PBM. Under basal conditions or in the presence of IFN- $\gamma$ , CM released more TNF- $\alpha$  than PBM or THP-1 cells ( $p < 0.05$ ). In addition, PBM released more TNF- $\alpha$  than THP-1 cells ( $p < 0.05$ ). IFN- $\gamma$  did not significantly augment the release of TNF- $\alpha$  by these cells ( $p > 0.05$ ). Thus, IFN- $\gamma$  and TNF- $\alpha$  induced equivalent gp91-*phox* gene expression in THP-1 cells compared with CM or PBM but did not bring about equivalent NADPH oxidase activity. TNF- $\alpha$  release was higher in more mature cells. This partial divergence of gp91-*phox* gene expression, NADPH oxidase activity, and TNF- $\alpha$  release is probably a consequence of different events of myeloid cell biology and relates at least in part to cell differentiation state.

## INTRODUCTION

HUMAN COLOSTRUM MACROPHAGES (CM) are mature monocytic cells that protect the mother against mastitis<sup>(1,2)</sup> and augment the immature immune system of the neonate.<sup>(3)</sup> These cells synthesize defense factors, including complement components (factors B, C2, C3, and C4),<sup>(4,5)</sup> lactoferrin, lysozyme,<sup>(3,6)</sup> granulocyte-macrophage colony-stimulating factor (GM-CSF),<sup>(7)</sup> leukotriene B<sub>4</sub> (LTB<sub>4</sub>), LTC<sub>4</sub>,<sup>(8)</sup> transforming growth factor- $\beta$  (TGF- $\beta$ ),<sup>(9)</sup> prostaglandin E<sub>2</sub> (PGE<sub>2</sub>).<sup>(10)</sup> They represent 30%–80% of the human colostrum leukocyte population, which varies from 500 to 10,000 cells/ml.<sup>(3)</sup> CM cells

express MHC class II molecules,<sup>(11)</sup> develop antibody-dependent cellular cytotoxicity,<sup>(12,13)</sup> ingest and kill bacteria,<sup>(14)</sup> and display the morphology and motility of activated cells.<sup>(3)</sup> Colostrum is an important and accessible source of human macrophages.

Professional phagocytes, such as neutrophils, eosinophils, monocytes, and macrophages, share a metabolic function termed the “respiratory burst,” which is triggered by inflammatory stimuli or infection. These stimuli activate the NADPH oxidase system, which catalyzes the donation of one electron from NADPH to molecular oxygen, converting it into superoxide anion (O<sub>2</sub><sup>-</sup>), which is rapidly converted to secondary toxic

<sup>1</sup>Departments of Pediatrics and Pharmacology and Center for Investigation in Pediatrics, State University of Campinas Medical School, Campinas, Brazil

<sup>2</sup>Department of Pediatrics, University of Massachusetts Medical School, Worcester, MA 01655

<sup>3</sup>Department of Immunology, Institute of Biomedical Sciences, University of São Paulo, Brazil.

oxygen species, such as hydrogen peroxide ( $H_2O_2$ ), hydroxyl radical ( $OH^\bullet$ ), and hypochlorous acid ( $HOCl$ ), all of which are important microbicidal agents.<sup>(15-17)</sup>

The phagocyte NADPH oxidase is a multicomponent enzyme system. The final electron donor is a heterodimeric flavocytochrome b, composed of a 91-kDa glycoprotein (gp91-*phox*) and a 22-kDa polypeptide (p22-*phox*). Following activation, the flavocytochrome b associates with Rap1A<sup>(15,18)</sup> and other cytosolic proteins, including p47-*phox*, p67-*phox*, p40-*phox*, and p21Rac.<sup>(19,20)</sup> The gp91-*phox* and p22-*phox* genes (also termed *CYBB* and *CYBA*, respectively) undergo parallel induction by various cytokines, including interferon- $\gamma$  (IFN- $\gamma$ ), in monocyte-derived macrophages and granulocytes.<sup>(21,22)</sup> In the human myelomonocytic THP-1 cell line, IFN- $\gamma$  and tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ) cause differentiation followed by a synergistic upregulation of the NADPH oxidase system, reflected by an enhancement of superoxide release, cytochrome *b*<sub>558</sub> content, and expression of the gp91-*phox* and p47-*phox* genes.<sup>(23)</sup>

The aim of the current work was to investigate the effects of IFN- $\gamma$  and TNF- $\alpha$  on the NADPH oxidase system of human myeloid cells in different developmental stages, analyzing both superoxide release and gp91-*phox* gene expression in CM, peripheral blood monocyte (PBM), and THP-1 cells. In addition, we analyzed the effect of IFN- $\gamma$  on the release of TNF- $\alpha$  by these cells.

## MATERIALS AND METHODS

### Cell isolation

Colostrum samples were manually collected and centrifuged (160g for 15 min at 4°C) to separate cells from the fat and aqueous layers. The cell button was resuspended in RPMI 1640 medium and centrifuged over a Ficoll-Paque density gradient 1077.<sup>(24)</sup> The upper layer containing mononuclear leukocytes was transferred to polystyrene plates, where CM were isolated by adherence. Monocytes were isolated by centrifugation of blood samples of the same donor over Ficoll-Paque and adherence on polystyrene plates. Colostrum and blood samples were collected up to 72 h postpartum. Macrophages/monocytes were 98% pure and showed >90% viability.

### Cell culture

Macrophages, PBM, and THP-1 cells were cultured under endotoxin-free conditions (<10 pg/ml) in RPMI 1640 medium supplemented with 10% heat-inactivated fetal bovine serum (FBS), 2 nM L-glutamine, 100 U/ml penicillin, and 100  $\mu$ g/ml streptomycin, at 37°C in a humid atmosphere with 5%  $CO_2$ .<sup>(25)</sup> IFN- $\gamma$  (100 U/ml) alone or in combination with TNF- $\alpha$  (1000 U/ml) was added to cultures for 48 h as indicated.

### NADPH oxidase activity

Superoxide release was quantified by superoxide dismutase (SOD)-inhibitable cytochrome *c* reduction as described McCord and Fridovich,<sup>(26)</sup> modified by Condino-Neto et al.<sup>(23)</sup> Briefly, on the day of the experiment, plates CM, PBM, or THP-1 cells were centrifuged, and the supernatant was re-

moved. The cells were incubated in Hanks' balanced salt solution (HBSS) (without phenol red) containing cytochrome *c* (50  $\mu$ M) for 1 h at 37°C in a humidified 5%  $CO_2$  atmosphere. Half of the wells received SOD (60 U/ml) at the beginning of the incubation. In another set of identical plates, phorbol myristate acetate (PMA) (30 nM) was added at the start of the incubation as an activator of superoxide release. After incubation, all plates were placed on ice, and the other half of the wells received SOD (60 U/ml). The plates were centrifuged again, and the absorbance of the supernatants was monitored at 550 nm. The results were calculated from the difference between the absorbance of the tubes with or without SOD during the incubation and expressed as nmol superoxide/ $10^6$  cells/h.

### gp91-phox gene expression

Expression of the gp91-*phox* gene (MIM.300481) was assessed by relative RT-PCR analysis using an Ambion QuantumRNA 18S Internal Standards kit (Ambion, Austin, TX). Total cell RNA was extracted by the guanidine HCl method.<sup>(27)</sup> Reverse transcription was performed with SuperScript II RT (GIBCO BRL, Gaithersburg, MD) and random hexamers. The cDNA was amplified by 30 cycles of PCR with oligonucleotide primers specific for gp91-*phox* (nucleotides [nt] 32 [exon 1]-443 [exon 5]) (GenBank accession number NM 000397). The PCR products were analyzed by electrophoresis in ethidium bromide-stained agarose gels. Relative gene expression was measured by Image Master software (Amersham Pharmacia Biotech, Piscataway, NJ).

### TNF- $\alpha$ release

For ELISA assays of TNF- $\alpha$ , 96-well plates were incubated with anti-TNF- $\alpha$  capture antibody overnight at ambient temperature. After receiving blocking solution, plates were incubated with cell culture supernatant, followed by detection antibody. After this incubation, plates were incubated with streptavidin-horseradish peroxidase (HRP), substrate solution, and stopping solution, respectively. The optical density (OD) was monitored in an ELISA reader at 450 nm. Results were expressed in picograms of TNF- $\alpha$ / $10^6$  cells (DuoSet Human TNF- $\alpha$ , R&D Systems, Minneapolis, MD).


### Statistics

Descriptive statistics were performed, and the results are represented by bar graphs showing mean  $\pm$  standard deviation (SD) values. Kruskal-Wallis and Mann-Whitney U-test were used for comparisons among groups;  $p < 0.05$  was considered significant.<sup>(28,29)</sup>

## RESULTS

### NADPH oxidase activity

Spontaneous superoxide release by CM ( $n = 13$ ) cultured in basal conditions was significantly higher compared with PBM ( $n = 20$ ) or THP-1 cells ( $n = 8$ ) ( $p < 0.05$ , Kruskal-Wallis test) (Fig. 1). The addition of IFN- $\gamma$  alone or in combination with TNF- $\alpha$  caused a significant increase in spontaneous superoxide release by PBM and THP-1 cells ( $p < 0.05$ , Mann-Whitney




**FIG. 1.** Bar graph showing mean  $\pm$  SD of spontaneous superoxide release by CM ( $n = 13$ ), PBM ( $n = 20$ ), and THP-1 cells ( $n = 8$ ) (nmol of superoxide/ $10^6$  cells/h). Cells were cultured in basal conditions or with IFN- $\gamma$  100 U/ml alone or in combination with TNF- $\alpha$  1000 U/ml for 2 days.  $p < 0.05$ . CM control  $>$  PBM control  $>$  THP-1 cells ( $p < 0.05$ , Kruskal-Wallis test); PBM control  $<$  PBM IFN- $\gamma$ /TNF- $\alpha$  ( $p < 0.05$ , Mann-Whitney); THP-1 control  $<$  THP-1 IFN- $\gamma$ /TNF- $\alpha$  ( $p < 0.05$ , Mann-Whitney).

test) (Fig. 1), but not by CM ( $p > 0.05$ , Mann-Whitney test) (Fig. 1).

PMA (30 nM)-stimulated superoxide production was measured in CM ( $n = 26$ ), PBM ( $n = 37$ ), and THP-1 ( $n = 31$ ) cells

cultured in basal conditions. Results were similar for CM and PBM ( $p > 0.05$ , Kruskal-Wallis test) (Fig. 2), but superoxide release by THP-1 cells was significantly lower compared with CM or PBM ( $p < 0.05$ , Kruskal-Wallis test) (Fig. 2). The ad-



**FIG. 2.** Bar graph showing mean  $\pm$  SD values of PMA-stimulated superoxide anion release (nmol of superoxide/ $10^6$  cells/h) by CM ( $n = 26$ ), PBM ( $n = 37$ ), and THP-1 cells ( $n = 31$ ). Cells were cultured in basal conditions or with IFN- $\gamma$  100 U/ml alone or in combination with TNF- $\alpha$  1000 U/ml for 2 days.  $*p < 0.05$ . THP-1 control  $<$  CM control = PBM control ( $p < 0.05$ , Kruskal-Wallis test); CM IFN- $\gamma$ /TNF- $\alpha$   $>$  CM control ( $p < 0.05$ , Mann-Whitney); PBM IFN- $\gamma$ /TNF- $\alpha$   $>$  PBM control ( $p < 0.05$ , Mann-Whitney); THP-1 IFN- $\gamma$ /TNF- $\alpha$   $>$  THP-1 control ( $p < 0.05$ , Mann-Whitney); THP-1 IFN- $\gamma$ /TNF- $\alpha$   $<$  CM IFN- $\gamma$ /TNF- $\alpha$  = PBM IFN- $\gamma$ /TNF- $\alpha$  ( $p < 0.05$ , Kruskal-Wallis test).

dition of IFN- $\gamma$  or IFN- $\gamma$  and TNF- $\alpha$  caused a significant increase in PMA-stimulated superoxide release by CM, PBM, and THP-1 cells ( $p < 0.05$ , Mann-Whitney test) (Fig. 2). However, PMA-stimulated superoxide by THP-1 cells cultured with IFN- $\gamma$  or IFN- $\gamma$  and TNF- $\alpha$  remained consistently lower compared with CM or PBM ( $p < 0.05$ , Kruskal-Wallis test) (Fig. 2).

*Gene expression studies*


Under basal conditions, gp91-*phox* gene expression was significantly higher in CM ( $n = 13$ ) and PBM ( $n = 15$ ) compared with THP-1 cells ( $n = 9$ ) ( $p < 0.05$ , Kruskal-Wallis test) (Fig. 3). The addition of IFN- $\gamma$  or of IFN- $\gamma$  and TNF- $\alpha$  caused a significant increase in gp91-*phox* gene expression in THP-1 cells ( $p < 0.05$ , Mann-Whitney test) (Fig. 3) but not in CM or PBM ( $p > 0.05$ , Mann-Whitney test) (Fig. 3). Thus, IFN- $\gamma$  in combination with TNF- $\alpha$  induced THP-1 cells to express equivalent levels of gp91-*phox* transcripts compared with CM or PBM, but without equivalent NADPH oxidase activity.

*TNF- $\alpha$  release*


Under basal conditions, the TNF- $\alpha$  release by CM ( $n = 6$ ) and PBM ( $n = 7$ ) was significantly higher compared with THP-1 cells ( $n = 6$ ) ( $p < 0.05$ , Kruskal-Wallis Test) (Fig. 4). Culturing CM, PBM, and THP-1 cells with IFN- $\gamma$  for 48 h caused a modest increase in the TNF- $\alpha$  release by these cells ( $p > 0.05$ , Mann-Whitney test) (Fig. 4). In addition, after 48 h culture with IFN- $\gamma$ , CM ( $n = 5$ ) and PBM ( $n = 7$ ) released higher levels of TNF- $\alpha$  compared with THP-1 ( $n = 6$ ) cells ( $p < 0.05$ , Kruskal-Wallis Test) (Fig. 4).

**DISCUSSION**

In the current model, THP-1 cells represent an early phase of mononuclear phagocyte development, PBM are in an intermediate state of development, and CM represent tissue cells in a terminal maturation state. Our results show that under basal



**FIG. 3.** (A) Ethidium bromide-stained agarose gel of RT-PCR products representing gp91-*phox* gene expression in CM, PBM, and THP-1 cells. Cells were cultured in basal conditions or with IFN- $\gamma$  100 U/ml and TNF- $\alpha$  1000 U/ml for 2 days. The upper band represents gp91-*phox* transcripts, and the lower band represents 18S rRNA normalization controls. Lane 1, base pair marker (100 bp); lane 2, CM control; lane 3, PBM control; lane 4, CM + IFN- $\gamma$ /TNF- $\alpha$ ; lane 5, PBM + IFN- $\gamma$ /TNF- $\alpha$ ; lane 6, THP-1 cell control; lane 7, THP-1 cells + IFN- $\gamma$ /TNF- $\alpha$ . (B) Bar graphs showing mean  $\pm$  SD for relative expression of the gp91-*phox* gene in CM ( $n = 13$ ), PBM ( $n = 15$ ) and THP-1 cells ( $n = 9$ ). Cells were cultured in basal conditions and with IFN- $\gamma$  100 U/ml alone or in combination with TNF- $\alpha$  1000 U/ml. CM = PBM > THP-1 (control) ( $p < 0.05$ , Kruskal-Wallis test); THP-1 IFN- $\gamma$ /TNF- $\alpha$  > THP-1 control ( $p < 0.05$ , Mann-Whitney). Asterisks indicate that control < IFN- $\gamma$  < IFN- $\gamma$ /TNF- $\alpha$ .



**FIG. 4.** Bar graphs showing mean  $\pm$  SD values of TNF- $\alpha$  release (pg TNF- $\alpha$ /10<sup>6</sup> cells) by CM ( $n = 11$ ), PBM ( $n = 14$ ), and THP-1 cells ( $n = 12$ ). Cells were cultured in basal conditions or with IFN- $\gamma$  100 U/ml. \* $p < 0.05$ . THP-1 control < PBM control < CM control ( $p < 0.05$ , Kruskal-Wallis test); THP-1 IFN- $\gamma$  < PBM IFN- $\gamma$  < CM IFN- $\gamma$  ( $p < 0.05$ , Kruskal-Wallis test); CM control > PBM control ( $p < 0.05$ , Mann-Whitney); CM control > THP-1 control ( $p < 0.05$ , Mann-Whitney); PBM control > THP-1 control ( $p < 0.05$ , Mann-Whitney); CM IFN- $\gamma$  > PBM IFN- $\gamma$  ( $p < 0.05$ , Mann-Whitney); CM IFN- $\gamma$  > THP-1 IFN- $\gamma$  ( $p < 0.05$ , Mann-Whitney); PBM IFN- $\gamma$  > THP-1 IFN- $\gamma$  ( $p < 0.05$ , Mann-Whitney).

conditions, CM spontaneously release more superoxide than PBM and THP-1 cells. The addition of IFN- $\gamma$  or of IFN- $\gamma$  plus TNF- $\alpha$  caused a significant increase in spontaneous superoxide release by PBM and THP-1 cells and also increased PMA-stimulated superoxide release by CM, PBM, and THP-1 cells. IFN- $\gamma$  alone or in combination with TNF- $\alpha$  did not enhance gp91-*phox* gene expression by CM or PBM, unlike THP-1 cells, in which the cytokines induced a significant increase in the expression of this gene encoding a key component of the NADPH oxidase complex. Accordingly, CM released more TNF- $\alpha$  than PBM, which in turn released more TNF- $\alpha$  than THP-1 cells, showing that TNF- $\alpha$  release correlates with the mononuclear phagocyte maturation state.

The results obtained with PBM confirmed previous observations<sup>(30,31)</sup> in which culturing human monocytes with IFN- $\gamma$  and TNF- $\alpha$  for 48 h caused them to release more superoxide. Cummings et al.,<sup>(32)</sup> measuring superoxide release by both CM and PBM freshly isolated from the same individual, also found that spontaneous superoxide release from CM exceeds that from PBM, but after PMA stimulation, the values became equivalent.

The current results comparing stimulated superoxide release from CM and PBM also concur with previous observations. Tsuda et al.<sup>(33)</sup> showed that CM stimulated by opsonized zymosan or PMA release similar amounts of superoxide anion and H<sub>2</sub>O<sub>2</sub> compared with monocytes. These findings were confirmed by Speer et al.,<sup>(14)</sup> who showed that freshly isolated, PMA-stimulated CM release superoxide and H<sub>2</sub>O<sub>2</sub> at rates similar to those of PBM. Furthermore, Pabst et al.<sup>(34)</sup> showed that

human milk macrophages, peritoneal macrophages, and human blood monocytes appeared to be already primed for a vigorous oxygen radical response.

The current experiments also confirm our previous finding<sup>(23)</sup> that culturing THP-1 cells with IFN- $\gamma$  alone or in combination with TNF- $\alpha$  for 48 h induces these cells to differentiate, to express increased NADPH oxidase activity and cytochrome *b* content, and to upregulate genes encoding NADPH oxidase components. Even with the considerable induction of NADPH oxidase activity in THP-1 cells by IFN- $\gamma$  and TNF- $\alpha$ , however, it is not sufficient to reach the levels expressed in PBM and CM.

The higher amounts of TNF- $\alpha$  release by more differentiated cells do not definitively account for these observations. The most probable explanation is that the NADPH oxidase system in mononuclear phagocytes is activated during its *in vivo* maturation by a large number of chemical stimuli with which THP-1 cells do not have contact during *in vitro* maturation. These stimuli include not only cytokines and chemokines but also NADPH oxidase products. The reactive oxygen species (ROS) regulate cellular growth, differentiation, proliferation, apoptosis, and gene expression.<sup>(35–38)</sup> Furthermore, THP-1 cells are a leukemia-derived, continuous cell line with distinct genotypic and phenotypic characteristics.

Our gene expression studies showed that the cytokines IFN- $\gamma$  and TNF- $\alpha$  upregulated gp91-*phox* expression in THP-1 cells, as shown in our previous work,<sup>(23)</sup> and reached levels equivalent to CM and PBM. However, the colostrum and blood-derived cells did not show a significant change in gp91-*phox* gene

expression in response to these cytokines. Notably, CM released the highest amounts of TNF- $\alpha$  compared with PBM or THP-1 cells, and the release of this cytokine was not subject to strong modulation by IFN- $\gamma$ , showing that other factors may influence this function *in vivo*.

In contrast, Newburger et al.<sup>(22)</sup> reported that human monocyte-derived macrophages cultured for 24 h with IFN- $\gamma$  show a 12-fold increase in gp91-*phox* gene expression and that treatment with lipopolysaccharide (LPS), TNF- $\alpha$ , or macrophage CSF (M-CSF) or GM-CSF increases gp91-*phox* and, to a lesser extent, p22-*phox* transcript levels. Cassatella et al.<sup>(39)</sup> also showed that a 24–48-h treatment with IFN- $\gamma$  increases gp91-*phox* levels in monocyte-derived macrophages. The differences between our results and those previously reported may be attributed to differences in culture time and cytokine concentration.

Thus, gp91-*phox* gene expression is already highly expressed in CM and PBM, probable because of the effects of stimuli received during maturation, such as interleukin-3 (IL-3), CSF-1, GM-CSF, IL-4, and IL-13,<sup>(40)</sup> and also because of the autocrine modulation by TNF- $\alpha$  and ROS. THP-1 cells are undifferentiated precursor of monocytes and macrophages and, thus, are able to respond more to IFN- $\gamma$  and TNF- $\alpha$ . In addition, their leukemic origin<sup>(23)</sup> may contribute to a different response compared with primary monocytes and macrophages.

Winston et al.<sup>(41)</sup> showed that TNF- $\alpha$  mediates macrophage differentiation and IFN- $\gamma$  appears to act as a molecular switch. In the presence of IFN- $\gamma$ , stimulation of macrophages with TNF- $\alpha$  causes macrophage differentiation along a pathway in which inducible nitric oxide synthase (iNOS) is expressed, whereas in the absence of IFN- $\gamma$ , stimulation of macrophages with TNF- $\alpha$  causes differentiation along a pathway in which insulin-like growth factor-1 (IGF-1) is expressed.

Our experiments show that IFN- $\gamma$  did not strongly modulate TNF- $\alpha$  release by CM, PBM, or THP-1 cells. Monocytes and macrophages are a major source of TNF- $\alpha$ .<sup>(42)</sup> Studies have shown that LPS-induced TNF- $\alpha$  release is, at least in part, a result of NF- $\kappa$ B activation.<sup>(43,44)</sup> LPS rapidly activates TNF- $\alpha$  gene transcription, which causes an increase in steady-state TNF- $\alpha$  mRNA levels, the translation of which accounts for the observed dramatic increase in secreted TNF- $\alpha$  protein.<sup>(45)</sup>

Our results also show that TNF- $\alpha$  release is directly related to the cell maturation state. Thus, TNF- $\alpha$  release in culture medium may provide an autocrine mechanism that can influence the NADPH oxidase system, potentiating cell responses to other cytokines.

The release of superoxide is a consequence of gene expression and downstream events, including translation, RNA and protein stability, receptor activation, and signal transduction, all of which can vary according to a cell's stage of differentiation. We conclude that regulation of the NADPH oxidase system in human myelomonocytic cells by IFN- $\gamma$  and TNF- $\alpha$  is a consequence of different events of cell biology and is related at least in part to cell differentiation. Better understanding of NADPH oxidase regulation will strengthen the basis for modulating its expression in conditions with high risk for oxidant-mediated inflammation and tissue injury. Specific knowledge of human CM function should shed light on the contribution of breastfeeding to the host defense of the neonate.

## ACKNOWLEDGMENTS

This work was supported by the Fundação de Amparo à Pesquisa do Estado de São Paulo (grants 98/16003-7 and 00/04818-8), National Institutes of Health (R01 DK54369), and the Fogarty International Center, National Institutes of Health (R03 TW00883).

## REFERENCES

- Balkwill FR, Hogg N. Characterization of human breast milk macrophages cytostatic for human cell lines. *J Immunol.* 1979;123:1451–1456.
- Outteridge PM, Lee CS. Cellular immunity in the mammary gland with particular reference to T, B lymphocytes and macrophages. *Adv Exp Med Biol.* 1981;137:513–534.
- Goldman AS, Goldblum RM. Transfer of maternal leukocytes to the infant by human milk. *Curr Top Microbiol Immunol.* 1997;222:205–213.
- Cole FS, Schneeberger EE, Lichtenberg NA, Colten HR. Complement biosynthesis in human breast-milk macrophages and blood monocytes. *Immunology.* 1982;46:429–441.
- Goldman AS, Thorpe LW, Goldblum RM, Hanson LA. Anti-inflammatory properties of human milk. *Acta Paediatr Scand.* 1986;75:689–695.
- Vassao RC, Carneiro-Sampaio MM. Phagocytic activity of human colostrum macrophages. *Braz J Med Biol Res.* 1989;22:457–464.
- Ichikawa M, Sugita M, Takahashi M, Satomi M, Takeshita T, Araki T, Takahashi H. Breast milk macrophages spontaneously produce granulocyte-macrophage colony-stimulating factor and differentiate into dendritic cells in the presence of exogenous interleukin-4 alone. *Immunology.* 2003;108:189–195.
- Shimizu T, Niklasson A, Strandvik B. Leukotriene B4 and C4 generation by human colostrum macrophages *in vitro*. *Biol Neonate.* 1993;63:65–69.
- Wagner CL, Forsythe DW, Pittard WB. Variation in the biochemical forms of transforming growth factor- $\alpha$  present in human milk and secreted by human milk macrophages. *Biol Neonate.* 1995;68:325–333.
- Blau H, Passwell JH, Levanon M, Davidson J, Kohen F, Ramot B. Studies on human milk macrophages: effect of activation on phagocytosis and secretion of prostaglandin E<sub>2</sub> and lysozyme. *Pediatr Res.* 1983;17:241–245.
- Leyva-Cobian F, Clemente J. Phenotypic characterization and functional activity of human milk macrophages. *Immunol Lett.* 1984;8:249–256.
- Mandyla H, Xanthou M, Maravelias C, Baum D, Matsaniotis N. Antibody dependent cytotoxicity of human colostrum phagocytes. *Pediatr Res.* 1982;16:995–999.
- Kohl S, Pickering LK, Cleary TG, Steinmetz KD, Loo LS. Human colostrum cytotoxicity. II. Relative defects in colostrum leukocyte cytotoxicity and inhibition of peripheral blood leukocyte cytotoxicity by colostrum. *J Infect Dis.* 1980;142:884–891.
- Speer CP, Gahr M, Pabst MJ. Phagocytosis-associated oxidative metabolism in human milk macrophages. *Acta Paediatr Scand.* 1986;75:444–451.
- Leusen JHW, Verhoeven AJ, Roos D. Interactions between the components of the human NADPH oxidase: a review about the intrigues in the *phox* family. *Front Biosci.* 1996;1:d72–90.
- Quinn MT, Gauss KA. Structure and regulation of the neutrophil respiratory burst oxidase: comparison with nonphagocyte oxidases. *J Leukocyte Biol.* 2004;76:760–781.
- Cathcart MK. Regulation of superoxide anion production by NADPH



- oxidase in monocytes/macrophages: contributions to atherosclerosis. *Arterioscler Thromb Vasc Biol.* 2004;24:23–28.
18. Nauseef WM. Assembly of the phagocyte NADPH oxidase. *Histochem Cell Biol.* 2004;122:277–291.
  19. Hausmann M, Spottl T, Andus T, Rothe G, Falk W, Scholmerich J, Herfarth H, Rogler G. Subtractive screening reveals up-regulation of NADPH oxidase expression in Crohn's disease intestinal macrophages. *Clin Exp Immunol.* 2001;125:48–55.
  20. Lacy P, Mahmudi-Azer S, Bablitz B, Gilchrist M, Fitzharris P, Cheng D, Man SF, Bokoch GM, Moqbel R. Expression and translocation of Rac2 in eosinophils during superoxide generation. *Immunology.* 1999;98:244–252.
  21. Newburger PE, Ezekowitz RA, Whitney C, Wright J, Orkin SH. Induction of phagocyte cytochrome b heavy chain gene expression by interferon gamma. *Proc Natl Acad Sci USA.* 1988;85:5215–5219.
  22. Newburger PE, Dai Q, Whitney C. *In vitro* regulation of human phagocyte cytochrome b heavy and light chain gene expression by bacterial lipopolysaccharide and recombinant human cytokines. *J Biol Chem.* 1991;266:16171–16177.
  23. Condino-Neto A, Whitney C, Newburger PE. Dexamethasone but not indomethacin inhibits human phagocyte nicotinamide adenine dinucleotide phosphate oxidase activity by down-regulating expression of genes encoding oxidase components. *J Immunol.* 1998;161:4960–4967.
  24. Boyum A. Isolation of mononuclear cells and granulocytes from human blood. Isolation of mononuclear cells by one centrifugation, and of granulocytes by combining centrifugation and sedimentation at 1 g. *Scand J Clin Lab Invest.* 1968;97(Suppl):77–89.
  25. Fischkoff SA. Graded increase in probability of eosinophilic differentiation of HL-60 promyelocytic leukemia cells induced by culture under alkaline conditions. *Leuk Res.* 1988;12:679–686.
  26. McCord JM, Fridovich I. Superoxide dismutase. An enzymic function for erythrocyte (hemocuprein). *J Biol Chem.* 1969;244:6049–6055.
  27. Ginsburg D, Handin RI, Bonthron DT, Donlon TA, Bruns GA, Latt SA, Orkin SH. Human von Willebrand factor (vWF): isolation of complementary DNA (cDNA) clones and chromosomal localization. *Science.* 1985;228:1401–1406.
  28. Milliken GA, Johnson, DE. *Analysis of Messy Data.* New York: Van Nostrand Reinhold Company; 1984.
  29. Montgomery DC. *Design and Analysis of Experiments.* 3rd ed. New York: John Wiley & Sons; 1991.
  30. Szeffler S, Norton CE, Ball B, Gross JM, Aida Y, Pabst MJ. IFN-gamma and LPS overcome glucocorticoid inhibition of priming for superoxide release in human monocytes. Evidence that secretion of IL-1 and tumor necrosis factor-alpha is not essential for monocyte priming. *J Immunol.* 1989;142:3985–3992.
  31. Dusi S, Donini, M, Lissandrini, D, Mazzi, P, Bianca, VD, Rossi, F. Mechanisms of expression of NADPH oxidase components in human cultured monocytes: role of cytokines and transcriptional regulators involved. *Eur J Immunol.* 2001;31:929–938.
  32. Cummings NP, Neifert MR, Pabst MJ, Johnston RB Jr. Oxidative metabolic response and microbicidal activity of human milk macrophages: effect of lipopolysaccharide and muramyl dipeptide. *Infect Immun.* 1985;49:435–439.
  33. Tsuda H, Takeshige K, Shibata Y, Minkami S. Oxygen metabolism of human colostrum macrophages: comparison with monocytes and polymorphonuclear leukocytes. *J Biochem.* 1984;95:1237–1245.
  34. Pabst MJ, Hedegaard HB, Johnston RB Jr. Cultured human monocytes require exposure to bacterial products to maintain an optimal oxygen radical response. *J Immunol.* 1982;128:123–128.
  35. Yang S, Hardaway M, Sun G, Ries WL, Key LL Jr. Superoxide generation and tyrosine kinase. *Biochem Cell Biol.* 2000;78:11–17.
  36. Irani K. Oxidant signaling in vascular cell growth, death, and survival: a review of the roles of reactive oxygen species in smooth muscle and endothelial cell mitogenic and apoptotic signaling. *Circ Res.* 2000;87:179–183.
  37. Sauer H, Wartenberg M, Hescheler J. Reactive oxygen species as intracellular messengers during cell growth and differentiation. *Cell Physiol Biochem.* 2001;11:173–186.
  38. Barbieri SS, Eligini S, Brambilla M, Tremoli E, Colli S. Reactive oxygen species mediate cyclooxygenase-2 induction during monocyte to macrophage differentiation: critical role of NADPH oxidase. *Cardiovasc Res.* 2003;60:187–197.
  39. Cassatella MA, Bazzoni F, Flynn RM, Dusi S, Trinchieri G, Rossi F. Molecular basis of interferon-gamma and lipopolysaccharide enhancement of phagocyte respiratory burst capability. Studies on the gene expression of several NADPH oxidase components. *J Biol Chem.* 1990;265:20241–20246.
  40. Skalnik DG. Transcriptional mechanisms regulating myeloid-specific genes. *Gene.* 2002;284:1–21.
  41. Winston BW, Krein PM, Mowat C, Huang Y. Cytokine-induced macrophage differentiation: a tale of 2 genes. *Clin Invest Med.* 1999;22:236–255.
  42. Vassalli P. The pathophysiology of tumor necrosis factors. *Annu Rev Immunol.* 1992;10:411–452.
  43. Udalova IA, Knight JC, Vidal V, Nedospasov SA, Kwiatkowski D. Complex NF- $\kappa$ B interactions at the distal tumor necrosis factor promoter region in human monocytes. *J Biol Chem.* 1998;273:21178–21186.
  44. Liu H, Sidiropoulos P, Song G, Pagliari LJ, Birrer MJ, Stein B, Anrather J, Pope RM. TNF-alpha gene expression in macrophages: regulation by NF-kappa B is independent of c-Jun or C/EBP beta. *J Immunol.* 2000;164:4277–4285.
  45. Raabe T, Bukrinsky M, Currie RA. Relative contribution of transcription and translation to the induction of tumor necrosis factor-alpha by lipopolysaccharide. *J Biol Chem.* 1998;273:974–980.

Address reprint requests to:  
 Dr. Antonio Condino-Neto  
 Department of Immunology  
 Institute of Biomedical Sciences  
 University of São Paulo  
 1730 Lineu Prestes Avenue  
 São Paulo, SP 05508-900  
 Brazil

Tel: 55 19 3289-3103  
 Fax: 55 19 3289-8638  
 E-mail: condino@icb.usp.br

Received 15 September 2004/Accepted 5 April 2005

**This article has been cited by:**

1. Gorazd Tompa, Aire Laine, Anne Pihlanto, Hannu Korhonen, Irena Rogelj, Pertti Marnila. 2010. Chemiluminescence of non-differentiated THP-1 promonocytes: developing an assay for screening anti-inflammatory milk proteins and peptides. *Luminescence* n/a-n/a. [[CrossRef](#)]
2. Miho Sakai, Andreas Vonderheit, Xun Wei, Claudia Küttel, Andreas Stemmer. 2009. A novel biofuel cell harvesting energy from activated human macrophages. *Biosensors and Bioelectronics* **25**:1, 68-75. [[CrossRef](#)]
3. Grace Y. Sun, Lloyd A. Horrocks, Akhlaq A. Farooqui. 2007. The roles of NADPH oxidase and phospholipases A<sub>2</sub> in oxidative and inflammatory responses in neurodegenerative diseases. *Journal of Neurochemistry*, ahead of print070611013409004. [[CrossRef](#)]
4. Livia E. Marçal, Péricles M. Dias-da-Motta, Jussara Rehder, Ronei L. Mamoni, Maria Heloísa S.L. Blotta, Constance B. Whitney, Peter E. Newburger, Fernando F. Costa, Sara T.O. Saad, Antonio Condino-Neto. 2007. Up-regulation of NADPH oxidase components and increased production of interferon-gamma by leukocytes from sickle cell disease patients. *American Journal of Hematology* . [[CrossRef](#)]
5. Qing Lan, Tongzhang Zheng, Min Shen, Yawei Zhang, Sophia S. Wang, Shelia H. Zahm, Theodore R. Holford, Brian Leaderer, Peter Boyle, Stephen Chanock. 2007. Genetic polymorphisms in the oxidative stress pathway and susceptibility to non-Hodgkin lymphoma. *Human Genetics* **121**:2, 161. [[CrossRef](#)]
6. Karen Y. Stokes . 2006. Microvascular Responses to Hypercholesterolemia: The Interactions Between Innate and Adaptive Immune Responses. *Antioxidants & Redox Signaling* **8**:7-8, 1141-1151. [[Abstract](#)] [[PDF](#)] [[PDF Plus](#)]