

May 22nd, 4:30 PM - 6:00 PM

Use of the BRFSS Follow-up Surveys to Assess Awareness of the Massachusetts Medicaid Cessation Benefit

Mark Paskowsky
Massachusetts Department of Public Health

Lois Keithly
Massachusetts Department of Public Health

Lili Chen
University of Massachusetts Medical School

See next page for additional authors

Follow this and additional works at: https://escholarship.umassmed.edu/cts_retreat

Part of the [Community Health and Preventive Medicine Commons](#)

Paskowsky, Mark; Keithly, Lois; Chen, Lili; and Li, Wenjun, "Use of the BRFSS Follow-up Surveys to Assess Awareness of the Massachusetts Medicaid Cessation Benefit" (2012). *UMass Center for Clinical and Translational Science Research Retreat*. 51.
https://escholarship.umassmed.edu/cts_retreat/2012/posters/51

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.

Presenter Information

Mark Paskowsky, Lois Keithly, Lili Chen, and Wenjun Li

Creative Commons License

Creative

Commons

This work is licensed under a [Creative Commons Attribution-Noncommercial-Share Alike 3.0 License](#).

Attribution-

Noncommercial-

Share

Alike

3.0

License

USE OF THE BRFSS FOLLOW-UP SURVEYS TO ASSESS AWARENESS OF THE MASSACHUSETTS MEDICAID CESSATION BENEFIT

Mark Paskowsky¹, MS, Lois Keithly¹, PhD, Lili Chen², MS, Wenjun Li², PhD

¹Massachusetts (MA) Tobacco Cessation and Prevention Program, MA Department of Public Health, Boston, MA; ²Division of Preventive and Behavioral Medicine, University of Massachusetts Medical School, Worcester, MA

Contact: Kevin.Kane1@umassmed.edu 508-856-3973

Background: Conducting a BRFSS follow-up survey is an effective method for evaluating special program initiatives. Massachusetts Tobacco Control Program (MTCP) staff designed a brief telephone questionnaire to measure awareness of a new tobacco cessation benefit for the Medicaid population and to identify barriers to utilization of the benefit.

Methods: A telephone call back was conducted by the Massachusetts Department of Public Health (MDPH) from April 2007 – October 2007, 10-16 months after implementation of the smoking cessation benefit. It consisted of a sample of BRFSS respondents between the ages of 18 and 64 years interviewed between March 2007 and September 2007 who agreed to be called back for a follow-up survey. The data consists mainly of smoking and smoking cessation-related questions, insurance related questions, and questions regarding healthcare utilization.

Results: Sixty-two percent of BRFSS respondents provided consent to participate in the follow-up survey. Of those, forty-three percent responded to the survey. Of the 2,399 follow-up survey completions, 487 were Medicaid members. Survey results show that Medicaid smokers at follow-up had a high awareness of quitline services (64%) and the benefit (32%). Compared to non-Medicaid smokers, more Medicaid smokers reported intentions to use tobacco cessation medication if referred by a health care provider (55% vs. 47%, $p=.04$).

Conclusions: Using the BRFSS call-back survey approach made it possible to administer the survey to smokers in the Medicaid population without a laborious and expensive process of screening Medicaid eligible smokers from the general population. Thus, the approach is cost-effective and population-based. The results showed high awareness of quitline services and smoking cessation of the Medicaid smoking cessation benefit, and, most importantly, a higher rate of reported intention to use cessation medication among the Medicaid smoking population.