

May 22nd, 4:30 PM - 6:00 PM

Use of the BRFSS Follow-up Surveys to Assess Awareness of the Massachusetts Medicaid Cessation Benefit

Mark Paskowsky

Massachusetts Department of Public Health, Mark.Paskowsky@state.ma.us

Lois Keithly

Massachusetts Department of Public Health

Lili Chen

University of Massachusetts Medical School, lili.chen@umassmed.edu

See next page for additional authors

Follow this and additional works at: http://escholarship.umassmed.edu/cts_retreat



Part of the [Community Health and Preventive Medicine Commons](#)

Mark Paskowsky, Lois Keithly, Lili Chen, and Wenjun Li, "Use of the BRFSS Follow-up Surveys to Assess Awareness of the Massachusetts Medicaid Cessation Benefit" (May 22, 2012). *UMass Center for Clinical and Translational Science Research Retreat*. Paper 51.

http://escholarship.umassmed.edu/cts_retreat/2012/posters/51

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.

Presenter Information

Mark Paskowsky, Lois Keithly, Lili Chen, and Wenjun Li

Creative Commons License

This work is licensed under a [Creative Commons Attribution-Noncommercial-Share Alike 3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/).

USE OF THE BRFSS FOLLOW-UP SURVEYS TO ASSESS AWARENESS OF THE MASSACHUSETTS MEDICAID CESSATION BENEFIT

Mark Paskowsky¹, MS, Lois Keithly¹, PhD, Lili Chen², MS, Wenjun Li², PhD

¹Massachusetts (MA) Tobacco Cessation and Prevention Program, MA Department of Public Health, Boston, MA; ²Division of Preventive and Behavioral Medicine, University of Massachusetts Medical School, Worcester, MA

Contact: Kevin.Kane1@umassmed.edu 508-856-3973

Background: Conducting a BRFSS follow-up survey is an effective method for evaluating special program initiatives. Massachusetts Tobacco Control Program (MTCP) staff designed a brief telephone questionnaire to measure awareness of a new tobacco cessation benefit for the Medicaid population and to identify barriers to utilization of the benefit.

Methods: A telephone call back was conducted by the Massachusetts Department of Public Health (MDPH) from April 2007 – October 2007, 10-16 months after implementation of the smoking cessation benefit. It consisted of a sample of BRFSS respondents between the ages of 18 and 64 years interviewed between March 2007 and September 2007 who agreed to be called back for a follow-up survey. The data consists mainly of smoking and smoking cessation-related questions, insurance related questions, and questions regarding healthcare utilization.

Results: Sixty-two percent of BRFSS respondents provided consent to participate in the follow-up survey. Of those, forty-three percent responded to the survey. Of the 2,399 follow-up survey completions, 487 were Medicaid members. Survey results show that Medicaid smokers at follow-up had a high awareness of quitline services (64%) and the benefit (32%). Compared to non-Medicaid smokers, more Medicaid smokers reported intentions to use tobacco cessation medication if referred by a health care provider (55% vs. 47%, $p=.04$).

Conclusions: Using the BRFSS call-back survey approach made it possible to administer the survey to smokers in the Medicaid population without a laborious and expensive process of screening Medicaid eligible smokers from the general population. Thus, the approach is cost-effective and population-based. The results showed high awareness of quitline services and smoking cessation of the Medicaid smoking cessation benefit, and, most importantly, a higher rate of reported intention to use cessation medication among the Medicaid smoking population.