

2-2003

Targeted disruption of the mouse Mel(1b) melatonin receptor

Xiaowei Jin
Harvard Medical School

Charlotte von Gall
University of Massachusetts Medical School, Worcester

Rick L. Pieschl
Bristol-Myers Squibb Pharmaceutical Research Institute

See next page for additional authors

Follow this and additional works at: http://escholarship.umassmed.edu/neurobiology_pp

 Part of the [Neuroscience and Neurobiology Commons](#)

Repository Citation

Jin, Xiaowei; von Gall, Charlotte; Pieschl, Rick L.; Gribkoff, Valentin K.; Stehle, Jorg H.; Reppert, Steven M.; and Weaver, David R., "Targeted disruption of the mouse Mel(1b) melatonin receptor" (2003). *Neurobiology Publications and Presentations*. 86.
http://escholarship.umassmed.edu/neurobiology_pp/86

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in Neurobiology Publications and Presentations by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.

Targeted disruption of the mouse Mel(1b) melatonin receptor

Authors

Xiaowei Jin, Charlotte von Gall, Rick L. Pieschl, Valentin K. Gribkoff, Jorg H. Stehle, Steven M. Reppert, and David R. Weaver

Comments

Citation: Mol Cell Biol. 2003 Feb;23(3):1054-60. [Link to article on publisher's website](#)

Publisher PDF posted as allowed by the publisher's author rights policy at http://journals.asm.org/site/misc/ASM_Author_Statement.xhtml.

**Targeted Disruption of the Mouse Mel_{1b}
Melatonin Receptor**

Xiaowei Jin, Charlotte von Gall, Rick L. Pieschl, Valentin K. Gribkoff, Jorg H. Stehle, Steven M. Reppert and David R. Weaver

Mol. Cell. Biol. 2003, 23(3):1054. DOI:
10.1128/MCB.23.3.1054-1060.2003.

Updated information and services can be found at:
<http://mcb.asm.org/content/23/3/1054>

These include:

REFERENCES

This article cites 43 articles, 13 of which can be accessed free at: <http://mcb.asm.org/content/23/3/1054#ref-list-1>

CONTENT ALERTS

Receive: RSS Feeds, eTOCs, free email alerts (when new articles cite this article), [more»](#)

Information about commercial reprint orders: <http://journals.asm.org/site/misc/reprints.xhtml>
To subscribe to to another ASM Journal go to: <http://journals.asm.org/site/subscriptions/>

Targeted Disruption of the Mouse Mel_{1b} Melatonin Receptor

Xiaowei Jin,^{1†} Charlotte von Gall,^{2,3‡} Rick L. Pieschl,⁴ Valentin K. Gribkoff,⁴
Jorg H. Stehle,² Steven M. Reppert,^{1,3‡} and David R. Weaver^{1,3*}

Laboratory of Developmental Chronobiology, MassGeneral Hospital for Children, Massachusetts General Hospital and Harvard Medical School, Boston, Massachusetts 02114¹; Anatomisches Institut II, Johann Wolfgang Goethe-Universität Frankfurt, D-60590 Frankfurt, Germany²; Department of Neurobiology, University of Massachusetts Medical School, Worcester, Massachusetts 01605³; and Neuroscience Drug Discovery, Bristol-Myers Squibb Pharmaceutical Research Institute, Wallingford, Connecticut 06492⁴

Received 29 August 2002/Returned for modification 16 October 2002/Accepted 7 November 2002

Two high-affinity, G protein-coupled melatonin receptor subtypes have been identified in mammals. Targeted disruption of the Mel_{1a} melatonin receptor prevents some, but not all, responses to the hormone, suggesting functional redundancy among receptor subtypes (Liu et al., *Neuron* 19:91-102, 1997). In the present work, the mouse Mel_{1b} melatonin receptor cDNA was isolated and characterized, and the gene has been disrupted. The cDNA encodes a receptor with high affinity for melatonin and a pharmacological profile consistent with its assignment as encoding a melatonin receptor. Mice with targeted disruption of the Mel_{1b} receptor have no obvious circadian phenotype. Melatonin suppressed multiunit electrical activity in the suprachiasmatic nucleus (SCN) in Mel_{1b} receptor-deficient mice as effectively as in wild-type controls. The neuropeptide, pituitary adenylyl cyclase activating peptide, increases the level of phosphorylated cyclic AMP response element binding protein (CREB) in SCN slices, and melatonin reduces this effect. The Mel_{1a} receptor subtype mediates this inhibitory response at moderate ligand concentrations (1 nM). A residual response apparent in Mel_{1a} receptor-deficient C3H mice at higher melatonin concentrations (100 nM) is absent in Mel_{1a}-Mel_{1b} double-mutant mice, indicating that the Mel_{1b} receptor mediates this effect of melatonin. These data indicate that there is a limited functional redundancy between the receptor subtypes in the SCN. Mice with targeted disruption of melatonin receptor subtypes will allow molecular dissection of other melatonin receptor-mediated responses.

The hormone, melatonin, is produced rhythmically in the vertebrate pineal gland (13). Rhythmic melatonin production plays a critical role in the regulation of reproduction in seasonally breeding mammals (1, 24). In nonmammalian vertebrates, melatonin also plays a major role in the regulation of circadian rhythms (4, 46). While endogenous melatonin appears to play only a subtle role in the regulation of circadian rhythms in mammals, exogenous melatonin influences circadian rhythms in several rodent species and in humans (2, 3, 9, 21, 22, 39, 46, 47). While the magnitude of this phase shifting effect is small for adult mammals, it is nevertheless critical in some situations; e.g., daily melatonin administration appears to be useful for entrainment of non-24-h circadian cycles to the 24-hour day for blind individuals (22).

Two high-affinity receptors for melatonin have been identified in mammals (for reviews, see references 33 and 43). These receptors, the Mel_{1a} and Mel_{1b} receptor subtypes, are members of the G protein-coupled receptor superfamily (29, 30). A melatonin receptor-related receptor (also called H9 and GPR50) has sequence homology to the Mel_{1a} and Mel_{1b} re-

ceptors but does not bind melatonin (7, 16, 32). In addition to these three members of the gene family in mammals, nonmammalian species have a third high-affinity melatonin receptor subtype, the Mel_{1c} receptor (31). The mammalian Mel_{1a} and Mel_{1b} receptors are also called the MT1 and MT2 receptors, respectively [10]. The inability of this MT nomenclature system to accommodate the nonmammalian Mel_{1c} receptor has resulted in persistence of dual nomenclature systems; here, we use the original nomenclature.

The relative importance of the mammalian melatonin receptor subtypes in mediating circadian responses to the hormone is unclear. The circadian clocks of neonatal hamsters are very efficiently set by even a single injection of melatonin, and several physiological and in vitro responses of the suprachiasmatic nucleus (SCN) to melatonin have been observed in hamsters (25, 36, 39, 47). Remarkably, however, the Mel_{1b} receptor gene of several hamster species, including those with robust circadian and reproductive responses to the hormone, does not encode a functional melatonin receptor (47; see also GenBank accession number AY145849). Pharmacological studies with mice, however, suggest that the mouse Mel_{1b} receptor mediates circadian responses (9, 19). Other data, derived from mice with targeted disruption of the Mel_{1a} melatonin receptor, indicate that this subtype mediates an acute suppressive effect of melatonin on SCN neuronal firing, and they reveal apparent redundancy of receptor subtypes in mediating the phase shifting response to melatonin (23).

To determine whether the mouse Mel_{1b} receptor contributes

* Corresponding author. Present address: Department of Neurobiology, Lazare Research Building, Room 723, University of Massachusetts Medical School, 364 Plantation St., Worcester, MA 01605-2324. Phone: (508) 856-2495. Fax: (508) 856-6266. E-mail: david.weaver@umassmed.edu.

† Present address: CombinatoRx Inc., Boston, MA 02118.

‡ Present address: Department of Neurobiology, University of Massachusetts Medical School, Worcester MA 01605-2324.




FIG. 1. Mel_{1b} receptor gene, targeting construct, and genotyping strategies. (A) Schematic of the Mel_{1b} receptor gene and the targeting construct. A neomycin resistance gene driven by the phosphoglycerate kinase promoter (PGK-Neo cassette) was inserted in reverse orientation at the *SmaI* site within exon 1 of the mouse Mel_{1b} receptor gene. The 5' and 3' arms of the targeting construct were 2.2 and 7 kb, respectively. (B) Genotyping by Southern blot. A *Bam*HI restriction site introduced with the PGK-Neo cassette formed the basis for genotyping by Southern blotting, using a probe (solid bar in panel A) located outside the 5' end of the construct. Following digestion with *Bam*HI, the probe hybridizes to a ca. 10-kb band in wild-type DNA and a ca. 7-kb band for the targeted allele. Genotypes are shown above the lanes. Lanes illustrate the hybridization pattern of wild-type (W), heterozygous mutant (H), and homozygous Mel_{1b} -receptor mutant (knockout [K]) mice. Approximate sizes of the genomic fragments are indicated at the left. (C) Strategy for genotyping by PCR. Magnified view of the relationship of the primers (arrows) to the nucleotide sequences of the Mel_{1b} receptor gene and the PGK-Neo cassette. (D) Genotyping by PCR. Amplification of mouse genomic DNA with a cocktail of three primers led to amplification of distinct bands representing the wild-type and targeted alleles. Genotypes are shown above the lanes; designations are as in panel B. Size markers on the left indicate bands at 910, 540, 426, 409, 266, and 166 bp, generated by pUC19 and digested with *Dde*I.

to responses to the hormone, the mouse Mel_{1b} receptor cDNA was isolated. Finding that the mouse Mel_{1b} receptor cDNA encodes a high-affinity melatonin receptor, we then disrupted the Mel_{1b} receptor gene. Analysis of the mutant mice indicates that the Mel_{1b} receptor encodes a functionally relevant melatonin receptor. The Mel_{1b} receptor appears to mediate a response to the hormone occurring at higher ligand concentrations in mice lacking the Mel_{1a} melatonin receptor.

MATERIALS AND METHODS

Isolation and characterization of the mouse Mel_{1b} receptor cDNA. A fragment of the mouse Mel_{1b} receptor cDNA isolated by PCR has been previously reported (47). A probe based on this published sequence was used to screen mouse genomic DNA libraries, resulting in isolation of two overlapping genomic clones containing exon 2 (A341 and A351). Genomic DNA encoding exon 1 was isolated from a bacterial artificial chromosome library (Research Genetics, Inc.) screened with the most 5' end of the intronic sequence from clone A341. In addition, PCR amplification of mouse brain cDNA was performed using primers spanning the intron, to confirm the splicing pattern, as well as along the cDNA to confirm the sequence. A full-length cDNA was isolated by PCR using an overlap-PCR method to "stitch" together exons 1 and 2, producing a silent mutation at the splice site. The integrity of the cDNA construct was confirmed by sequence analysis and compared to the sequence of genomic clones.

Genomic DNA sequence corresponding to the mouse Mel_{1b} receptor locus has recently been deposited in GenBank. GenBank accession no. NW_000351 represents a genomic contig including the mouse Mel_{1b} receptor locus. The deduced amino acid sequence of the receptor cDNA (GenBank accession no. XM_146818), based on inferred splicing patterns using GenomeScan, differs from that determined above due to the detection of two apparent exons within intron 1, disruption of exon 2 by an apparent intron, and splicing to an additional

exon just 5' of the actual stop codon. Both homology with other members of the melatonin receptor gene family and the RT-PCR data support the conclusion that the Mel_{1b} receptor locus consists of two exons.

When the nucleotide sequences are translated using a two-exon model, the deduced amino acid sequence of the Mel_{1b} receptor cDNA reported in GenBank (AAL85489 and NW_000351) still differs from our sequence, due to the apparent insertion of two nucleotides in exon 2 in the portion of sequence encoding the carboxyl terminus of the protein in the GenBank entry. The accuracy of the sequence reported here is supported by homology of our deduced amino acid sequence with the human and Northern pike Mel_{1b} receptor sequences (12) and is confirmed by the identity of our sequence with the sequence entry in the Celera database (entries mCT7575 and mCG8325). (The only amino acid difference between the sequence reported here and in the Celera database is T237 of our sequence, which is E237 in the Celera and other sequences.) We cannot exclude the possibility that strain differences or other polymorphisms in the genetic sequence exist.

Ligand binding characteristics. To compare the affinity and specificity of the mouse Mel_{1a} and Mel_{1b} receptors, the characteristics of these receptor cDNAs were examined in parallel in transiently transfected COS7 cells. Ligand binding studies were conducted using the radioligand 2-[125 I]iodomelatonin (125 I-MEL) as previously described (29, 34). Saturation binding experiments were conducted with ligand concentrations ranging from 10 to 1,800 pM. Data were analyzed using Kaledograph software. Competitive inhibition studies were conducted using 125 I-MEL concentrations of 110 to 150 pM. K_i values were calculated from 50% inhibitory concentration (IC_{50}) values using the Cheng-Prusoff equation (5): $K_i = IC_{50} / (1 + ([\text{ligand}] / K_d))$.

Targeting disruption of the Mel_{1b} receptor gene. A targeting construct was generated in which the first exon of the Mel_{1b} receptor gene was disrupted by insertion of a neomycin resistance cassette at a *Sma*I site in exon 1 (Fig. 1). The cassette (PGK-Neo) consists of the phosphoglycerate kinase (PGK) promoter and a neomycin resistance gene and was placed in reverse orientation. The 5' arm was 2.2 kb; the 3' arm was 7 kb. The targeting construct was introduced into

J1 embryonic stem cells by electroporation at the Massachusetts General Hospital Knockout Core facility (En Li, director). For screening, DNA from neomycin-resistant colonies was digested with *Bam*HI and assessed by Southern blotting, using a nonoverlapping 0.8-kb *Ssr*I fragment flanking the 5' end of the construct as a probe. The targeting event was confirmed by digestion of embryonic stem cell DNA with *Eco*RI, followed by hybridization with a nonoverlapping probe (based on exon 2) from the 3' flanking region as a probe (data not shown). A clone with the appropriate banding pattern on Southern blot analysis (#83) was microinjected into blastocysts by the Knockout Core facility. Chimeric mice were bred to females of strain 129/sv, provided by En Li. Heterozygous offspring were bred together to generate isogenic *Mel_{1b}* receptor-deficient and wild-type lines on the 129/sv background.

Southern blot analysis of tail DNA confirmed the targeting event (Fig. 1B). For colony maintenance, mice were genotyped using PCR (Fig. 1C and D). Reactions were performed using a mixture of three primers, allowing detection of both wild-type and targeted alleles in a single reaction (Fig. 1C). A common forward primer in exon 1 (mus1b5'-4, CCAGGCCCTGTGACTGCCCGGG) and a gene-specific reverse primer from intron 1 (mus1b3'-3, 5' CTGCCACT GAGGACAGAACAGG 3') amplified a 272-bp fragment from the wild-type allele. The common forward primer and a reverse primer based on the sequence of the 3' end of the PGK-Neo cassette (neo6-2, TGCCCCAAAGGCTACC CGCTTCC) amplified a ca. 550-bp fragment from the targeted allele. Amplification of genomic DNA extracted from tail biopsies was performed using *Taq* DNA polymerase and a program of 3 min at 94°C for initial denaturation, followed by 35 cycles of denaturation at 94°C (30 s), annealing at 60°C (30 s), and extension at 72°C (1 min). Products were separated on agarose gels and visualized with ethidium bromide using UV trans-illumination (Fig. 1D).

Generation of melatonin receptor-deficient lines on a C3H/He genetic background. Mice with targeted disruption of the *Mel_{1a}* receptor have been previously described (23). In these mice, exon 1 of the *Mel_{1a}* receptor gene has been replaced with the PGK-neomycin cassette. To generate animals with the receptor targeting events on a consistent genetic background, a *Mel_{1a}* receptor-deficient male mouse with a hybrid, 129/sv × C57BL/6 genetic background was backcrossed to a C3H/He strain female (C3H, Charles River Laboratories). The C3H strain was used because this strain has been used extensively by others in examining circadian behavioral responses to melatonin (2, 3, 9), and this strain has rhythmic melatonin production (unlike most other inbred strains of mice; see reference 11). In each generation, heterozygous males were selected and crossed to C3H females. After backcrossing was done for 10 generations, heterozygous male and female mice were interbred to produce wild-type and *Mel_{1a}* receptor-deficient lines which produce rhythmic melatonin as previously reported (42).

Similarly, a male *Mel_{1b}* receptor-deficient mouse (on the isogenic, 129/sv genetic background) was bred to a C3H female. Heterozygous male offspring from each generation were backcrossed to C3H females for six generations of backcrossing, at which time male and female heterozygotes were interbred to produce wild-type and *Mel_{1b}* receptor-deficient lines on the C3H background. Finally, *Mel_{1a}* receptor-deficient mice on the C3H genetic background were crossed with the *Mel_{1b}* receptor-deficient mice on the C3H background to generate *Mel_{1a}/Mel_{1b}* double-mutant mice on a C3H genetic background.

Multiunit recordings. Multiunit recordings of SCN electrical activity were performed as previously described (14, 23). Briefly, hypothalamic slices were prepared from adult male mice maintained in a 12-h light:12-h dark lighting cycle. Animals were euthanized 2 to 5 h after lights-on. Brains were rapidly dissected and placed into artificial cerebrospinal fluid medium (ACSF) containing 116.3 mM NaCl, 5.4 mM KCl, 1.0 mM NaH₂PO₄, 26.2 mM NaHCO₃, 1.8 mM CaCl₂, 0.8 mM MgSO₄, 24.6 mM dextrose, and 5-mg/liter gentamicin sulfate (pH 7.5). Slices (400 μm thick) were prepared using a tissue chopper. Electrical activity was recorded using a Teflon-coated platinum-iridium wire electrode as previously described (14, 23).

Experiments were performed on the second day in vitro, >24 h after preparation of the slices. Slices were exposed to vehicle (0.1% ethanol in ACSF) and then to escalating concentrations of melatonin (0.1 to 100 nM) in ACSF containing a maximum of 0.1% ethanol. Data are expressed as percent change in electrical activity at the 100-nM dose. Experiments were conducted without knowledge of the genotypes of the animals. These studies were conducted on wild-type and homozygous *Mel_{1b}*-mutant mice on the 129/sv genetic background.

Cyclic AMP response element binding protein (CREB) phosphorylation. Pituitary adenylyl cyclase activating peptide (PACAP) is a transmitter of the retinohypothalamic tract, the pathway that conveys light-dark information from the retina to the SCN (17). PACAP appears to modulate the effects of glutamate on the SCN (17, 40). In vitro application of PACAP to SCN slices induces phosphorylation of CREB on serine residue 133 (20, 40, 41). Phosphorylated CREB (pCREB) is thought to play an important role in the photic regulation of

gene expression in the SCN, including the light-induced expression of *c-fos* and the *mPeriod* genes (14, 38, 40). Melatonin reduces PACAP-induced pCREB immunoreactivity in SCN slices in a dose-dependent manner (20, 40, 41, 42). Melatonin may thus subtly modulate photic sensitivity of the SCN circadian clock.

Melatonin inhibition of the PACAP-induced CREB phosphorylation state was studied as previously described (41). Briefly, 400-μm coronal brain slices containing the SCN were collected 2 to 6 h after lights-on and maintained in oxygenated medium (145 mM NaCl, 5 mM KCl, 1.8 mM CaCl₂, 0.8 mM MgCl₂, 10 mM HEPES, and 10 mM glucose [pH 7.35]). At the time corresponding to Zeitgeber Time 10 (ZT10, e.g., 10 h after lights-on in the colony room), slices were treated with PACAP (100 nM) or vehicle. Slices were pretreated with melatonin (1 or 100 nM) or vehicle for 15 min prior to PACAP application. Fifteen minutes after PACAP application, slices were fixed with 4% paraformaldehyde. Slices were fixed for 12 to 16 h and then cryo-protected (20% sucrose in phosphate-buffered saline) and sectioned at 14-μm thickness on a cryostat. Sections were mounted on gelatin-coated slides and stored at -20°C until processed for immunohistochemical detection of CREB phosphorylated at serine 133 (pCREB). pCREB was detected by use of a commercially available, polyclonal antibody (Upstate Biotechnology, Lake Placid, N.Y.) at a dilution of 1:1,000. Immunoreaction was visualized with a standard avidin-biotin labeling method (Vector Labs, Burlingame, Calif.) with diaminobenzidine as a chromogen.

Semiquantitative analysis was performed as previously described (41). Briefly, images were digitized (VIDAS, Kontron, Germany), and background staining was defined as the lower threshold. Within the area of the SCN, all cell nuclei showing a pCREB immunoreaction exceeding the threshold were marked. Immunoreactive nuclei were then counted in three sections per animal in a blind manner. The average of six values (number of immunoreactive nuclei per unilateral SCN per 14-μm section) was calculated for each animal.

It is not known whether melatonin alters the number of nuclei immunoreactive for pCREB by a reduction in the rate of phosphorylation of CREB or by activation of a phosphatase. In this report, "CREB phosphorylation" is used to indicate the static state of CREB phosphorylation (e.g., density of pCREB-immunoreactive nuclei), not the rate at which CREB is phosphorylated.

Nucleotide sequence accession number. The cDNA sequence for the mouse *Mel_{1b}* receptor that was determined in this study has been deposited in GenBank (accession number AY145850).

RESULTS

Isolation and characterization of the mouse *Mel_{1b}* receptor cDNA. The mouse *Mel_{1b}* receptor cDNA sequence was determined from genomic DNA and confirmed by amplification of the receptor cDNA from brain RNA (see Materials and Methods for discussion of discrepancies between our sequence and other database entries).

The deduced amino acid sequence of the *Mel_{1b}* receptor has features shared by other members of the melatonin receptor subfamily of G protein-coupled receptors (Fig. 2). Specifically, the amino acid sequence of the mouse *Mel_{1b}* receptor is highly related to the human *Mel_{1b}* receptor, with which it shares 79% identity. The other closely related sequences identified by BLAST searches of the GenBank database are fragments of the *Mel_{1b}* receptor cDNA from other species, including the Northern pike *Mel_{1b}* receptor cDNA (12). Amino acid identity of the mouse *Mel_{1a}* and *Mel_{1b}* receptors is 50.8%.

The mouse *Mel_{1b}* receptor sequence also has several features conserved in the melatonin receptor family within the superfamily of G protein-coupled receptors. A highly conserved sequence just downstream from the third transmembrane domain (NRYC[C/Y]ICHS) is distinctive; most other receptor families have DRY or ERY in place of NRY (33, 45). In transmembrane domain 7, the melatonin receptor family shares a characteristic motif (NA[I/V][I/V]Y) rather than NPXXY. The presence of potential a N-linked glycosylation site in the amino-terminal, putative extracellular domain of the

```

MOUSE 1B  MPENSSIPNCCSEAGSLAARPSWSGSAGARPPVTARAPWVAPMLSTVVVTTAVDFVGNLLVILSVLRNRKLRNAGNLFVVSALADLVIALYYPYPLILVA
HUMAN 1B  MSENSFPANCCSEAGGWAIRPGWSGAGSARPSRTPRPVWAPALSAVLIIVTTAVDVVGNLLVILSVLRNRKLRNAGNLFVVSALADLVVAFYYPYPLILVA
MOUSE 1A  MKGNVSELLNATQQAPGGEGGRPRPSWLASTLAFILIFTIIVVDILGNLLVILSVYRNKLRNAGNIFVVSALADLVVAVYYPYPLVLTLS
HUMAN 1A  MQQNGSALPNASQ---PVLRGD GARPSWLASALACVLIFTIIVVDILGNLLVILSVYRNKLRNAGNIFVVSALADLVVAIYYPYPLVLTMS
Consensus  -----RP-W-A--L-----T--VD--GNLLVILSV--RN--KLRN--GN--F-VSLA--ADLV--A--YYPYPL--L--

MOUSE 1B  IIRDGWWLGEAHCKASAFV MGLSVIGSVFNITAIAINRYCCICHSTTYHRVCSHWYTPYISLVWLLTLVALVFNFFVGSLEYDPRIYSCFTFIQTASTQY
HUMAN 1B  IFYDGWALGEBHCKASAFV MGLSVIGSVFNITAIAINRYCYICHSMAYHRIYRRWHTPLHICLIWLLTVALLPNFFVGSLEYDPRIYSCFTFIQTASTQY
MOUSE 1A  ILNNGWNLGYLHCQVSAFLMGLSVIGSVFNITGIAMNRYCYICHSLKYDKIYSNKNSLCYVFLIWMLLTIAIMPNLQTGTLQYDPRIYSCFTFTQSVSSAY
HUMAN 1A  IFNNGWNLGYLHCQVSGFLMGLSVIGSVFNITGIAINRYCYICHSLKYDKLYSKNSLCYVLLIWLTLAVALPNLRAGTLQYDPRIYSCFTFAQSVSSAY
Consensus  I---GW-LG--HC--S-L-MGLSVIGS--FNIT--IA--NRYC--ICHS--Y-----W-LT--A--PN---G-L-YDPRIYSCFTF-Q--S--Y

MOUSE 1B  TAAVVAIHFLLPMAVVSFCYLRIWVVLVQARRKAKATRKLRRLRPSDLRSFLTMFAVVFVFAICWAPLNCIGLAVAINPEAMALQVPEGLFVTSYFLAYFN
HUMAN 1B  TAAVVVHFLLPPIAVVSFCYLRIWVVLVQARRKAKPESRLCKLPSDLRSFLTMFVVVFVFAICWAPLNCIGLAVAINPEAMAPQIPEGLFVTSYLLAYFN
MOUSE 1A  TIAVVVFHFIVPMIIVIFCYLRIWVVLVQVRRRVKPDNKPKLKPQDFRNFVTFMVVFLFAICWAPLNLIGLIVASDPATMVPRIPFWLFWASYFLAYFN
HUMAN 1A  TIAVVVFHFIVPMIIVIFCYLRIWVVLVQVRRRVKPDNRKPKLKPQDFRNFVTFMVVFLFAICWAPLNFIFGLAVASDPASMPVRIPEWLFVASYMAYFN
Consensus  T-AVV--HF--P---V-YCYLRIW-LVLQ-R---K-----L-P---R-F-TMF-VFVFAICWAPLN-IGL-VA--P--M----PE-LFV-SY--AYFN

MOUSE 1B  SCLNAIVYGLLNQNFREYKRILLAIWNTRRCIQHASKHCLTEERQGPPTPAARATVPVKEGAL 364
HUMAN 1B  SCLNAIVYGLLNQNFREYKRILLALWNRPHCIQDASKGSHAEGLSQPAPPI--IGVQHQADAL 362
MOUSE 1A  SCLNAIYGLLNQNFREYKIIIVSLCTAKMPFVSSNEEADKICKKPSLIP--NNNLIKVDSV 353
HUMAN 1A  SCLNAIYGLLNQNFREYRRIIVSLCTARVFFVDSNDVADRKKPKSPLMT--NNNVVKVDSV 350
Consensus  SCLNAI-YGLLNQNFREY--I-----S-----P-----
    
```

FIG. 2. The mouse Mel_{1b} receptor sequence. The sequence of the mouse Mel_{1b} receptor has been aligned with other members of the melatonin receptor gene family: mouse Mel_{1b} (present work), human Mel_{1b} (30), mouse Mel_{1a} (34) (NP_032665), and human Mel_{1a} receptor (29) (NP_005949). Putative transmembrane domains are indicated by underlining in the sequence for the mouse Mel_{1b} receptor. Gaps introduced to optimize alignment are indicated by dashes. Consensus sequence is defined as residues that are identical in all four of the sequences shown.

protein is also a common feature of the melatonin receptor family. Finally, with respect to gene structure, the location of the splice site between exons 1 and 2 is also conserved among all mammalian melatonin receptor genes that have been examined (32, 34, 47).

To determine the relative affinity and specificity of mouse melatonin receptors, the characteristics of the mouse Mel_{1a} and Mel_{1b} receptor cDNAs were examined in parallel. ¹²⁵I-MEL binding to transfected COS7 cells revealed that the affinity of the mouse Mel_{1b} receptor is significantly lower than the affinity of the mouse Mel_{1a} receptor (Mel_{1b}, K_d (mean ± standard error of the mean [SEM]) = 199.9 ± 13.4 pM [n = 4]; Mel_{1a}, K_D = 59.3 ± 18.3 pM [n = 2]). Competition experiments with several drugs revealed that the rank order of potency was as expected for a member of the melatonin receptor gene family, e.g., 2-iodomelatonin > melatonin > 6-chloromelatonin >> N-acetylserotonin (Table 1). The relative affinity of the receptor subtypes as well as the rank order of drug potency for inhibiting ¹²⁵I-MEL binding are similar to the characteristics of the human Mel_{1a} and Mel_{1b} receptors (30).

It is noteworthy that the mouse Mel_{1b} receptor encodes a functional receptor, as inferred from its sequence and confirmed in radioreceptor assays. This is in contrast to the Mel_{1b} receptor gene in several hamster species, including Siberian and Syrian hamsters, in which nonsense mutations disrupt the receptor cDNA (see reference 47 and GenBank accession number AY145849).

Targeted disruption of the mouse Mel_{1b} receptor gene. Mice with targeted disruption of the Mel_{1b} receptor gene are viable and fertile. This is not unexpected; most inbred strains of mice do not produce melatonin as a result of genetic defects in the synthetic enzymes (11, 35), and removal of the pineal gland, the source of circulating melatonin, does not have major physiological effects (except in species in which melatonin is involved in the seasonal regulation of reproduction).

Observations of locomotor activity rhythms of Mel_{1b} receptor-deficient mice and wild-type controls placed in constant

darkness revealed no robust circadian phenotype; e.g., activity rhythms persisted in homozygous Mel_{1b} receptor-deficient mice in constant conditions, with a period length similar to that observed in controls (data not shown). Preliminary attempts to use daily or repeated daily injections of melatonin to phase shift the locomotor activity rhythm of wild-type mice were unsuccessful, preventing us from employing an in vivo assay of melatonin receptor function. Instead, in vitro assays of melatonin receptor function were employed.

Melatonin inhibits multiunit activity via the Mel_{1a} receptor. Inhibition of SCN multiunit activity by melatonin was unaffected in mice with targeted disruption of the Mel_{1b} receptor (Fig. 3). In contrast, the acute suppressive effect of melatonin was absent in mice homozygous for targeted disruption of the Mel_{1a} receptor (23; [data replotted for comparison in Fig. 3]). Thus, the Mel_{1a} receptor alone appears to mediate the suppressive effect of melatonin on SCN neuronal firing rate.

Both melatonin receptor subtypes contribute to regulation of CREB phosphorylation. Previous studies have shown that melatonin reduces the levels of PACAP-induced CREB phosphorylation at low concentrations (20, 40, 41). In wild-type mice, physiological concentrations of melatonin (≤1 nM) inhibit the level of phosphorylated CREB by >70%. In mice with targeted disruption of the Mel_{1a} melatonin receptor, the re-

TABLE 1. Inhibition constants for mouse Mel_{1a} and Mel_{1b} receptor cDNAs^a

Compound	K _i (no. of experiments) for:	
	Mel _{1a}	Mel _{1b}
2-Iodomelatonin	0.045 ± 0.005 (4)	0.159 ± .017 (4)
Melatonin	0.438 ± 0.169 (3)	2.853 ± 0.665 (3)
6-Chloromelatonin	4.053 ± 1.371 (3)	8.903 ± 1.640 (3)
N-acetylserotonin	638 (1)	296 (1)

^a Inhibition constants (K_i values) are expressed as means ± SEM, in nanomolar concentrations.




FIG. 3. Inhibition of SCN multiunit activity by melatonin. Inhibition of SCN electrical activity by melatonin (0.1 to 100 nM) was assessed during midday. Data are plotted as the percent inhibition of electrical activity observed at the 100 nM dose relative to the level of electrical activity from the same slice exposed to vehicle. The left set of bars shows that the inhibition of SCN electrical activity is absent in Mel_{1a} receptor-deficient (-/-) mice (data replotted from reference 23; sample sizes 7 to 11 per group). Right bars show inhibition of SCN activity in experiments conducted for this study, using Mel_{1b} receptor-deficient mice (-/-) ($n = 6$) and wild-type control (+/+) ($n = 11$) mice. The inhibition of electrical activity in each group was significant ($P < 0.0001$, one-sample t tests, versus 0% inhibition), and there was no significant difference between the Mel_{1b} -/- and Mel_{1b} +/+ groups (t test, $P = 0.2$).

sponse to 1 nM melatonin is absent, but a residual effect of melatonin is apparent at higher melatonin concentrations (Fig. 4A, consistent with previously published results [41]). This higher-threshold response to melatonin appears to be mediated by the Mel_{1b} receptor, as the Mel_{1b}-preferring antagonist, 4P-PDOT (8), blocks the high-threshold response (41). In mice with targeted disruption of both Mel_{1a} and Mel_{1b} receptor genes, the residual response to high concentrations of melatonin is completely absent (Fig. 4B). These data indicate a dual regulation of the CREB phosphorylation state in the SCN: the Mel_{1a} receptor mediates responses to low concentrations of the hormone, while the residual response at higher concentrations is mediated by the Mel_{1b} receptor. The presence of a functional Mel_{1a} receptor that would mediate low-threshold responses precluded assessing the effect of disruption of the Mel_{1b} receptor alone.

DISCUSSION

Melatonin has subtle actions on circadian rhythmicity in mice and in other mammals (46). Mice with targeted disruption of either Mel_{1a} or Mel_{1b} receptor subtype, or the double mutants, do not have a striking overt phenotype. Examination of melatonin receptor-mediated responses does reveal a functional effect of each of the receptor subtypes, however. The Mel_{1a} receptor is necessary for the suppressive effect of melatonin on SCN neuronal firing (23). Targeted disruption of the Mel_{1b} receptor did not alter this response to melatonin. With respect to inhibition of the PACAP-induced CREB phosphorylation state, melatonin acts through the Mel_{1a} receptor at low




FIG. 4. Inhibition of CREB phosphorylation levels by melatonin. Semiquantitative densitometric analysis shows that melatonin reduces the number of pCREB-immunoreactive nuclei in PACAP-treated SCN slices. Values represent labeled nuclei per unilateral SCN and are the mean \pm SEM for four to five animals per group. In wild-type mice, melatonin (1 nM) reduces PACAP-induced CREB phosphorylation by $>70\%$ (41). In contrast, a 1 nM concentration had no significant effect on mice with targeted disruption of the Mel_{1a} melatonin receptor (Mel_{1a} -/-) (A). At higher melatonin concentrations (100 nM), a more complete suppression occurs even in mice with disruption of the Mel_{1a} receptor subtype (A) (far right). This residual, higher-threshold response is absent in mice with targeted disruption of both the Mel_{1a} and Mel_{1b} receptors (B). + + +, significant difference between vehicle control and PACAP treatment ($P < 0.002$, t test); **, significant difference versus PACAP group ($P < 0.01$, Dunnett's test); NS, not significantly different from PACAP group ($P > 0.05$, Dunnett's test).

concentrations of the hormone (41) (present results). An effect of higher concentrations of melatonin is apparent in Mel_{1a} receptor-deficient mice, and this response is blocked by treatment with the Mel_{1b} receptor antagonist, 4P-PDOT (41). Consistent with this dual regulation, we report here that the SCN of Mel_{1a}-Mel_{1b} double-mutant mice is completely unresponsive to melatonin, even at the higher (100 nM) melatonin concentrations.

One of the *in vitro* assays used previously to assess the effects of melatonin on the circadian clock involves monitoring the effect of melatonin on the phase of single-unit firing rate rhythms in SCN slices (19, 23, 26, 47). Liu et al. (23) found that melatonin and 2-iodomelatonin phase-shifted the circadian

rhythm of SCN single-unit activity even in mice with targeted disruption of the Mel_{1a} receptor, although the magnitude of response was lower in the Mel_{1a} receptor-deficient mice at low ligand concentrations. The response persisting in Mel_{1a} receptor-deficient mice was mediated by a pertussis toxin-sensitive mechanism, indirectly implicating Mel_{1b} receptors as mediating the higher-threshold response. More recently, Hunt et al. (19) reported that the Mel_{1b} receptor subtype mediates the phase-shifting effect of melatonin in the rat SCN, based on studies using subtype-selective antagonists. In contrast, SCN slices from hamsters lacking a functional Mel_{1b} receptor (due to a naturally occurring nonsense mutation) respond to melatonin with phase shifts in single-unit activity (47). These diverse results from the single-unit assay indicate either that there are pronounced species differences in the melatonin receptor subtype mediating the phase shifting response or, more likely, that there is functional redundancy between the receptor subtypes.

While this assay system has been used successfully by our associates and others in the past, in preliminary studies we were unable to obtain reliable single-unit activity rhythms, even with commercially reared wild-type mice, at the time we attempted to examine the responses of Mel_{1b} receptor-deficient mice (C. Liu and S. M. Reppert, unpublished data). In the absence of a robust rhythm with clearly defined phase, assessment of the impact of melatonin on circadian phase was not possible.

We were also unable, in preliminary experiments, to demonstrate a consistent effect of melatonin on locomotor activity rhythms *in vivo*, even in wild-type mice with the C3H genetic background. This strain has previously been reported to exhibit phase shifts in response to melatonin treatment (2, 3, 9). The mean phase shift occurring in the paradigm described by Dubocovich and colleagues is a less than 1.0-h total shift with daily injections of melatonin for 3 days. Behavioral arousal (such as would occur coincident with handling the animal for injection) has an effect on the rodent circadian system that appears to be as robust as the effect of melatonin (18, 28). In our hands, melatonin injection paradigms did not elicit consistently detectable phase shifts in mice. We were thus unable to use the mutant mice to directly test the hypothesis (9) that the Mel_{1b} receptor mediates behavioral phase shifts.

Data from other systems suggest that both Mel_{1a} and Mel_{1b} receptors contribute to responses to the hormone. Vascular responses to melatonin occurring at low concentrations are mediated primarily by the Mel_{1a} receptor, with a Mel_{1b}-mediated response occurring at higher concentrations (6, 37). This directly parallels our findings regarding the dose-related inhibition of PACAP-induced CREB phosphorylation in the SCN. The acute suppressive effect of melatonin on SCN neuronal firing is mediated by the Mel_{1a} receptor. The Mel_{1b} receptor appears to mediate inhibition of firing in the rat hippocampus, which, unlike the mouse hippocampus, has a significant level of melatonin receptor binding (44). A melatonin-regulated rhythm in *mPeriod1* gene expression in the pars tuberalis of the pituitary is abolished by removal of the pineal gland in hamsters (27); the absence of a functional Mel_{1b} melatonin receptor gene in this species implicates the Mel_{1a} receptor. Consistent with this proposal, pinealectomy or targeted disruption of the Mel_{1a} receptor gene in mice that have rhythmic

pineal melatonin synthesis disrupts rhythmic *mPer1* expression in the pars tuberalis (42). These findings suggest that the expression of melatonin receptor subtypes varies by tissue and indicate the possibility of functional redundancy when the receptor subtypes are coexpressed.

In summary, targeted disruption of the Mel_{1b} receptor does not have a profound impact on the circadian system of mice. Our data from examining melatonin inhibition of the level of CREB phosphorylation in PACAP-treated SCN slices indicate that the Mel_{1a} receptor mediates the majority of this response to the hormone. At higher melatonin concentrations, a residual impact of melatonin, acting via the Mel_{1b} receptor, is apparent. This difference in the relative contribution of the receptor subtypes is likely not due to differences in the affinity of the receptors *per se*, but instead may derive from differences in the levels of receptor expression directed by the two genes or by differences in coupling to second messenger pathways. The Mel_{1a} receptor accounts for all the melatonin receptor binding detected in mouse brain by *in vitro* autoradiography (23). The Mel_{1a} receptor, being much more highly expressed, may mediate most responses to the hormone in mice. The Mel_{1b} receptor can also contribute to responses in those tissues in which it is expressed. The availability of mice with targeted disruption of Mel_{1a}, Mel_{1b}, and both receptor subtypes will allow a genetic approach to dissection of additional melatonin receptor-mediated responses.

ACKNOWLEDGMENTS

We thank Chen Liu, Donald Hodges, and Stefanie Rassnick for assistance with preliminary studies and Camala Capodice and Christopher Lambert for technical assistance.

This work was supported by grants from the NIH (DK42125 to S.M.R. and AG09975 to D.R.W.), the Deutsche Forschungsgemeinschaft (to J.H.S and C.V.G.), the Paul und Ursula Klein-Stiftung and the Heinrich und Fritz Reise-Stiftung (to J.H.S), and a sponsored research agreement from Bristol-Myers Squibb. X.J. was supported in part by NIH Postdoctoral fellowship F32 MH12067. C.V.G. was supported in part by the Emmy-Noether Programm of the Deutsche Forschungsgemeinschaft.

REFERENCES

- Bartness, T. J., E. L. Bittman, M. H. Hastings, J. B. Powers, and B. D. Goldman. 1993. Timed melatonin infusion paradigm for melatonin delivery: what has it taught us about the melatonin signal, its reception, and the photoperiodic control of seasonal responses? *J. Pineal Res.* **15**:161–190.
- Benloucif, S., M. I. Masana, K. Yun, and M. L. Dubocovich. 1999. Interactions between light and melatonin on the circadian clock of mice. *J. Biol. Rhythms* **14**:281–289.
- Benloucif, S., and M. L. Dubocovich. 1996. Melatonin and light induce phase shifts of circadian activity rhythms in the C3H/HeN mouse. *J. Biol. Rhythms* **11**:113–125.
- Cassone, V. M. 1990. Effects of melatonin on vertebrate circadian systems. *Trends Neurosci.* **13**:457–464.
- Cheng, Y., and W. H. Prusoff. 1973. Relationship between the inhibition constant (K_i) and the concentration of inhibitor which causes 50 per cent inhibition (I_{50}) of an enzymatic reaction. *Biochem. Pharmacol.* **22**:3099–3108.
- Doolen, S., D. N. Krause, M. L. Dubocovich, and S. P. Duckles. 1998. Melatonin mediates two distinct responses in vascular smooth muscle. *Eur. J. Pharmacol.* **345**:67–69.
- Drew, J. E., P. Barrett, L. M. Williams, S. Conway, and P. J. Morgan. 1998. The ovine melatonin-related receptor: cloning and preliminary distribution and binding studies. *J. Neuroendocrinol.* **10**:651–661.
- Dubocovich, M. L., M. I. Masana, S. Jacob, and D. M. Sauri. 1997. Melatonin receptor antagonists that differentiate between the human Mel_{1a} and Mel_{1b} recombinant subtypes are used to assess the pharmacological profile of the rabbit retina ML1 presynaptic heteroreceptor. *Naunyn-Schmiedeberg's Arch. Pharmacol.* **355**:365–375.
- Dubocovich, M. L., K. Yun, W. M. Al-Ghoul, S. Benloucif, and M. I. Masana.

1998. Selective MT₂ melatonin receptor antagonists block melatonin-mediated phase advances of circadian rhythms. *FASEB J.* **12**:1211–1220.
10. **Dubocovich, M. L., D. P. Cardinali, P. Delagrange, D. N. Krause, A. D. Strosberg, D. Sugden, and F. D. Yocca.** 2000. The IUPHAR compendium of receptor characterisation and classification, 2nd ed., p. 187–193. IUPHAR Media, London, United Kingdom.
 11. **Ebihara, S., T. Marks, D. J. Hudson, and M. Menaker.** 1986. Genetic control of melatonin synthesis in the pineal gland of the mouse. *Science* **231**:491–493.
 12. **Gaildrat, P., F. Becq, and J. Falcon.** 2002. First cloning and functional characterization of a melatonin receptor in fish brain: a novel one? *J. Pineal Res.* **32**:74–84.
 13. **Ganguly, S., S. L. Coon, and D. C. Klein.** 2002. Control of melatonin synthesis in the mammalian pineal gland: the critical role of serotonin acetylation. *Cell Tissue Res.* **309**:127–138.
 14. **Gribkoff, V. K., R. L. Pieschl, T. A. Wisialowski, A. N. van den Pol, and F. D. Yocca.** 1998. Phase shifting of circadian rhythms and depression of neuronal activity in the rat suprachiasmatic nucleus by neuropeptide Y: mediation by different receptor subtypes. *J. Neurosci.* **18**:3014–3022.
 15. **Ginty, D. D., J. M. Kornhauser, M. A. Thompson, H. Bading, K. E. Mayo, J. S. Takahashi, and M. E. Greenberg.** 1993. Regulation of CREB phosphorylation in the suprachiasmatic nucleus by light and a circadian clock. *Science* **260**:238–241.
 16. **Gubitza, A. K., and S. M. Reppert.** 1999. Assignment of the melatonin-related receptor to human chromosome X (GPR50) and mouse chromosome X (Gpr50). *Genomics* **55**:248–251.
 17. **Hannibal, J.** 2002. Neurotransmitters of the retinohypothalamic tract. *Cell Tissue Res.* **309**:73–88.
 18. **Hastings, M. H., S. M. Mead, R. R. Vindlacheruvu, F. J. Ebling, E. S. Maywood, and J. Grosse.** 1992. Non-photic phase shifting of the circadian activity rhythm of Syrian hamsters: the relative potency of arousal and melatonin. *Brain Res.* **591**:20–26.
 19. **Hunt, A. E., W. M. Al-Ghoul, M. U. Gillette, and M. L. Dubocovich.** 2001. Activation of MT₂ melatonin receptors in rat suprachiasmatic nucleus phase advances the circadian clock. *Am. J. Physiol.* **280**:C110–C118.
 20. **Kopp, M., H. Meissl, and H. W. Korf.** 1997. The pituitary adenylate cyclase-activating polypeptide-induced phosphorylation of the transcription factor CREB (cAMP response element binding protein) in the rat suprachiasmatic nucleus is inhibited by melatonin. *Neurosci. Lett.* **227**:145–148.
 21. **Lewy, A. J., S. Ahmed, J. M. Jackson, and R. L. Sack.** 1992. Melatonin shifts human circadian rhythms according to a phase-response curve. *Chronobiol. Int.* **9**:380–392.
 22. **Lewy, A. J., V. K. Bauer, B. P. Hasler, A. R. Kendall, M. L. Pires, and R. L. Sack.** 2001. Capturing the circadian rhythms of free-running blind people with 0.5 mg melatonin. *Brain Res.* **918**:96–100.
 23. **Liu, C., D. R. Weaver, X. Jin, L. P. Shearman, R. L. Pieschl, V. K. Gribkoff, and S. M. Reppert.** 1997. Molecular dissection of two distinct actions of melatonin on the suprachiasmatic circadian clock. *Neuron* **19**:91–102.
 24. **Malpoux, B., M. Migaud, H. Tricoire, and P. Chemineau.** 2001. Biology of mammalian photoperiodism and the critical role of the pineal gland and melatonin. *J. Biol. Rhythms* **16**:336–347.
 25. **Margraf, R. R., and G. R. Lynch.** 1993. Melatonin injections affect circadian behavior and SCN neurophysiology in Djungarian hamsters. *Am. J. Physiol.* **264**:R615–R621.
 26. **McArthur, A. J., A. E. Hunt, and M. U. Gillette.** 1997. Melatonin action and signal transduction in the rat suprachiasmatic circadian clock: activation of protein kinase C at dusk and dawn. *Endocrinology* **138**:627–634.
 27. **Messenger, S., M. L. Garabette, M. H. Hastings, and D. G. Hazlerigg.** 2001. Tissue-specific abolition of Per1 expression in the pars tuberalis by pinealectomy in the Syrian hamster. *NeuroReport* **12**:1–4.
 28. **Reebs, S. G., and N. Mrosovsky.** 1989. Effects of induced wheel running on the circadian activity rhythms of Syrian hamsters: entrainment and phase response curve. *J. Biol. Rhythms* **4**:39–48.
 29. **Reppert, S. M., D. R. Weaver, and T. Ebisawa.** 1994. Cloning and characterization of a mammalian melatonin receptor that mediates reproductive and circadian responses. *Neuron* **13**:1177–1185.
 30. **Reppert, S. M., C. Godson, C. D. Mahle, D. R. Weaver, S. A. Slangenaupt, and J. F. Gusella.** 1995. Molecular characterization of a second melatonin receptor expressed in human retina and brain: the Mel_{1b}-melatonin receptor. *Proc. Natl. Acad. Sci. USA* **92**:8734–8738.
 31. **Reppert, S. M., D. R. Weaver, V. M. Cassone, C. Godson, and L. F. Kola-kowski, Jr.** 1995. Melatonin receptors are for the birds: molecular analysis of two receptor subtypes differentially expressed in chick brain. *Neuron* **15**:1003–1015.
 32. **Reppert, S. M., D. R. Weaver, T. Ebisawa, C. D. Mahle, and L. F. Kola-kowski, Jr.** 1996. Cloning of a melatonin-related receptor from human pituitary. *FEBS Lett.* **386**:219–224.
 33. **Reppert, S. M., D. R. Weaver, and C. Godson.** 1996. Melatonin receptors step into the light: cloning and classification of subtypes. *Trends Pharmacol. Sci.* **17**:100–102.
 34. **Roca, A. L., C. Godson, D. R. Weaver, and S. M. Reppert.** 1997. Structure, characterization, and expression of the gene encoding the mouse Mel_{1a} melatonin receptor. *Endocrinology* **137**:3469–3477.
 35. **Roseboom, P. H., M. A. A. Nambodiri, D. B. Zimonjic, N. C. Popescu, I. R. Rodriguez, J. A. Gastel, and D. C. Klein.** 1998. Natural melatonin 'knock-down' in C57BL/6J mice: rare mechanism truncates serotonin N-acetyltransferase. *Mol. Brain Res.* **63**:189–197.
 36. **Rusak, B., and G. D. Yu.** 1993. Regulation of melatonin-sensitivity and firing-rate rhythms of hamster suprachiasmatic nucleus neurons: pinealectomy effects. *Brain Res.* **602**:200–204.
 37. **Ting, K. N., N. A. Blaylock, and D. Sugden.** 1999. Molecular and pharmacological evidence for MT₁ receptor subtypes in tail artery of juvenile Wistar rats. *Br. J. Pharmacol.* **127**:987–995.
 38. **Travnickova-Bendova, Z., N. Cermakian, S. M. Reppert, P. Sassone-Corsi.** 2002. Bimodal regulation of mPeriod promoters by CREB-dependent signaling and CLOCK/BMAL1 activity. *Proc. Natl. Acad. Sci. USA* **99**:7728–7733.
 39. **Viswanathan, N., and F. C. Davis.** 1997. Single prenatal injections of melatonin or the D1-dopamine receptor agonist SKF 38393 to pregnant hamsters sets the offspring's circadian rhythms to phases 180 degrees apart. *J. Comp. Physiol. [A]* **180**:339–346.
 40. **von Gall, C., G. Duffield, M. Hastings, M. D. Kopp, F. Dehghani, H. W. Korf, and J. H. Stehle.** 1998. CREB in the mouse SCN: a molecular integrator coding the phase adjusting stimuli of light, glutamate, PACAP and melatonin for clockwork access. *J. Neurosci.* **18**:10389–10397.
 41. **von Gall, C., D. R. Weaver, M. Kock, H. W. Korf, and J. H. Stehle.** 2000. Melatonin limits transcriptional impact of phosphoCREB in the mouse SCN via the Mel_{1a} receptor. *NeuroReport* **11**:1803–1807.
 42. **von Gall, C., M. L. Garabette, C. A. Kell, S. Frenzel, F. Dehghani, P. M. Schumm-Draeger, D. R. Weaver, H. W. Korf, M. H. Hastings, and J. H. Stehle.** 2002. Rhythmic gene expression in pituitary depends on heterologous sensitization by the neurohormone melatonin. *Nat. Neurosci.* **5**:234–238.
 43. **von Gall, C., J. H. Stehle, and D. R. Weaver.** 2002. Mammalian melatonin receptors: molecular biology and signal transduction. *Cell Tissue Res.* **309**:151–162.
 44. **Wan, Q., H. Y. Man, F. Liu, J. Braunton, N. B. Niznik, S. F. Pang, G. M. Brown, and Y. T. Wang.** 1999. Differential modulation of GABAA receptor function by Mel_{1a} and Mel_{1b} receptors. *Nat. Neurosci.* **2**:401–403.
 45. **Watson, S., and S. Arkininstall.** 1994. The G-protein linked receptor factbook. Academic Press, San Diego, Calif.
 46. **Weaver, D. R.** 1999. Melatonin and circadian rhythmicity in vertebrates. Physiological roles and pharmacological effects, p. 197–262 *In* F. W. Turek and P. C. Zee (ed.), Regulation of sleep and circadian rhythms. Marcel Dekker, Inc., New York, N.Y.
 47. **Weaver, D. R., C. Liu, and S. M. Reppert.** 1996. Nature's knockout: the Mel_{1b} receptor is not necessary for circadian or reproductive responses in Siberian hamsters. *Mol. Endocrinol.* **10**:1478–1487.