

3-11-2006

# Counter-selectable marker for bacterial-based interaction trap systems

Xiangdong Meng

*University of Massachusetts Medical School*

Robin M. Smith

*University of Massachusetts Medical School, robin.smith@umassmed.edu*

Astrid V. Giesecke

*Massachusetts General Hospital*

*See next page for additional authors*

Follow this and additional works at: [http://escholarship.umassmed.edu/pgfe\\_pp](http://escholarship.umassmed.edu/pgfe_pp)

 Part of the [Genetics and Genomics Commons](#)

## Repository Citation

Meng, Xiangdong; Smith, Robin M.; Giesecke, Astrid V.; Joung, J. Keith; and Wolfe, Scot A., "Counter-selectable marker for bacterial-based interaction trap systems" (2006). *Program in Gene Function and Expression Publications and Presentations*. 124.  
[http://escholarship.umassmed.edu/pgfe\\_pp/124](http://escholarship.umassmed.edu/pgfe_pp/124)

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in Program in Gene Function and Expression Publications and Presentations by an authorized administrator of eScholarship@UMMS. For more information, please contact [Lisa.Palmer@umassmed.edu](mailto:Lisa.Palmer@umassmed.edu).

---

# Counter-selectable marker for bacterial-based interaction trap systems

## **Authors**

Xiangdong Meng, Robin M. Smith, Astrid V. Giesecke, J. Keith Joung, and Scot A. Wolfe

## **Rights and Permissions**

Citation: *Biotechniques*. 2006 Feb;40(2):179-84. [Link to article on publisher's website](#)

# Counter-selectable marker for bacterial-based interaction trap systems

Xiangdong Meng<sup>1</sup>, Robin M. Smith<sup>1</sup>, Astrid V. Giesecke<sup>2</sup>, J. Keith Joung<sup>2</sup>, and Scot A. Wolfe<sup>1</sup>

<sup>1</sup>University of Massachusetts Medical School, Worcester and <sup>2</sup>Massachusetts General Hospital, Charlestown, MA, USA

*BioTechniques* 40:179-184 (February 2006)  
doi 10.2144/000112049

Counter-selectable markers can be used in two-hybrid systems to search libraries for a protein or compound that interferes with a macromolecular interaction or to identify macromolecules from a population that cannot mediate a particular interaction. In this report, we describe the adaptation of the yeast *URA3/5-FOA* counter-selection system for use in bacterial interaction trap experiments. Two different *URA3* reporter systems were developed that allow robust counter-selection: (i) a single copy *F'* episome reporter and (ii) a co-cis-trans *HIS3-URA3* reporter vector. The *HIS3-URA3* reporter can be used for either positive or negative selections in appropriate bacterial strains. These reagents extend the utility of the bacterial two-hybrid system as an alternative to its yeast-based counterpart.

## INTRODUCTION

Many in vivo interaction trap systems have been developed for the identification of protein-protein, protein-DNA, and protein-RNA interactions. These methods typically involve the use of a "bait" (a specific protein, DNA, or RNA sequence) to identify an interacting partner from a library of different "prey." These experiments can take the form of one-hybrid (1,2), two-hybrid (3), or three-hybrid (4) selections and can be performed in yeast (3,5) or bacteria (6–9). Interaction trap systems were originally developed for use in *Saccharomyces cerevisiae*, with interacting partners identified by positive selection. Positive selection is typically achieved by inactivating a gene, such as *HIS3*, required for survival under certain specific growth conditions, and then introducing a reporter system that drives expression of a functional copy of the deleted gene if a desired interaction occurs between the bait and prey (1). For *HIS3*, the stringency of the selection can be manipulated by varying the concentration of 3-amino triazole (3-AT), a competitive inhibitor of *HIS3*, in the media.

The *URA3* gene (orotidine-5'-phosphate decarboxylase) was origi-

nally developed as a counter-selectable marker for yeast (10,11). The endogenous *URA3* gene can be inactivated, and a reporter cassette introduced in which expression of *URA3* is dependent on the presence of a bait/prey interaction. If *URA3* is expressed, the uracil biosynthesis pathway will be functional, and 5-fluoro-orotic acid (5-FOA), which is added to the media, is metabolized into a suicide substrate for the essential thymidylate synthase enzyme. *URA3/5-FOA* systems have been used successfully in yeast interaction trap systems to counter-select against bait/prey interactions in vivo (5,12). The stringency of this counter-selection can be titrated by varying expression of the *URA3* gene; higher levels of *URA3* expression increase the sensitivity of a cell to 5-FOA (12).

*pyrF* is the *Escherichia coli* homolog of *URA3*, and deletion of *pyrF* can also be complemented by expression of the yeast *URA3* gene (13,14). The *URA3/5-FOA* system has been used to select against the expression of a functional *URA3* gene in bacteria (14), but it has not been employed in the context of an interaction trap selection system in bacteria. For certain applications, bacteria provide a superior in vivo selection system to yeast (7,15,16). The

transformation efficiency of bacteria is approximately 1000 times greater than yeast, which permits the interrogation of much larger libraries. Furthermore, fundamental differences in mechanism between prokaryotic and eukaryotic transcriptional regulation reduce the possibility of interactions between baits or preys of eukaryotic origin and components of the bacterial transcriptional system that could lead to constitutive activation or repression of the reporter gene.

A counter-selection system for bacteria would provide a mechanism for eliminating self-activating baits/preys from libraries (5,17) and would facilitate the identification of mutations that inhibit a protein-protein, protein-DNA, or protein-RNA interaction, thereby permitting the mapping of important functional residues at a recognition interface (5,12). Given the potential utility of a counter-selectable marker in bacteria, we developed a *URA3* reporter system for bacterial-based interaction trap systems (Figure 1). This counter-selectable marker eliminates functional bait-prey interactions with a low breakthrough rate. We also constructed a tandem *HIS3-URA3* reporter system that provides the flexibility of performing positive or negative selections using a single reporter construct.

## MATERIALS AND METHODS

### $\Delta$ *pyrF* Bacterial Strain Construction

The *pyrF* gene was deleted from two different *E. coli* strains, KJ1C and US0, which bear a preexisting deletion in the *hisB* gene (Reference 7 and J. Hurt, J.K. Joung, and C.O. Pabo, unpublished data). The US0 strain is isogenic to KJ1C, except that it lacks a *Tn10* insertion positioned near the *lac* operon deletion present in both strains. The *pyrF* gene was inactivated in both strains using a  $\lambda$  red recombination system (18). Primers 5'-GCCCATCA TCAAGAAGGTCTGGTCATGAC GTTAAGTCTTCATCTTCTTCC GTGTAGGCTGGAGCTGCTTCG-3' and 5'-TCATGCACTCCGCTGTAA AGAGGCGTTGATCGCTTTCAGC GTCTGCGCTGGACATATGAATAT

CCTCCTTAG-3' were used to amplify a region of the pKD4 recombination plasmid with about 50 bp of homology to *pyrF* (bolded letters in the primer sequence) at each end. The PCR product was digested with *DpnI* to remove any plasmid template, gel-purified, and transformed into electrocompetent KJ1C and US0 cells containing a  $\lambda$  red helper plasmid (pKD46). The transformants were recovered in SOB media with 10 mM L-arabinose and screened for recombination on kanamycin plates. Candidate insertions in the *pyrF* gene were confirmed by PCR using primers that amplify *pyrF* (5'-AGGGGAGAAATCGCAACTGT-3', 5'-CTTATCGCCCTGGATTCAA-3'). The kanamycin gene was removed from positive clones by FLP-mediated recombination by introducing plasmid pCP20. The scars remaining from the recombination process in the KJ1C and US0 strains were confirmed by sequencing the PCR-amplified region of the genome. The F' episome from XL1-Blue (Stratagene, La Jolla, CA, USA), which contains *lacI<sup>q</sup>* was introduced into US0 $\Delta$ *hisB* $\Delta$ *pyrF* through conjugation followed by selection on minimal medium containing 0.1% histidine, 0.1 mM uracil, 10  $\mu$ g/mL tetracycline, and 2.5 mM 5-FOA.

### Counter-Selection Medium

Yeast extract minimal medium (YM) is composed of M9 minimal medium supplemented with 10  $\mu$ M ZnSO<sub>4</sub>, 100  $\mu$ M CaCl<sub>2</sub>, 1 mM MgSO<sub>4</sub>, 10  $\mu$ g/mL thiamine, 25  $\mu$ g/mL kanamycin, 30  $\mu$ g/mL chloramphenicol, 200  $\mu$ M uracil, 0.1% histidine, 0.01% yeast extract, and a mixture of 17 amino acids (excluding histidine, methionine, and cysteine) (19). 5-FOA (2.5 and 0.2 mM) was used for the counter-selections of pH3U3 and F' episome reporter, respectively. The composition of NM media has been previously described (19). For selective plates, 1.8% agar was added to the medium mixture.

### Plasmids and Strains Used in the Bacterial Counter-Selection System


The bacterial counter-selection system is derived from a previously described bacterial two-hybrid system (7). Zif268 and ZnFp53 DNA binding domains (DBDs) (20,21) were expressed as direct fusions to the  $\alpha$  subunit of RNA polymerase ( $\alpha$ -Zif268 or  $\alpha$ -ZnFp53) using a modified pACL- $\alpha$ Gal4 vector (7), which contains a p15A origin of replication and chloramphenicol resistance marker. DNA encoding a short linker and each DNA

binding domain was substituted for Gal4 between the *NotI* and *AvrII* sites. An additional ampicillin resistance marker was introduced into the  $\alpha$ -ZnFp53 expression plasmid to distinguish it from the  $\alpha$ -Zif268 plasmid.

Two types of *URA3* reporters were developed for the counter-selection system: (i) the *URA3* F' episome reporter and (ii) the pH3U3 reporter. The *URA3* F' episome reporter is architecturally identical to the previously reported *HIS3* (7) and *lacZ* (16) F' episome reporters. The p1566-*URA3-aadA* plasmid for creating the F' episome reporter was constructed by substituting *URA3* from *S. cerevisiae* for *HIS3* between the *BamHI* and *XbaI* restriction sites in p1566-*HIS3-aadA*, which has a p15A origin of replication and kanamycin resistance marker. DNA binding sequences for Zif268 or ZnFp53 (Zif268: 5'-GCGTGGGCG-3'; ZnF53: 5'-GGGACACGT-3') (20,21) were introduced into p1566-*URA3-aadA* centered 62 bp upstream of the transcription start site of the weak *lac* promoter (7). The *URA3* reporter cassettes were integrated into the F' episome of the CSH100 strain via recombination and then introduced into the KJ1C $\Delta$ *hisB* $\Delta$ *pyrF* strains by conjugation to make the *URA3* F' episome reporter strains (7,16,22). An alternate plasmid-based reporter (pH3U3), which contains a *HIS3-URA3* cistron was constructed that allows both positive and negative selections (23). Reporters containing the Zif268 or ZnFp53 sites in pH3U3 were generated, and the selections using these vectors were performed in the US0 $\Delta$ *hisB* $\Delta$ *pyrF* strain containing the F' episome from XL1-Blue.

### URA3/5-FOA Counter-Selection Procedure

Electrocompetent cells were prepared of KJ1C $\Delta$ *hisB* $\Delta$ *pyrF* strains bearing different F' episome *URA3* reporters and of US0 $\Delta$ *hisB* $\Delta$ *pyrF* strains bearing different pH3U3 reporters. These competent cells were transformed with the  $\alpha$ -Zif268 or  $\alpha$ -ZnFp53 expression vector and recovered in 1 mL SOC for 1 h at 37°C. Following recovery, these cells were pelleted and resuspended in 1 mL YM



**Figure 1.** Overview of the bacterial-based *URA3* counter-selection system. Survival of US0 $\Delta$ *hisB* $\Delta$ *pyrF* transformants in the presence of 5-fluoro-orotic acid (5-FOA) depends on the absence of an interaction between the DNA binding domain (DBD) and a DNA sequence upstream of the *URA3* reporter in a one-hybrid configuration (top). If the DBD binds upstream of the promoter, it will recruit RNA polymerase through a direct fusion to an RNA polymerase subunit and stimulate *URA3* expression. In principle, the stronger the interaction between the DBD and a sequence upstream of the promoter, the less tolerant an individual cell will be to 5-FOA due to increased expression of *URA3* (12).

containing 25  $\mu\text{g}/\text{mL}$  kanamycin and 30  $\mu\text{g}/\text{mL}$  chloramphenicol and grown for 1 h at 37°C. The cells were pelleted and washed three times with 1 mL sterile water and then resuspended in 1 mL YM for counter-selection on YM plates containing 25  $\mu\text{g}/\text{mL}$  kanamycin, 30  $\mu\text{g}/\text{mL}$  chloramphenicol, 10  $\mu\text{M}$  isopropyl- $\beta$ -D-thiogalactopyranoside (IPTG), and either 0.2 or 2.5 mM 5-FOA. Plates were incubated at 37°C overnight.

## RESULTS AND DISCUSSION


### Creation of $\Delta\text{pyrF}$ Bacterial Strains

*pyrF*, which is the bacterial homolog of *URA3*, was inactivated in two  $\Delta\text{hisB}$  *E. coli* strains, KJ1C and US0, which were previously developed for HIS3/3-AT-based positive selection (7). The *pyrF* gene was interrupted using the  $\lambda$  red homologous recombination protocol described by Datsenko and

Wanner (18). PCR analysis of the KJ1C and US0 isolates constructed using this procedure confirmed that the majority of the *pyrF* gene was deleted from the genome (data not shown). These new  $\Delta\text{hisB}\Delta\text{pyrF}$  strains are resistant to 5-FOA and require uracil and histidine for growth on NM media (Figure 2A).


### Evaluation of the URA3/5-FOA Bacterial Counter-Selection System

The effectiveness of URA3/5-FOA counter-selection system in bacteria was evaluated using a single copy reporter integrated into the F' episome. This arrangement is homologous to the *HIS3* and *lacZ* reporters previously developed for the bacterial two-hybrid system (7,16). Cells containing the *URA3* reporter at its basal level of activity can tolerate media containing 0.2 mM 5-FOA. Two *URA3* reporters were constructed (*Zif268-URA3* and *ZnFp53-URA3*) that contain a binding site for the corresponding zinc finger proteins, *Zif268* or *ZnFp53* (20,21), upstream of the promoter (Figure 1). The *Zif268-URA3* and *ZnFp53-URA3* F' episome reporters should be selectively activated by the *Zif268* and *ZnFp53* proteins, respectively, when these proteins are expressed as fusions to the  $\alpha$ -subunit of RNA polymerase (6,24). Electrocompetent cells containing each *URA3* reporter were transformed with expression vectors for  $\alpha$ -*Zif268* or  $\alpha$ -*ZnFp53*. Following recovery, cells containing the four different combinations of reporters and DBDs were challenged to survive on minimal medium plates containing 0.2 mM 5-FOA. Cells containing matched reporters and DBDs (*ZnFp53-URA3*/ $\alpha$ -*ZnFp53* or *Zif268-URA3*/ $\alpha$ -*Zif268*) displayed a significantly reduced survival rate (about 1 in  $10^4$  cells), whereas cells containing mismatched reporters and DBDs displayed nearly complete survival (Figures 2B and 3). A mixed population of expression vectors (where the  $\alpha$ -*Zif268*/ $\alpha$ -*ZnFp53* ratio is  $10^3$  to 1) was used to test the ability of the URA3/5-FOA counter-selection to isolate a desired subpopulation through selection. This mixture was transformed into the selection strain containing the *Zif268-URA3* reporter. Approximately 85% of the



**Figure 2. Deletion of the *pyrF* gene in the US0 bacterial strain inactivates the uracil biosynthesis pathway, permitting *URA3* to be used as a counter-selectable marker.** (A) Tenfold serial dilutions of cells from one of the US0 $\Delta\text{hisB}\Delta\text{pyrF}$  strains and the parent US0 $\Delta\text{hisB}$  strain (WT) were spotted in 5- $\mu\text{L}$  drops on NM minimal plates with histidine and either containing uracil, lacking uracil, or containing uracil and 2.5 mM 5-fluoro-orotic acid (5-FOA). The  $\Delta\text{pyrF}$  strain requires uracil in the media to survive and tolerates the presence of 5-FOA. (B) Counter-selection using the *Zif268-URA3* F' episome reporter strain. Left plate, approximately  $5 \times 10^5$  cells of the *Zif268-URA3* F' episome reporter strain containing the  $\alpha$ -*Zif268* expression vector were plated on one 150  $\times$  15 mm 0.2 mM 5-FOA plate, and 80 colonies survived. Right plate, approximately  $7 \times 10^5$  cells of the *Zif268-URA3* F' episome reporter strain containing a mixture of the  $\alpha$ -*Zif268* and  $\alpha$ -*ZnFp53* expression vectors at a ratio of  $10^3$  to 1 were plated on one 150  $\times$  15 mm 0.2 mM 5-FOA plate, and about 450 colonies survived. The  $\alpha$ -*ZnFp53* expression vector contains an extra antibiotic marker ( $\text{amp}^R$ ) that allows cells containing this vector to be distinguished from cells containing the  $\alpha$ -*Zif268* expression vector. Based on the presence of this extra marker, approximately 85% of surviving colonies contained the  $\alpha$ -*ZnFp53* expression vector. The 5-FOA counter-selection using the F' episome *URA3* reporter was effective at cell densities as high as  $10^7$  cells on a 150  $\times$  15 mm round plate.





**Figure 3. Activation of the *URA3* F' episome reporter allows survival in the presence of 6-azauracil (6-AU) and eliminates cells in the presence of 5-fluoro-orotic acid (5-FOA).** Fivefold serial dilutions of *KJ1CΔhisBΔpyrF* cells containing each of four combinations of F' episome reporters and DNA binding domains (DBDs) were spotted in 5- $\mu$ L drops on plates with or without selective media in the following order: row 1, the *Zif268-URA3/α-Zif268*; row 2, the *Zif268-URA3/α-ZnFp53*; row 3, the *ZnFp53-URA3/α-Zif268*; and row 4, *ZnFp53-URA3/α-ZnFp53*. (A) Cells were plated on NM media containing either uracil or 6-AU, and colonies were observed on the 6-AU plate only in the matched reporter/DBD sets (rows 1 and 4), although some toxicity was observed even in these sets at high cell densities. (B) Cells were plated on either yeast extract tryptone medium (2 $\times$ YT) or yeast extract minimal medium (YM) containing 0.2 mM 5-FOA. In this case, nearly complete survival is observed for the mismatched sets (rows 2 and 3), whereas effective elimination of the matched sets is observed (rows 1 and 4).

cells that survived the counter-selection contained the  $\alpha$ -ZnFp53 construct (Figure 2B). The increased activity of the *URA3* F' episome reporter in the matched reporter/DBD combinations was confirmed by titrating cells on uracil-deficient media containing 6-azauracil (6-AU), a compound that is metabolized into a competitive inhibitor of the *URA3* enzyme (25,26). As expected, under these conditions, only matched reporters and DBDs (*ZnFp53-URA3/α-ZnFp53* or *Zif268-URA3/α-Zif268*) produced colonies (Figure 3).

A new bacterial counter-selection reporter vector was constructed to decrease the breakthrough rate of 5-FOA counter-selections and to allow positive (*HIS3*) or negative (*URA3*) selections to be performed with the same construct. The pH3U3 reporter plasmid contains a synthetic *HIS3-URA3* cistron expressed from the weak *lac* promoter (23). Combinations of pH3U3 reporter plasmids (containing either *Zif268* or *ZnFp53* binding sites) and expression plasmids ( $\alpha$ -*Zif268* or  $\alpha$ -*ZnFp53*) were introduced into the *US0ΔhisBΔpyrF* strain to examine the effectiveness of 5-FOA counter-selection. Following recovery, the four different combinations of transformants were titrated on 2.5 mM 5-FOA YM plates. A higher efficiency of killing was observed for the *URA3/5-FOA* counter-selection when using the pH3U3 reporter than when using the F' episome *URA3* reporter (Figure 4A). Cells containing matched reporters and DBDs (*Zif268-pH3U3/α-Zif268* or *ZnFp53-pH3U3/α-ZnFp53*) displayed a lower breakthrough rate (about 1 in  $10^5$  cells), while cells containing mismatched reporters and DBDs (*Zif268-pH3U3/α-ZnFp53* or *ZnFp53-pH3U3/α-Zif268*) were fully tolerant of these conditions. The opposite result is obtained when a *HIS3/3-AT* selection is performed; only cells that contain matched reporters and DBDs survive in the presence of 2 mM 3-AT (data not shown).

A mixed population of expression vectors (where  $\alpha$ -*Zif268/α-ZnFp53* ratio is 1 to  $10^3$ ) was used to test the ability of the *URA3/5-FOA* counter-selection to isolate a desired subpopulation through selection. This

mixture was transformed into the *US0ΔhisBΔpyrF* strain containing the ZnFp53-pH3U3 reporter, and counter-selections were performed as described above. Only approximately 1 in 10<sup>3</sup> cells survived the selection, which is consistent with the number of Zif268 clones in the population. A similar result was achieved when the ratio of the two vectors was reversed, and the mixture was transformed into the *US0ΔhisBΔpyrF* strain containing the Zif268-pH3U3 reporter. Expression plasmids from nine of the surviving colonies of each selection were prepared and tested by analytical restriction enzyme digestion to identify the transcription factor

expressed in each clone (Figure 4B). All nine clones selected using the ZnFp53-pH3U3 reporter contained the Zif268 transcription factor, and all nine clones selected using the Zif268-pH3U3 reporter contained the ZnFp53 transcription factor. As with the F' episome reporter, approximately 10<sup>7</sup> cells can be challenged on a single selection plate without increased rates of breakthrough.

In summary, we have developed a counter-selectable marker for the bacterial interaction trap system that could be used to search libraries for a protein or compound that interferes with an interaction or to isolate a macromolecule that does not mediate

a certain protein-protein, protein-DNA, or protein-RNA interaction. The *URA3* reporter can also be used to remove self-activating clones from a population (23). The *URA3/5-FOA* counter-selection system can be used in conjunction with an F' episome reporter or as a linked *HIS3-URA3* reporter vector. The *HIS3-URA3* reporter provides the advantage that it can also be used for *HIS3/3-AT*-based positive selections using the *ΔhisBΔpyrF* strains described herein. These reagents continue to extend the utility of the bacterial interaction trap system as a powerful alternative to its yeast-based counterpart.

#### ACKNOWLEDGMENTS


We thank the *E. coli* genetic stock center for the  $\lambda$  red recombination system, Jessica Hurt and Carl Pabo for plasmids and strains, and Peter Pryciak for the *URA3* gene. S.A.W. and X.M. were supported in part by the Concern Foundation and National Institutes of Health (NIH) grant no. R01GM068110, R.M.S. was supported in part by NIH grant no. R01GM068110, and A.V.G. and J.K.J. were supported in part by start-up funds from the Massachusetts General Hospital (MGH) Department of Pathology and NIH grant no. R01GM072621. This work was supported in part by NIH grant nos. R01GM068110 (to S.A.W.), R01GM069906 (to J.K.J.), and R01GM072621 (to J.K.J.).

#### COMPETING INTERESTS STATEMENT

X.M., J.K.J., and S.A.W. declare that they have a patent application pending on related technology. All other authors declare no competing interests.

#### REFERENCES

1. Wang, M.M. and R.R. Reed. 1993. Molecular cloning of the olfactory neuronal transcription factor Olf-1 by genetic selection in yeast. *Nature* 364:121-126.
2. Li, J.J. and I. Herskowitz. 1993. Isolation of ORC6, a component of the yeast origin recognition complex by a one-hybrid system. *Science* 262:1870-1874.



**Figure 4. Counter-selection using the pH3U3 reporter system.** (A) Counter-selection efficiency of 5-fluoro-orotic acid (5-FOA) using a pH3U3 reporter. Cells ( $1.3 \times 10^6$ ) containing ZnFp53 expression vector and pH3U3 reporter with ZnFp53 binding site were plated on either a yeast extract tryptone medium (2x YT) plate (no counter-selection) or a yeast extract minimal medium (YM) plate with 2.5 mM 5-FOA. A lawn of cells was produced on the 2x YT plate, while only 20 colonies were apparent on the 5-FOA plate. (B) Selection of a desired expression vector from a mixed population via *URA3/5-FOA* counter-selection. Cells containing Zif268-pH3U3 reporter were transformed with a mixture of  $\alpha$ -Zif268/ $\alpha$ -ZnFp53 expression vectors at a ratio of 10<sup>3</sup> to 1. Approximately 10<sup>5</sup> of these cells were challenged to grow on YM with 2.5 mM 5-FOA. Fewer than 100 colonies survived the counter-selection. Plasmid DNA from nine colonies was isolated and analyzed by restriction enzyme digestion (*NotI*). All nine clones contained the  $\alpha$ -ZnFp53 vector (lanes 3–12) as evidenced by the presence of the expected 4700- and 250-bp fragments. Lanes 1 and 2 are positive controls of  $\alpha$ -ZnFp53 and  $\alpha$ -Zif268 vector, respectively, cut by *NotI*. The corresponding experiment was performed with the ZnFp53-pH3U3 reporter, where the ratio of the mixture of  $\alpha$ -Zif268/ $\alpha$ -ZnFp53 expression vectors was inverted. Plasmid DNA from nine colonies was isolated and analyzed by restriction enzyme digestion (*NotI*). All nine clones contained the  $\alpha$ -Zif268 vector as evidenced by the presence of the expected 3900-bp fragment (lanes 13–22). The extra bands visible in lanes 13–22 are from the pH3U3 reporter vector. In each gel, the left unmarked lane is the 1-kb DNA ladder from New England Biolabs (Ipswich, MA, USA), and the right unmarked lane is the 100-bp DNA ladder (also from New England Biolabs).

3. **Fields, S.** 1993. The two-hybrid system to detect protein-protein interactions. *Methods* 5:116-124.
4. **SenGupta, D.J., B. Zhang, B. Kraemer, P. Pochart, S. Fields, and M. Wickens.** 1996. A three-hybrid system to detect RNA-protein interactions in vivo. *Proc. Natl. Acad. Sci. USA* 93:8496-8501.
5. **Vidal, M. and P. Legrain.** 1999. Yeast forward and reverse 'n'-hybrid systems. *Nucleic Acids Res.* 27:919-929.
6. **Dove, S.L., J.K. Joung, and A. Hochschild.** 1997. Activation of prokaryotic transcription through arbitrary protein-protein contacts. *Nature* 386:627-630.
7. **Joung, J.K., E.I. Ramm, and C.O. Pabo.** 2000. A bacterial two-hybrid selection system for studying protein-DNA and protein-protein interactions. *Proc. Natl. Acad. Sci. USA* 97:7382-7387.
8. **Althoff, E.A. and V.W. Cornish.** 2002. A bacterial small-molecule three-hybrid system. *Angew. Chem. Int. Ed. Engl.* 41:2327-2330.
9. **Karimova, G., J. Pidoux, A. Ullmann, and D. Ladant.** 1998. A bacterial two-hybrid system based on a reconstituted signal transduction pathway. *Proc. Natl. Acad. Sci. USA* 95:5752-5756.
10. **Boeke, J.D., F. LaCroute, and G.R. Fink.** 1984. A positive selection for mutants lacking orotidine-5'-phosphate decarboxylase activity in yeast: 5-fluoro-orotic acid resistance. *Mol. Gen. Genet.* 197:345-346.
11. **Boeke, J.D., J. Trueheart, G. Natsoulis, and G.R. Fink.** 1987. 5-Fluoroorotic acid as a selective agent in yeast molecular genetics. *Methods Enzymol.* 154:164-175.
12. **Vidal, M., R.K. Brachmann, A. Fattaey, E. Harlow, and J.D. Boeke.** 1996. Reverse two-hybrid and one-hybrid systems to detect dissociation of protein-protein and DNA-protein interactions. *Proc. Natl. Acad. Sci. USA* 93:10315-10320.
13. **Bach, M.L., F. Lacroute, and D. Botstein.** 1979. Evidence for transcriptional regulation of orotidine-5'-phosphate decarboxylase in yeast by hybridization of mRNA to the yeast structural gene cloned in *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* 76:386-390.
14. **Broschard, T.H., A. Lebrun-Garcia, and R.P. Fuchs.** 1998. Mutagenic specificity of the food mutagen 2-amino-3-methylimidazo[4,5-f]quinoline in *Escherichia coli* using the yeast URA3 gene as a target. *Carcinogenesis* 19:305-310.
15. **Serebriiskii, I.G., R. Fang, E. Latypova, R. Hopkins, C. Vinson, J.K. Joung, and E.A. Golemis.** 2005. A combined yeast/bacteria two-hybrid system: development and evaluation. *Mol. Cell. Proteomics* 4:819-826.
16. **Hurt, J.A., S.A. Thibodeau, A.S. Hirsh, C.O. Pabo, and J.K. Joung.** 2003. Highly specific zinc finger proteins obtained by directed domain shuffling and cell-based selection. *Proc. Natl. Acad. Sci. USA* 100:12271-12276.
17. **Walhout, A.J. and M. Vidal.** 1999. A genetic strategy to eliminate self-activator baits prior to high-throughput yeast two-hybrid screens. *Genome Res.* 9:1128-1134.
18. **Datsenko, K.A. and B.L. Wanner.** 2000. One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc. Natl. Acad. Sci. USA* 97:6640-6645.
19. **Serebriiskii, I. and J. Joung.** 2001. Yeast and bacterial two-hybrid selection systems for studying protein-protein interactions, p. 93-142. *In* E. Golemis (Ed.), *Protein-Protein Interactions: A Molecular Cloning Manual*. CSH Laboratory Press, Cold Spring Harbor, NY.
20. **Greisman, H.A. and C.O. Pabo.** 1997. A general strategy for selecting high-affinity zinc finger proteins for diverse DNA target sites. *Science* 275:657-661.
21. **Wolfe, S.A., H.A. Greisman, E.I. Ramm, and C.O. Pabo.** 1999. Analysis of zinc fingers optimized via phage display: evaluating the utility of a recognition code. *J. Mol. Biol.* 285:1917-1934.
22. **Whipple, F.W.** 1998. Genetic analysis of prokaryotic and eukaryotic DNA-binding proteins in *Escherichia coli*. *Nucleic Acids Res.* 26:3700-3706.
23. **Meng, X., M.H. Brodsky, and S.A. Wolfe.** 2005. A bacterial one-hybrid system for determining the DNA-binding specificity of transcription factors. *Nat. Biotechnol.* 23:988-994.
24. **Dove, S.L. and A. Hochschild.** 1998. Conversion of the omega subunit of *Escherichia coli* RNA polymerase into a transcriptional activator or an activation target. *Genes Dev.* 12:745-754.
25. **Levine, H.L., R.S. Brody, and F.H. Westheimer.** 1980. Inhibition of orotidine-5'-phosphate decarboxylase by 1-(5'-phospho-beta-d-ribofuranosyl)barbituric acid, 6-azauridine 5'-phosphate, and uridine 5'-phosphate. *Biochemistry* 19:4993-4999.
26. **Le Douarin, B., B. Pierrat, E. vom Baur, P. Chambon, and R. Losson.** 1995. A new version of the two-hybrid assay for detection of protein-protein interactions. *Nucleic Acids Res.* 23:876-878.

Received 22 July 2005; accepted 30 August 2005.

Address correspondence to Scot A. Wolfe, Program in Gene Function and Expression, Department of Biochemistry and Molecular Pharmacology, University of Massachusetts Medical School, 364 Plantation St., Worcester, MA 01605, USA. e-mail: scot.wolfe@umassmed.edu

To purchase reprints  
of this article, contact  
[Reprints@BioTechniques.com](mailto:Reprints@BioTechniques.com)