

6-27-2001

# T cell receptor Vbeta gene usage in Thai children with dengue virus infection

Susan J. Gagnon

*University of Massachusetts Medical School*

Anita M. Leporati

*University of Massachusetts Medical School, Anita.Leporati@umassmed.edu*

Sharone Green

*University of Massachusetts Medical School, sharone.green@umassmed.edu*

*See next page for additional authors*

Follow this and additional works at: <http://escholarship.umassmed.edu/oapubs>

 Part of the [Immunology and Infectious Disease Commons](#), and the [Pediatrics Commons](#)

---

## Repository Citation

Gagnon, Susan J.; Leporati, Anita M.; Green, Sharone; Kalayanarooj, Siripen; Vaughn, David W.; Stephens, Henry A. F.; Suntayakorn, Saroj; Kurane, Ichiro; Ennis, Francis A.; and Rothman, Alan L., "T cell receptor Vbeta gene usage in Thai children with dengue virus infection" (2001). *Open Access Articles*. 163.

<http://escholarship.umassmed.edu/oapubs/163>

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in Open Access Articles by an authorized administrator of eScholarship@UMMS. For more information, please contact [Lisa.Palmer@umassmed.edu](mailto:Lisa.Palmer@umassmed.edu).

---

# T cell receptor Vbeta gene usage in Thai children with dengue virus infection

## **Authors**

Susan J. Gagnon, Anita M. Leporati, Sharone Green, Siripen Kalayanarooj, David W. Vaughn, Henry A. F. Stephens, Saroj Suntayakorn, Ichiro Kurane, Francis A. Ennis, and Alan L. Rothman

## **Rights and Permissions**

Citation: Am J Trop Med Hyg. 2001 Jan-Feb;64(1-2):41-8.

## T CELL RECEPTOR V $\beta$ GENE USAGE IN THAI CHILDREN WITH DENGUE VIRUS INFECTION

SUSAN J. GAGNON, ANITA LEPORATI, SHARONE GREEN, SIRIPEN KALAYANAROOJ, DAVID W. VAUGHN, HENRY A. F. STEPHENS, SAROJ SUNTAYAKORN, ICHIRO KURANE, FRANCIS A. ENNIS, AND ALAN L. ROTHMAN  
*Center for Infectious Disease and Vaccine Research, University of Massachusetts Medical School, Worcester, Massachusetts; Queen Sirikit National Institute of Child Health (formerly Bangkok Children's Hospital), Bangkok, Thailand; Armed Forces Research Institute of Medical Sciences, Bangkok, Thailand; Department of Transfusion Medicine, Siriraj Hospital, Bangkok, Thailand; Kamphaeng Phet Provincial Hospital, Kamphaeng Phet, Thailand; Department of Virology I, National Institute of Infectious Diseases, Tokyo, Japan*

**Abstract.** T lymphocyte activation during dengue is thought to contribute to the pathogenesis of dengue hemorrhagic fever (DHF). We examined the T cell receptor V $\beta$  gene usage by a reverse transcriptase-polymerase chain reaction assay during infection and after recovery in 13 children with DHF and 13 children with dengue fever (DF). There was no deletion of specific V $\beta$  gene families. We detected significant expansions in usage of single V $\beta$  families in six subjects with DHF and three subjects with DF over the course of infection, but these did not show an association with clinical diagnosis, viral serotype, or HLA alleles. Differences in V $\beta$  gene usage between subjects with DHF and subjects with DF were of borderline significance. These data suggest that the differences in T cell activation in DHF and DF are quantitative rather than qualitative and that T cells are activated by conventional antigen(s) and not a viral superantigen.

### INTRODUCTION

Infection with dengue virus (DV), a mosquito-borne flavivirus, is a major health problem for many tropical and subtropical areas of the world, where as many as 100 million DV infections occur yearly.<sup>1</sup> This virus exists as four distinct serotypes;<sup>2</sup> infection with one DV serotype induces long-lived immunity against reinfection with that serotype. However, epidemiologic and laboratory studies indicate that reinfection with a heterologous DV serotype is more likely to result in the potentially life-threatening form of disease termed dengue hemorrhagic fever (DHF), characterized by increased capillary permeability.<sup>1,3–5</sup> It has been proposed that DHF reflects an immunopathologic response to DV infection.<sup>6</sup> Studies have demonstrated that levels of circulating soluble CD8 and soluble interleukin-2 receptors (sIL-2R), markers of immune activation, are higher in children with DHF than in those with dengue fever (DF).<sup>7,8</sup> Additionally, a high level of CD69 expression has been observed on circulating peripheral blood mononuclear cells (PBMCs) of children with DHF.<sup>9</sup>

The biological basis for the increased T cell activation in DHF has not been determined. Expansion of T cell subsets bearing T cell receptors (TCRs) using particular V $\beta$  gene families has been associated with severe disease in other infectious or autoimmune disorders. Specific examples include toxic shock syndrome, leprosy, infection with human immunodeficiency virus (HIV), Crohn's disease, and reactive arthritis.<sup>10–15</sup> *In vitro* analysis of TCR usage by a panel of DV-specific CD4<sup>+</sup> cytotoxic T lymphocyte clones generated from one individual after a primary dengue-4 virus infection demonstrated that eight of 19 clones expressed V $\beta$ 17.<sup>16</sup> A similar expansion of V $\beta$  17-expressing cells was observed following bulk culture stimulation of this donor's PBMCs with D4 antigen but not following stimulation with an anti-CD3 antibody. These data provided preliminary evidence that DV-infection might result in expansion of T cells bearing a particular V $\beta$  chain.

We therefore sought to determine whether an expansion of T lymphocytes expressing a particular V $\beta$  gene occurs *in*

*vivo* upon DV infection and whether preferential V $\beta$  region usage correlates with disease severity. We analyzed the V $\beta$  gene usage in PBMCs obtained during and after the acute infection from 13 subjects with DHF and 13 with DF using a semi-quantitative polymerase chain reaction (PCR) method.<sup>16</sup> We confirmed that this method yields highly reproducible results. We detected expansions of T cells expressing particular V $\beta$  genes during acute DV infection in some subjects. However, we did not find consistent expansions of specific V $\beta$  gene families among the group of subjects as a whole or significant differences in V $\beta$  usage between patients with DHF and those with DF.

### MATERIALS AND METHODS

**Clinical study design and sample collection.** Thai children between the ages of six months and 14 years were enrolled in the Dengue Hemorrhagic Fever Project, as previously reported.<sup>17</sup> Written informed consent was obtained from the subjects' parents or guardians. The study protocol was approved by Institutional Review Boards established by the Ministry of Public Health, Thailand, the Surgeon General's Office of the Department of the Army, and the University of Massachusetts. We selected at random 13 children with DHF and 13 children with DF from among the study population enrolled between April 1994 and November 1995. Dengue virus infections were confirmed by serologic testing and virus isolation as described.<sup>18</sup> Clinical diagnoses of DHF and DF were assigned according to World Health Organization criteria.<sup>17,19</sup>

Blood samples were obtained daily during illness and at a follow-up visit six months after the acute DV infection.<sup>17</sup> The PBMCs were isolated by density gradient centrifugation using Histopaque (Sigma, St. Louis, MO). They were washed and counted, and  $2 \times 10^6$  cells were then placed into an Eppendorf tube (Brinkman Instruments, Westbury, NY) and washed twice with phosphate-buffered saline. Finally, the cells were pelleted and 1.0 ml of solution D (4 M guanidinium isothiocyanate, 25 mM sodium citrate pH 7.0, 2%

[w/v] sarcosyl, and 0.1 M 2-mercaptoethanol) was added. Samples were then stored at  $-70^{\circ}\text{C}$ , shipped to the University of Massachusetts Medical School on dry ice, and kept at  $-70^{\circ}\text{C}$  until analysis. We selected for this study blood samples obtained on study day 2 (one day after enrollment), fever day +1 (one day after defervescence), and study day 180 (follow-up visit). All samples were analyzed under code.

**Isolation of RNA and first-strand cDNA synthesis.** Total cellular RNA was isolated by the acid guanidinium-phenol-chloroform method.<sup>20</sup> During this procedure, precipitation with isopropanol was performed with glycogen as a carrier.<sup>21</sup> The RNA was treated with DNaseI for 2–3 hr to degrade the DNA, with RNAGuard (Pharmacia Biotech, Inc., Piscataway, NJ) to prevent RNA degradation. First-strand cDNA was synthesized from this RNA using random hexamer primers (Pharmacia Biotech, Inc.) and Moloney murine leukemia virus reverse transcriptase (Gibco-BRL, Gaithersburg, MD) for 4–16 hr, and the reaction was terminated by heating samples at  $100^{\circ}\text{C}$  for 10 min.

**Analysis of TCR V $\beta$  gene usage by PCR.** Analysis of TCR V $\beta$  gene usage was performed as previously described.<sup>16</sup> Twenty-six TCR  $\beta$ -chain variable region-specific 5' oligonucleotide primers and one TCR  $\beta$ -chain constant region-specific 3' oligonucleotide primer were synthesized at the DNA Synthesis Facility at the University of Massachusetts Medical School on the basis of previous reports.<sup>16</sup>  $\beta$ -actin-specific oligonucleotide 5' and 3' primers were also prepared. The 3' primers were end-labeled with  $^{32}\text{P}$  using T4 polynucleotide kinase (Promega Corp., Madison, WI) and purified using Chroma Spin-10 columns (Clontech Laboratories, Inc., Palo Alto, CA). A volume of labeled 3' primer equivalent to  $10^5$  counts/min (cpm) was used in each reaction. We used 26 cycles of a PCR at  $95^{\circ}\text{C}$  for 1 min,  $55^{\circ}\text{C}$  for 1 min, and  $72^{\circ}\text{C}$  for 1 min. In the first cycle, a longer denaturing step (1.5 min) was used.

Following the PCR, total reaction mixtures were subjected to electrophoresis on a 5% polyacrylamide gel. The radioactivity of each amplified fragment was examined with a Betascope (Betagen, Mountain View, CA) for 100 min. The percentage of each TCR V $\beta$  subfamily expression was calculated by the following formula: expression of TCR V $\beta_n$  =  $100 \times (\text{relative expression of TCR V}\beta_n / \text{sum of the relative expression of all TCR V}\beta \text{ subfamilies})$ , where relative expression of TCR V $\beta_n$  = cpm of TCR V $\beta_n$  product/cpm of  $\beta$ -actin product amplified in the same reaction tube.

If no  $\beta$ -actin product was detected in one or more of the PCRs, the relative expression of that TCR V $\beta$  was determined from a partial repeat analysis of the sample as follows. Reactions that failed to yield a  $\beta$ -actin product were retested with at least six reactions that yielded both  $\beta$ -actin and TCR V $\beta$  products, including some with high and low relative expression. Missing values were then interpolated from the linear regression equation. The median correlation coefficient ( $r$ ) of repeated analyses was 0.97, with a range of 0.85–1.00.

**HLA typing.** HLA Class I serotyping was performed on peripheral blood lymphocytes within 24 hr of collection, using a standard two-stage microlymphocytotoxicity test for HLA-A, B, and C, as previously described.<sup>22</sup> HLA class II molecular typing of the DRB1, DQA1, DQB1, and DPB1 gene loci was performed on genomic DNA, using the 11th

and 12th International Histocompatibility Workshop primers, sequence-specific oligonucleotide probes and protocols as previously described.<sup>22,23</sup>

**Statistical analysis.** Pearson correlation was used to compare the expression of all TCR V $\beta$  gene families in serial analyses of the same PBMC samples. For comparison of TCR V $\beta$  expression in different PBMC samples, percentage values were first transformed using the logit function,  $f(x) = \log(x/(1-x))$ , using 2% as a minimum value for expression of each TCR V $\beta$  gene family (to avoid skewing of results because of small absolute changes in expression of low abundance TCR V $\beta$  transcripts). Changes in expression of specific TCR V $\beta$  gene families between two time points for an individual subject were considered significant if the differences in transformed values fell outside the mean  $\pm 3$  SD of differences for all TCR V $\beta$  gene families between those two time points. We compared the expression of each TCR V $\beta$  gene family between different time points for the study population as a whole using a paired  $t$ -test. We compared the expression of each TCR V $\beta$  gene family at each time point and changes in expression of each TCR V $\beta$  gene family between time points in subjects with DHF with the corresponding values in subjects with DF using a  $t$ -test. We set  $P < 0.01$  as the criterion for statistical significance, to adjust in part for the number of comparisons.

## RESULTS

**Characteristics of the study population.** Table 1 shows the clinical, serologic, and virologic information on the study subjects. Five of 13 DF patients had primary DV infections, while all of the DHF patients were experiencing secondary infections. All four serotypes of DV were represented in both the DF and DHF groups. Among the subjects studied with DHF, there were six with DHF grade 1, six with DHF grade 2, and one with DHF grade 3. Data on HLA class I alleles were available for 23 subjects and data on HLA class II alleles were available for 24 subjects (Table 1). The mean age of the subjects studied was 8.5 years (95% confidence interval [CI] = 7.3–9.7), and was not significantly different between those children with DHF (mean = 7.9 years) and those with DF (mean = 9.1 years).

**Reproducibility of the PCR assay.** We first analyzed the reproducibility of the PCR assay for measurement of TCR V $\beta$  gene usage by repeated analysis of cDNA prepared from seven blood samples. The mean correlation coefficient for the comparison of the results from repeated PCR amplification of the same specimens was 0.97 (95% CI = 0.96–0.99).

**Changes in TCR V $\beta$  gene usage associated with acute dengue virus infection.** To determine whether acute dengue virus infection causes significant shifts in the TCR V $\beta$  gene usage in PBMCs, we examined the TCR V $\beta$  gene usage of PBMCs obtained from each subject on study day 2, fever day +1 (one day after defervescence), and study day 180. For one patient in each group, only the study day 2 specimen was available for analysis. The mean interval between study day 2 and fever day +1 for the study population as a whole was 2.2 days (95% CI = 1.9–2.5), and was not significantly different between those children with DHF (mean = 2.2 days) and those with DF (mean = 2.3 days).

TABLE 1  
Characteristics of the study population

Subject	Diagnosis	Serologic response	Virus sero-type	HLA class I antigens	HLA class II alleles			
					DRB1*	DQA1*	DQB1*	DPB1*
C95-031	DF	Primary	1	A11.1, A33, B58, B60, C3, C7.2	0301, 0901	03, 0501	0201, 03032	0501, 1301
C95-039	DF	Primary	1	A1, A11.1, B56, B57, C6, C7.2	1502, 0701	0101, 0201	0501, 03032	1401, 0901
C94-067	DF	Primary	2	A1, A2, B39, B51, C7, C1	1501, 1502	0101, 0102	0501, 0601	0501, 1301
C95-026	DF	Primary	3	—	—	—	—	—
C95-029	DF	Primary	3	A24, A34, B7, B75, C1, C6	1502, 1202	0102, 0102	0502, 0601	0901, 2101
C94-076	DF	Secondary	1	A24, B18, B55, C3, C7	1202, 0405	03, 0601	0301, 0401	1301, 3101
C94-123	DF	Secondary	2	A11.1, A24, B70, B75, C1	1202, 1202	0601, 0601	0301, 0301	0201, 0301
K94-014	DF	Secondary	2	A2, A24, B46, C1	1502, 1202	0101, 0102	0501, 0502	0501, 1301
K94-031	DF	Secondary	3	A2, A11.1, B18, B46, C1, C7	1502, 0901	0101, 03	0501, 03032	0501, 1301
C94-062	DF	Secondary	4	A2, A11.1, B7, B51, C7, C14	1101, 1202	0501, 0601	0301, 0301	1301, 1301
C94-082	DF	Secondary	4	A2, A9.3, B18, B62, C3, C7	1101, 1202	0501, 0601	0301, 0301	0301, 1401
K94-001	DF	Secondary	4	A2, B46, B77, C1	0701, 0901	0201, 03	0201, 03032	0201, 0501
K94-012	DF	Secondary	4	A2, A11.1, B46, B75, C1	1502, 1202	0101, 0102	0501, 0502	0501, 1301
C94-045	DHF 1	Secondary	1	—	1501, 1401	0101, 0102	0502, 0601	0201, 1301
K95-006	DHF 1	Secondary	2	A11.1, A33, B56, B58, C3	1501, 0301	0501, 0102	0601, 0201	0401, 2801
K95-004	DHF 1	Secondary	3	A2, A11.2, B62, B46	1401, 1501	—	0501, 0502	—
K95-005	DHF 1	Secondary	3	A2, A11.1, B35, B38, C4, C7	1501, 1502	0102, 0102	0502, 0502	1401, 0402
C94-094	DHF 1	Secondary	4	A2, A11.1, B7, B13, C3, C7	1501, 1101	0102, 0501	0502, 0301	0201, 0202
K95-001	DHF 1	Secondary	4	A2, A33, B51, B58, C3	1202, 0301	0501, 0601	0201, 0301	0401, 0501
C94-095	DHF 2	Secondary	2	A2, A24, B60, B62, C1	1101, 1202	0501, 0601	0301, 0301	0501, 0501
C94-107	DHF 2	Secondary	2	A2, A24, B46, B75, C1	1401, 1501	0101, 0102	0502, 0601	0202, 2801
C94-070	DHF 2	Secondary	3	A11.1, A24, B38, B60, C7	1502, 0901	0101, 03	0501, 03032	0501, 2801
C94-136	DHF 2	Secondary	3	—	—	—	—	—
C94-050	DHF 2	Secondary	4	A11.1, A33, B51, B61, C1	1001, 1001	0101, 0101	0501, 0501	0401, 2801
C94-073	DHF 2	Secondary	4	A2, A33, B51, B75, C1	1501, 0901	0102, 03	0601, 03032	0402, 0402
C94-134	DHF 3	Secondary	2	A11.1, A28, B27, B57, C3, C6	1202, 0701	0201, 0601	0301, 03032	0401, 1301

DF = dengue fever; DHF = dengue hemorrhagic fever (number indicates severity grade).  
\* Allele was determined using molecular typing.

We did not note deletion of cells using any particular TCR V $\beta$  gene family following acute dengue virus infection in the study population. However, we detected TCR transcripts using V $\beta$ 20 in only one of the 26 subjects studied, including those with primary dengue virus infection. We were able to amplify TCR V $\beta$ 20 transcripts from PBMCs obtained from healthy donors from the United States and from six of the seven control Thai subjects using the same experimental procedures.

The results of our statistical analysis of the changes in TCR V $\beta$  gene usage for the study population as a whole are summarized in Table 2. All analyses for which a paired *t*-test showed  $P < 0.05$  are shown; however, using our selected cutoff of  $P < 0.01$ , there were few TCR V $\beta$  gene families that showed significant changes in usage. From study day 2 to fever day +1 there was a statistically significant decrease in usage of V $\beta$ 5.1. From fever day +1 to study day 180, there were statistically significant increases in usage of V $\beta$ 2, V $\beta$ 5.1, and V $\beta$ 6 and statistically significant decreases in usage of V $\beta$ 1 and V $\beta$ 13.1. From study day 2 to study day 180, there was a statistically significant decrease in usage of V $\beta$ 18.

Statistically significant changes in usage of one or more TCR V $\beta$  gene families during the acute infection were apparent in 12 subjects. The most common such finding was the expansion in use of a particular TCR V $\beta$  gene between study day 2 and fever day +1. We found expansion in use of a single TCR V $\beta$  gene family in six of 12 subjects with DHF and three of 12 subjects with DF ( $P$  not significant). These expansions generally did not persist in the PBMCs obtained six months after the acute infection. For example, Figure 1A shows one subject (C94-050) with DHF and sec-

ondary dengue-4 virus infection who demonstrated increased use of V $\beta$ 21 on fever day +1. Figure 1B shows one subject (C94-076) with DF and secondary dengue-1 virus infection who demonstrated increased use of V $\beta$ 24 on fever day +1. The TCR V $\beta$  gene family showing the greatest increase in usage differed among the study subjects with such expansions, and we did not identify any consistent associations with the serotype of virus causing infection or the HLA alleles of the subject.

#### Relationship of TCR V $\beta$ gene usage to disease severity.

We next compared the TCR V $\beta$  gene usage in the 13 subjects with DHF and the 13 subjects with DF (Figure 2). The TCR V $\beta$  gene usage at each of the three time points showed few differences between subjects with DHF and subjects with DF. V $\beta$ 2 gene usage was significantly higher in subjects with DF on fever day +1 (3.8% versus 2.5%;  $P = 0.006$ ). There was a trend toward higher V $\beta$ 24 gene usage in subjects with DHF on study day 2 (3.3% versus 2.6%;  $P = 0.016$ ), but this difference did not meet our criterion for significance. The results of our statistical analysis of the changes in TCR V $\beta$  gene usage in subjects with DHF and those with DF are summarized in Table 2. The change in V $\beta$ 1 gene usage between study day 2 and study day 180 was significantly different between subjects with DHF and subjects with DF (-1.3% versus +0.2%;  $P = 0.010$ ). None of the other differences between the groups met our criterion for statistical significance.

#### DISCUSSION

We analyzed the TCR V $\beta$  gene usage of PBMCs obtained at three time points during and after acute dengue virus in-

TABLE 2  
Summary of probability values for comparisons of interval changes in T cell receptor (TCR) gene usage in the study population<sup>a</sup>

Vβ	All subjects <sup>b</sup>			DHF versus DF <sup>c</sup>		
	Study day 2 to fever day +1	Fever day +1 to study day 180	Study day 2 to study day 10	Study day 2 to fever day +1	Fever day +1 to study day 180	Study day 2 to study day 10
1	0.022	(↓) 0.001 <sup>d</sup>	—	—	—	0.010
2	0.016	(↑) <0.001	—	—	—	—
3	—	—	—	—	—	—
4	—	0.038	—	—	—	—
5.1	(↓) 0.005	(↑) 0.002	—	—	—	—
5.2	—	—	—	—	0.021	—
6	—	(↑) 0.001	—	0.028	—	0.029
7	—	—	—	0.043	—	—
8	—	—	—	—	—	—
9	—	—	—	—	—	—
10	—	—	—	—	—	—
11	—	—	—	—	0.023	—
12	—	—	0.026	—	—	—
13.1	—	(↓) <0.001	0.036	—	—	—
13.2	—	—	—	—	—	0.037
14	—	—	—	—	—	—
15	—	—	—	0.036	—	—
16	—	—	—	—	—	—
17	—	—	—	—	—	—
18	—	—	(↓) 0.004	—	—	—
19	—	—	—	—	—	—
20	—	—	—	—	—	—
21	0.043	—	—	—	—	—
22	—	—	—	0.019	—	—
23	—	—	—	—	—	—
24	—	—	—	0.038	—	0.046

<sup>a</sup> All *P* values are 2-tailed. Dashes represent comparisons where *P* values are > 0.05. The direction of change in TCR gene usage is indicated only for comparisons showing *P* < 0.01.

<sup>b</sup> Logit-transformed values for TCR gene usage at the indicated time points were compared for all subjects combined by paired *t*-test.

<sup>c</sup> Changes in logit-transformed values for TCR gene usage were compared between subjects with dengue hemorrhagic fever (DHF) and those with dengue fever (DF) by *t*-test.

<sup>d</sup> (↑) or (↓), respectively, indicate an increase or decrease in Vβ expression.

fection. The earliest time point studied (study day 2) corresponded to a mean fever day value of  $-1.2$ , or slightly more than one day before defervescence. This was the closest approximation available to the viremic phase of infection,<sup>18</sup> and 24 of the 26 subjects studied had positive plasma virus culture at that time. The middle time point (fever day +1, one day after defervescence) corresponds closest to the period of plasma leakage in DHF.<sup>17</sup> The latest time point (study day 180) is the closest approximation to a baseline for each subject since we did not have access to PBMCs from before dengue virus infection in these subjects. Although the interval between the first two specimens was relatively short, we have evidence that significant immunologic changes occur during this period. In a separate group of study subjects, the expression of CD69 on CD8+ T cells of children with acute dengue virus infection was higher on study day 2 than on fever day +1, particularly in children with DHF.<sup>9</sup> On the other hand, the number of atypical lymphocytes in the peripheral blood of children with acute dengue virus infection was markedly higher on fever day +1.<sup>9,17</sup> Plasma levels of sCD8 and sIL-2R, markers of T cell activation, were elevated on study day 2 but increased further through fever day +1.<sup>7</sup>

We used a semi-quantitative PCR assay to measure the usage of 26 TCR Vβ gene families in the PBMC samples. Although some of the data were obtained from repeat analysis of the same samples and interpolation of missing data, we believe that our finding of a high interassay correlation justifies this approach.

Our analysis of the data tested several specific hypotheses.

First, we looked for increases in the usage of some TCR Vβ gene families in individual subjects during the acute infection, using the study day 180 specimen for comparison, as an indication of expansion of particular T cell populations. As shown in Figure 1, expansions in certain TCR Vβ subsets during the course of DV infection were observed in some subjects. This finding is in agreement with our *in vitro* data demonstrating preferential expansion of Vβ17-bearing T cells in the PBMCs of a dengue-immune donor upon stimulation with non-infectious DV antigen.<sup>16</sup> The TCR Vβ expansions were most often noted one day following defervescence (fever day +1), which supports the hypothesis that some of the atypical lymphocytes in the peripheral blood are proliferating DV-specific T cells.

The specific TCR Vβ gene families showing expansion during acute DV infection differed among the study subjects. We did not identify any consistent association between the TCR Vβ genes showing expansion and the clinical, virologic, or genetic profile of the subjects. We found statistically significant changes in expression of TCR Vβ1, Vβ2, Vβ5.1, Vβ6, Vβ13.1, and Vβ18 in the study population as a whole. Although we used a relatively conservative criterion for statistical significance of *P* < 0.01, these findings may still reflect type I error due to the large number of statistical comparisons performed. Therefore, we consider these to be preliminary observations that require confirmation in other study populations.

We did not detect deletions of any specific TCR Vβ gene families related to acute DV infection, regardless of the clinical severity of disease. This finding does not support the




FIGURE 1. T cell receptor (TCR) V $\beta$  gene usage during and after acute dengue virus infection in individual subjects. Fever day +1 is one day after defervescence. **A**, subject C94-050 (dengue hemorrhagic fever [DHF]) had a significant expansion of V $\beta$ 21 gene usage on fever day +1. **B**, subject C94-076 (dengue fever [DF]) had a significant expansion of V $\beta$ 24 gene usage on fever day +1.




FIGURE 2. T cell receptor (TCR) V $\beta$  gene usage during and after acute dengue virus infection in subjects with dengue hemorrhagic fever (DHF) and dengue fever (DF). Polymerase chain reaction of TCR V $\beta$  genes was performed on cDNA from 13 subjects with DHF and 13 subjects with DF. Values plotted are the mean  $\pm$  SD percent usage of 26 TCR V $\beta$  gene families at the three time points examined. Fever day +1 is one day after defervescence.

suggestion that a DV-encoded superantigen might be responsible for the increased T cell activation observed during DV infection, particularly in patients with DHF.<sup>7,8</sup> Superantigens can induce marked activation followed by deletion of T cells expressing a particular TCR V $\beta$  gene.<sup>24-26</sup> Our failure to detect such an effect in this study population is subject to several limitations. Only one of the subjects studied had grade 3 DHF, and none of the cases of DHF were associated with a primary DV infection; massive activation of T cells not specific for classical DV antigens caused by interaction with a superantigen might be more likely to play a role in those clinical situations. Also, we detected TCR V $\beta$ 20 transcripts in only one of the 26 subjects. This may reflect genetic differences between the Thai and United States populations, but we cannot exclude the possibility that T cells

using the V $\beta$ 20 gene were deleted earlier in infection or by a prior DV infection. Analysis of TCR V $\beta$  gene usage in the PBMCs of DV-naive Thai children would distinguish between these possibilities.

Lastly, we compared the TCR V $\beta$  gene usage in subjects with severe illness (DHF) with that in subjects with milder illness (DF). Although we found a significant difference in the usage of V $\beta$ 22 during the acute infection, the magnitude of this difference was small. Similarly, the statistically significant difference in the change in TCR V $\beta$ 1 gene usage from study day 2 to study day 180 between the two groups of subjects was small in magnitude. These differences may well reflect type I error. In general, our findings suggest that there are not qualitative differences in the T cell repertoire responding to acute DV infection between subjects with and



without plasma leakage, and that the increased T cell activation in DHF reflects quantitative differences in the T cell response.

Other groups have similarly failed to find differences between the TCR V $\beta$  repertoire in the PBMCs of HIV-infected subjects and that of control subjects. However, skewing of the TCR V $\beta$  repertoire has been observed in cells obtained from either the lungs<sup>14</sup> or lymph nodes<sup>27</sup> of HIV-infected patients. Others have also noted that alterations of TCR V $\beta$  gene usage occur much less frequently in PBMCs compared to the organs or tissues in which disease pathology is directly observed.<sup>28,29</sup> Dengue virus is believed to infect primarily monocytes and macrophages *in vivo*,<sup>30–32</sup> and a specific organ or tissue that preferentially supports DV replication has not been identified. However it is possible that during acute DV infection, notable biases in V $\beta$  expression may occur in tissues not examined in this study, such as lymph nodes or liver.

The pathogenesis of DHF is still poorly understood, but data suggest that increased levels of T cell activation play an important role. The cause of this heightened T cell activation is an active topic of investigation. To our knowledge, this study provides the first detailed analysis of TCR V $\beta$  gene usage in the PBMCs of patients with acute DV infections. Our results provide additional insight into the nature and timing of T cell responses that should be helpful in directing future studies into the pathophysiologic events in acute DV infection.

**Acknowledgments:** We thank Dr. Robert Lew for advice on statistical analyses; the nurses and pediatricians at the Queen Sirikit National Institute of Child Health and the Kamphaeng Phet Provincial Hospital, Thailand, for volunteer enrollment and patient management; the research nurses, technicians, technologists, and administrative personnel of the Department of Virology, Armed Forces Research Institute for Medical Sciences, Thailand, for specimen processing, data management, and serologic and virologic analyses; and Professor Dasnayane Chandanyingyong and the technicians at the Department of Transfusion Medicine, Siriraj Hospital, Thailand, for performing the HLA typing. We also thank the patients and their families for their participation in this study.

**Financial support:** This work was supported by a Program Project grant from the National Institutes of Health (NIH-P01AI34533) and support from the U.S. Army Medical Research and Materiel Command.

**Disclaimer:** The opinions contained herein are those of the authors and should not be construed as representing the official policies of the NIH or the Department of Defense.

**Authors' addresses:** Susan J. Gagnon, National Institute of Neurological Disorders and Stroke, National Institutes of Health, Building 10, Room 5N-206, Bethesda, MD 20892. Anita Leporati, Sharone Green, Francis A. Ennis, and Alan L. Rothman, Center for Infectious Disease and Vaccine Research, University of Massachusetts Medical School, Room S5-326, 55 Lake Avenue North, Worcester, MA 01655. Siripen Kalayanaroj, Queen Sirikit National Institute of Child Health, Bangkok, Thailand. David W. Vaughn, Walter Reed Army Institute of Research, Forest Glen Annex, Room 3A12A, Building 503, Robert Grant Road, Silver Spring, MD 20910. Henry A. F. Stephens, Tissue Typing Department, Guy's Hospital, 3rd Floor New Guy's House, St. Thomas Street, London SE1 9RT, United Kingdom. Saroj Suntayakorn, Kamphaeng Phet Provincial Hospital, Kamphaeng Phet, Thailand. Ichiro Kurane, National Institute of Infectious Diseases, 1-23-1 Toyama, Shinjuku-ku, Tokyo 162-8640, Japan.

**Reprint requests:** Alan L. Rothman, Center for Infectious Disease

and Vaccine Research, University of Massachusetts Medical School, Room S5-326, 55 Lake Avenue North, Worcester, MA 01655.

## REFERENCES

- Halstead S, 1988. Pathogenesis of dengue: challenges to molecular biology. *Science* 239: 476–481.
- Henchal EA, Putnak JR, 1990. The dengue viruses. *Clin Microbiol Rev* 3: 376–396.
- Burke DS, Nisalak A, Johnson DE, Scott RM, 1988. A prospective study of dengue infections in Bangkok. *Am J Trop Med Hyg* 38: 172–180.
- Halstead SB, 1980. Immunological parameters of Togavirus disease syndromes. Schlesinger RW, ed. *The Togaviruses: Biology, Structure, Replication*. New York: Academic Press, 107–173.
- Thein S, Aung MM, Shwe TN, Aye M, Zaw A, Aye K, Aye KM, Aaskov J, 1997. Risk factors in dengue shock syndrome. *Am J Trop Med Hyg* 56: 566–572.
- Rothman AL, Ennis FA, 1999. Immunopathogenesis of dengue hemorrhagic fever. *Virology* 257: 1–6.
- Green S, Vaughn DW, Kalayanaroj S, Nimmannitya S, Suntayakorn S, Nisalak A, Lew R, Innis BL, Kurane I, Rothman AL, Ennis FA, 1999. Early immune activation in acute dengue is related to development of plasma leakage and disease severity. *J Infect Dis* 179: 755–762.
- Kurane I, Innis BL, Nimmannitya S, Nisalak A, Meager A, Janus J, Ennis FA, 1991. Activation of T lymphocytes in dengue virus infections. High levels of soluble interleukin 2 receptor, soluble CD4, soluble CD8, interleukin 2, and interferon-gamma in sera of children with dengue. *J Clin Invest* 88: 1473–1480.
- Green S, Pichyangkul S, Vaughn DW, Kalayanaroj S, Nimmannitya S, Nisalak A, Kurane I, Rothman AL, Ennis FA, 1999. Early CD69 expression on peripheral blood lymphocytes from children with dengue hemorrhagic fever. *J Infect Dis* 150: 1429–1435.
- Choi Y, Lafferty JA, Clements JR, Todd JK, Gelfand EW, Kappler J, Marrack P, Kotzin BL, 1990. Selective expansion of T cells expressing V beta 2 in toxic shock syndrome. *J Exp Med* 172: 981–984.
- Schlievert PM, 1993. Role of superantigens in human disease. *J Infect Dis* 167: 997–1002.
- Allen RL, Gillespie GMA, Hall F, Edmonds S, Hall MA, Wordsworth BP, McMichael AJ, Bowness P, 1997. Multiple T cell expansions are found in the blood and synovial fluid of patients with reactive arthritis. *J Rheumatol* 24: 1750–1757.
- Ogawa H, Ito H, Takeda A, Kanazawa S, Yamamoto M, Nakamura H, Kimura Y, Yoshizaki K, Kishimoto T, 1997. Universal skew of T cell receptor (TCR) V $\beta$  usage for Crohn's disease. *Biochem Biophys Res Commun* 240: 545–551.
- Trentin L, Zambello R, Facco M, Sancetta R, Cerutti A, Milani A, Tassinari C, Crivellaro C, Cipriani A, Agostini C, Semenzato G, 1996. Skewing of the T cell receptor repertoire in the lung of patients with HIV-1 infection. *AIDS* 10: 729–737.
- Wang X, Golkar L, Uyemura K, Ohmen JD, Villahermosa LG, Fajardo TT, Cellona RV, Walsh GP, Modlin RL, 1993. T cells bearing V beta 6 T cell receptors in the cell-mediated immune response to *Mycobacterium leprae*. *J Immunol* 151: 7105–7116.
- Okamoto Y, Gagnon SJ, Kurane I, Leporati AM, Ennis FA, 1994. Preferential usage of T-cell receptor V beta 17 by dengue virus-specific human T lymphocytes in a donor with immunity to dengue virus type 4. *J Virol* 68: 7614–7619.
- Kalayanaroj S, Vaughn DW, Nimmannitya S, Green S, Suntayakorn S, Kunentrasai N, Viramitrachai W, Ratanachu-ek S, Kiatpolpoj S, Innis BL, Rothman AL, Nisalak A, Ennis FA, 1997. Early clinical and laboratory indicators of acute dengue illness. *J Infect Dis* 176: 313–321.
- Vaughn DW, Green S, Kalayanaroj S, Innis BL, Nimmannitya S, Suntayakorn S, Rothman AL, Ennis FA, Nisalak A, 1997. Dengue in the early febrile phase: viremia and antibody responses. *J Infect Dis* 176: 322–330.

19. Anonymous, 1986. *Dengue Haemorrhagic Fever: Diagnosis, Treatment and Control*. Geneva: World Health Organization.
20. Chomczynski P, Sacchi N, 1987. Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem* 162: 156–159.
21. Helms C, Graham MY, Dutchik JE, Olson MV, 1985. Laboratory methods. A new method for purifying lambda DNA from phage lysates. *DNA* 4: 39–49.
22. Chandanayingyong D, Stephens HAF, Klaythong R, Sirkong M, Udee S, Longta P, Chantangpol R, Bejrachandra S, Runruang E, 1997. HLA-A, -B, -DRB1, -DQA1 and DQB1 polymorphism in Thais. *Hum Immunol* 53: 174–182.
23. Stephens HAF, Brown AE, Chandanayingyong D, Webster HK, Sirkong M, Longta P, Vangseratthana R, Gordon DM, Lekmak S, Rungruang E, 1995. The presence of the HLA class II allele DPB1\*0501 in ethnic Thais correlates with an enhanced vaccine-induced antibody response to a malaria sporozoite vaccine. *Eur J Immunol* 25: 3142–3147.
24. Ferrero I, Anjuere F, Azcoitia I, Renno T, MacDonald HR, Ardavin C, 1998. Viral superantigen-induced negative selection of TCR transgenic CD4+ CD8+ thymocytes depends on activation, but not proliferation. *Blood* 91: 4248–4254.
25. Galelli A, Delcourt M, Wagner MC, Peumans W, Truffa-Bachi P, 1995. Selective expansion followed by profound deletion of mature V beta 8.3+ T cells in vivo after exposure to the superantigenic lectin *Urtica dioica* agglutinin. *J Immunol* 154: 2600–2611.
26. Renno T, Hahne M, MacDonald HR, 1995. Proliferation is a prerequisite for bacterial superantigen-induced T cell apoptosis in vivo. *J Exp Med* 181: 2283–2287.
27. Mion M, Indraccolo S, Feroli F, Minuzzo S, Masiero S, Zamarchi R, Barelli A, Borri A, Chieco-Bianchi L, Amadori A, 1997. TCR expression and clonality analysis in peripheral blood and lymph nodes of HIV-infected patients. *Hum Immunol* 57: 93–103.
28. Alam A, Lule J, Coppin H, Lambert N, Mazieres B, DePreval C, Cantagrel A, 1995. T-cell receptor variable region of the b-chain use in peripheral blood and multiple synovial membranes during rheumatoid arthritis. *Hum Immunol* 42: 331–339.
29. Duncan SR, Elias DJ, Roglic M, Pekny KW, Theofilopoulos AN, 1997. T-cell receptor biases and clonal proliferations in blood and pleural effusions of patients with lung cancer. *Hum Immunol* 53: 39–48.
30. Boonpucknavig S, Boonpucknavig V, Bhamarapavati N, Nimmannitya S, 1979. Immunofluorescence study of skin rash in patients with dengue hemorrhagic fever. *Arch Pathol Lab Med* 103: 463–466.
31. Halstead SB, 1989. Antibody, macrophages, dengue virus infection, shock, and hemorrhage: a pathogenetic cascade. *Rev Infect Dis* 11: S830–S839.
32. Theofilopoulos AN, Brandt WE, Russell PK, Dixon FT, 1976. Replication of dengue-2 virus in cultured human lymphoblastoid cells and subpopulations of human peripheral leukocytes. *J Immunol* 117: 953–961.