

# Stabilization and Localization of Xist RNA are Controlled by Separate Mechanisms and are Not Sufficient for X Inactivation

Christine Moulton Clemson,\* Jennifer C. Chow,‡ Carolyn J. Brown,‡ and Jeanne Bentley Lawrence\*

\*Department of Cell Biology, University of Massachusetts Medical Center, Worcester, Massachusetts 01655; and ‡Department of Medical Genetics, University of British Columbia, Vancouver, British Columbia, Canada V6T1Z3

**Abstract.** These studies address whether XIST RNA is properly localized to the X chromosome in somatic cells where human *XIST* expression is reactivated, but fails to result in X inactivation (Tinker, A.V., and C.J. Brown. 1998. *Nucl. Acids Res.* 26:2935–2940). Despite a nuclear RNA accumulation of normal abundance and stability, XIST RNA does not localize in reactivants or in naturally inactive human X chromosomes in mouse/human hybrid cells. The XIST transcripts are fully stabilized despite their inability to localize, and hence XIST RNA localization can be uncoupled from stabilization, indicating that these are separate steps controlled by distinct mechanisms. Mouse *Xist* RNA tightly localized to an active X chromosome, demon-

strating for the first time that the active X chromosome in somatic cells is competent to associate with *Xist* RNA. These results imply that species-specific factors, present even in mature, somatic cells that do not normally express *Xist*, are necessary for localization. When *Xist* RNA is properly localized to an active mouse X chromosome, X inactivation does not result. Therefore, there is not a strict correlation between *Xist* localization and chromatin inactivation. Moreover, expression, stabilization, and localization of *Xist* RNA are not sufficient for X inactivation. We hypothesize that chromosomal association of XIST RNA may initiate subsequent developmental events required to enact transcriptional silencing.

**X** inactivation, the process whereby an entire X chromosome is transcriptionally silenced in mammalian females (Lyon, 1961), has been studied for several decades, yet the underlying process remains largely undefined. A wealth of developmental evidence has implicated mouse *Xist* (and its human counterpart *XIST*) in the process of X inactivation. Recently it has been demonstrated that *Xist* is necessary for X inactivation during development, as mouse X chromosomes deleted for *Xist* failed to inactivate (Penny et al., 1996; Marahens et al., 1997). Transfection of ES cells further showed that *Xist* is sufficient for X inactivation, as an *Xist*-containing cosmid or YAC introduced onto an autosome causes the chromosome to manifest signs of inactivation (Herzing et al., 1997; Lee and Jaenisch, 1997).

The transcript from *XIST* is a large untranslated RNA, which we have shown to have a unique and highly specific location in the nucleus (Brown et al., 1992; Clemson et al., 1996). Unlike other nuclear RNAs, the bulk of the signal is

fully processed message that is stably maintained in a specific site in the nucleus, and is not detected in the cytoplasm. This abundant stable transcript maintains a tight association with the X chromosome, mirroring its shape and size as if painting the chromosome (Clemson et al., 1996). When *Xist* is expressed ectopically through ES cell transfection, the RNA shows the same association with autosomal chromatin after differentiation, which subsequently exhibits hallmarks of inactivation such as heterochromatinization, loss of acetylated histone H4 antibody staining, and inactivation of sequences close to the insertion site (Herzing et al., 1997; Lee and Jaenisch, 1997). The apparent structural association of *XIST/Xist* with the inactive X chromosome (Clemson et al., 1996; Lee et al., 1996; Lee and Jaenisch, 1997) coupled with the evidence revealing *Xist's* importance in the process of X inactivation (Lee et al., 1996; Penny et al., 1996; Herzing et al., 1997; Lee and Jaenisch, 1997) suggests that this RNA is a functional nuclear RNA that is most likely contributing to, by the act of its association, chromatin inactivation. As such, it is predicted that the precise localization of XIST RNA is critical for its function.

Although a spate of recent evidence has clearly demonstrated the absolute requirement for *Xist* in X inactivation

Address correspondence to Christine M. Clemson, Department of Cell Biology, University of Massachusetts Medical Center, 55 Lake Avenue North, Worcester, MA 01655-0106. Tel.: 508-856-6016; FAX: 508-856-5178; E-mail: chris@gene.ummed.edu

during mouse development, studies indicate that its human counterpart, *XIST*, is not essential for X inactivation in cultured somatic cells. Brown and Willard (1994) showed that *XIST* expression is apparently not necessary for maintaining X inactivation in mouse/human somatic cell hybrids, and Rack et al. (1994) showed that inactive X chromosomes in human leukemic cells that had lost the X-inactivation center maintained their inactivation. These results are surprising in light of the fact that *XIST* RNA is constitutively expressed in differentiated female cells, suggesting that it is functioning in the maintenance of chromosome inactivation. Two recent papers show that expression of *XIST* RNA is not sufficient to cause inactivation (Hansen et al., 1998; Tinker and Brown, 1998). In these reports, rodent/human somatic cell hybrids retaining the human active X chromosome were treated with several rounds of 5-azadeoxycytidine (5azadC),<sup>1</sup> a demethylating agent, to create clones that reactivated the dormant *XIST* gene. Using mouse/human hybrids, Tinker and Brown (1998) analyzed eight X-linked genes by reverse transcription (RT)-PCR for expression, with none of the genes showing signs of inactivation as a result of *XIST* expression, while Hansen et al. (1998) reactivated *XIST* expression in hamster-human hybrids, and detected no silencing of two X-linked genes. These results contrast with those that show that forced *Xist* expression caused by mutant methyltransferase results in silencing of X-linked genes in differentiating male ES cells and developing mice embryos (Panning and Jaenisch, 1996).

The requirement for *Xist* in X inactivation in developing mice embryos, coupled with the precise spatial positioning of the RNA to inactivated chromatin (Clemson et al., 1996; Lee et al., 1996; Panning and Jaenisch, 1996), strongly suggests that there is a correlation between *XIST* RNA binding and inactivation of chromatin; however, it is not known conclusively whether *Xist* localization results in X inactivation. Somatic cell hybrids where *XIST* is apparently neither necessary (Brown and Willard, 1994) nor sufficient (Tinker and Brown, 1998) for X inactivation allow us to investigate rigorously whether association of *XIST* RNA with the X chromosome corresponds with transcriptional silencing. Interestingly, our results reveal that the human *XIST* RNA shows aberrant positioning in mouse/human hybrid cells that express the normally dormant *XIST* gene by means of demethylation, in that the RNA is more disperse and spread across the entire nucleus. Even in hybrids that contain a naturally inactive human X chromosome, the *XIST* RNA does not show normal tight localization to the X chromosome. These results suggest that the impotence of *XIST* RNA in this system may be due to failure of the RNA to localize. Furthermore, these results imply that the ability to localize to the chromosome is not an innate feature of the *XIST* RNA, but is afforded by specific factors that are not fully substituted by mouse homologues. We also show that when the RNA is not restricted to the chromosome, it is still found as a large and stable entity, indicating that stabilization and localization

1. *Abbreviations used in this paper:* 5azadC, 5-azadeoxycytidine; rt, room temperature; RT, reverse transcription; Xa, active X chromosome; Xa-*XIST*<sup>+</sup>, reactivated X; Xi, inactive X chromosome; *XIST/Xist*, human/mouse Xi-specific transcript.

of *XIST* RNA are uncoupled. Using demethylation, we then forced expression of *Xist* RNA from the active murine X chromosome, and examined the relationship of the RNA to the X chromosome. Unlike its human counterpart, the murine *Xist* RNA strictly localized to the X chromosome in the mouse background. Since the reactivated *Xist* RNA correctly localized in a somatic cell, we conclude that its association to the X chromosome does not have to occur during development to mark the chromosome for *Xist* RNA binding. Furthermore, *Xist* RNA does not discriminate between active and inactive chromatin as it paints the active X chromosome completely. Additionally, this result shows that the factors required to deposit the RNA at the X chromosome are present and available in terminally differentiated cells that do not normally express *Xist*. Finally, we show that when *Xist* RNA is properly localized to an active mouse X chromosome, inactivation of the X chromosome does not result. This important result clearly demonstrates a lack of correlation between *Xist* RNA localization and inactivation, and also indicates that when removed from the normal developmental context, even with proper expression, stabilization, and localization, *Xist* is not able to cause X inactivation.

## Materials and Methods

### Cells, Cell Culture, and Treatment with 5-Azadeoxycytidine

Derivatives of the AHA-11aB1 active X chromosome-containing human/mouse somatic cell hybrid that expressed human *XIST* RNA were isolated after treatment with successive rounds of the demethylating agents 5-azadeoxycytidine or 5-azacytidine (Tinker and Brown, 1998). To create additional hybrid subclones that expressed the murine *Xist* gene, AHA-A5-2b cells (that stably expressed human *XIST* from the active X chromosome) or subclones that had undergone one round of 5-azadeoxycytidine treatment, but did not stably express human *XIST*, were plated at 10<sup>4</sup> cells/60-mm culture dish and allowed to attach for 4–6 h before being treated with 5-azadeoxycytidine (0.2 μg/ml) or 5-azacytidine (4 μM) for 24 h. Individual colonies were isolated by trypsinization in cloning cylinders, and were plated in 60-mm dishes. Cells were grown to confluence and subcloned, and RNA was isolated for RT-PCR examination of *Xist* expression using primers MX23b and MIX20 (Kay et al., 1993). RT was as described previously (Brown et al., 1990). *Xist*-positive clones were grown for in situ hybridization (see below), and were subjected to one to four additional rounds of treatment with demethylating agents. Cells were maintained in alpha-MEM with 7.5% FCS supplemented with penicillin/streptomycin (GIBCO BRL, Gaithersburg, MD).

### Cell Preparation for In Situ Hybridization

Our standard cell fixation has been described previously (Lawrence et al., 1989), and will be summarized briefly here. Monolayer cells grown on glass coverslips were extracted with 0.5% Triton X-100, 5% vanadyl ribonucleoside complex to preserve RNA (GIBCO BRL) in CSK buffer (Fey et al., 1986) for 2 min on ice. Cells were then fixed in 4% paraformaldehyde for 10 min at room temperature, and were stored in 70% ethanol at 4°C.

### DNA Probes

For fluorescence in situ RNA hybridization, the following probes were used: human *PGKI*, a genomic lambda clone for *PGKI* isolated from the American Type Culture Collection (Rockville, MD) X chromosome library LAOXNL01; *XIST* (human *XIST* probes generously provided by Hunt Willard, Case Western Reserve University): *XIST* G1A, an ~10-kb genomic plasmid spanning from the fourth intron to the 3' end of the human *XIST* gene; pXISTHbC1A, a 1.6-kb cDNA clone wholly contained within exon 1 of *XIST*; mouse *Xist*, a genomic clone spanning exons 5 and 6 (generously provided by Barbara Panning, Massachusetts Institute of

Technology); mouse *Zfx*: PDP1115, a 6.8-kb cDNA obtained from American Type Culture Collection (cat. 63069); and mouse *Pgk1*, a 1.8-kb cDNA clone from American Type Culture Collection (cat. 57222).

### *In Situ Hybridization and Detection*

Hybridization and detection was performed as described previously (Lawrence et al., 1988; Johnson et al., 1991; Xing et al., 1993), so will be detailed briefly here. DNA probes were nick-translated using biotin-11-dUTP or digoxigenin-16-dUTP (Boehringer Mannheim Corp., Indianapolis, IN). For exclusive RNA hybridization, cells were hybridized under nondenaturing conditions (such that cellular DNA was not accessible) overnight at 37°C in 50% formamide, 2× SSC using a probe concentration of 5 µg/ml. Human and mouse Cot-1 was included in the hybridization buffer for the human and murine RNA hybridizations, respectively. Posthybridization washes were performed as follows: 50% formamide, 2× SSC for 30 min at 37°C; 2× SSC for 30 min at 37°C; 1× SSC for 30 min at room temperature (rt) with agitation; and 4× SSC for 30 min at rt with agitation. Hybridization was detected with either antidigoxigenin antibody (Boehringer Mannheim Corp.) coupled with rhodamine or fluorescein at 200 µg/ml in 1% BSA, 4× SSC; or with fluorescein-conjugated avidin (Boehringer Mannheim) at 2.5 mg/ml in 1% BSA, 4× SSC for 1 h at 37°C. Postdetection washes were performed as follows: 4× SSC for 20 min at rt with agitation, 4× SSC, 0.1% Triton for 20 min at rt with agitation, and then 4× SSC for 20 min at rt with agitation. Simultaneous detection of whole X chromosome library and RNA was done as described previously (Clemson et al., 1996) with minor modifications. In brief, cells were extracted and fixed as described above. The cells were then denatured at 80°C for 5 min in 70% formamide, 2× SSC, and were immediately dehydrated through an ice-cold ethanol series. The Total Chromosome Library Coatasome X-digoxigenylated™ (Oncor Inc., Gaithersburg, MD) was used for human chromosome detection, while biotinylated Mouse Chromosome X Paint Probe™ (Oncor Inc.) was used for mouse X detection. 10 µl of the whole chromosome probe mixture was denatured for 10 min at 75°C, and was allowed to preanneal for 30 min at 37°C. Samples were then hybridized with the probes overnight at 37°C. Posthybridization washes, detection, and postdetection washes were performed as described above.

### *XIST Stability*

To determine the stability of the XIST RNA in hybrid and lymphoblast cells, recently expanded cultures were treated with 2 µg/ml actinomycin in DMSO. Control cultures received only the DMSO. Whole-cell RNA was harvested by acid guanidinium thiocyanate-phenol-chloroform extraction (Chomczynski and Sacchi, 1987), and was reverse-transcribed as above. cDNA was diluted 1/100, and RT-PCR was performed with primers for *XIST* (Duncan et al., 1993) and actin (Chariot and Castronovo, 1996). PCR was demonstrated to be in a linear range of amplification, and then products were quantitated with NIH Image (National Institutes of Health, Bethesda, MD) on scanned images of ethidium bromide-stained agarose gels. To determine the half-life of the *XIST* message, the ratio of *XIST* product to actin product was divided by the average ratio of *XIST* to actin in the DMSO-treated samples. The half-life of actin cytoplasmic mRNA has previously been estimated at 25 h (Sympson and Geoghan, 1990), and therefore should not have varied substantially over the 8-h time period used here. Stability of XIST RNA in hybrids was also determined by hybridization to slot blots, and yielded a similar half-life of 5 h (data not shown).

For quantitation of relative amounts of XIST RNA, hybridization of slot blots was performed as described previously (Brown et al., 1991). In brief, total cellular RNA was prepared using the guanidinium thiocyanate method. 5 µg of RNA from an inactive X-containing somatic cell hybrid (Willard et al., 1993) and the AHA-11aB1 active X-containing hybrid as well as its demethylated XIST<sup>+</sup> derivative was denatured in formaldehyde/formamide and transferred to a nitrocellulose membrane. The blots were hybridized with <sup>32</sup>P-labeled XIST14A cDNA probe (Brown et al., 1991). To control for loading equal amounts of the RNA, the blots were stripped and probed for the MIC2 gene.

### *Imaging*

Large numbers of cells were evaluated by direct examination using an Axio-phot microscope equipped with an 100× 1.4 NA Plan ApoChromat™ objective (Carl Zeiss Inc., Thornwood, NY) with a 2.5× photo eyepiece and multibandpass epifluorescence filters (Chroma, Brattleboro, VT).


Representative images were recorded with a CCD camera (200 series; Photometric, Tucson, AZ) with a pixel size of 19 µm and a 14-bit A/D converter (data acquisition system by G.W. Hannaway and Associates, Boulder, CO).

## **Results**

### *Human XIST Transcripts in Hybrid Cells Show Abnormal Nuclear Localization*

In other recent work (Tinker and Brown, 1998), subclones of AHA-11aB1 mouse/human somatic cells (AHA hybrids) were treated with three rounds of demethylation, each of which involved culturing cells for 24 h with 5azadC. After removing the 5azadC and culturing cells in normal media, many clones were isolated that stably expressed RNA from the normally dormant *XIST* gene on the active human X chromosome. Subsequent RT-PCR analysis showed that none of eight X-linked genes examined were inactivated as a result of *XIST* expression from the active X chromosome. As it was possible that some of the cells in the population examined might have lost *XIST* expression, and that the gene expression detected by RT-PCR could have been derived from this subset of cells, here we used in situ hybridization to confirm the lack of inactivation at a single cell level. Using genomic probes to the human genes, we examined expression of PGK-1 and XIST RNA in the AHA-A5-2b mouse/human hybrid cells that contain a human X chromosome expressing RNA from the hypomethylated *XIST* gene. PGK-1 RNA was detected in the majority of the cells that also expressed XIST RNA, confirming the PCR data from Tinker and Brown (1998) that the X chromosome is not inactivated as a result of reactivated *XIST* expression.

To begin our analysis of the correlation between *XIST* localization and inactivation of chromatin, we examined the distribution of the XIST RNA in AHA mouse/human hybrids that had previously been identified as stable *XIST* expressers by RT-PCR. Using in situ hybridization, we analyzed a total of six different hybrid clones, and found that almost every cell in each population was expressing XIST RNA. Furthermore, the hybrids continue to show stable *XIST* expression for over 2 yr after the original demethylation, confirming that transcriptional reactivation of the human *XIST* locus is quite stable (Tinker and Brown, 1998). The XIST RNA signal in the reactivated hybrid cells was similar in brightness and apparent abundance to normal diploid fibroblasts (Fig. 1, A and D), and the particulate nature of XIST RNA, which we have reported previously in normal fibroblasts (Clemson et al., 1996), is highly apparent in the AHA hybrids (Fig. 1, D–F). However, while the distribution of the XIST RNA in normal diploid fibroblasts is discrete and highly contained in a small area in the nucleus (Fig. 1 A), the reactivated XIST RNA in the hybrids was distinctly different (Fig. 1, D–F) in that it was more spread out, occupying a much larger area of the nucleus. This less-contained pattern revealed numerous tiny distinct particles that spread over a significant portion of the nucleus. In normal diploid fibroblasts, the XIST RNA associates closely with the inactive X chromosome in a pattern that is similar in both size and shape (Fig. 1, A–C; Clemson et al., 1996). While the dispersed



**Figure 2.** Reactivated and endogenous *XIST* expression is similar in hybrid cells. Total cellular RNA from the following cells were transferred to nitrocellulose and hybridized with an *XIST* cDNA probe: AHA-11aB1, the active X-con-

taining hybrid cells (*Xa*); AHA-2C-5C-9, the demethylated derivative that ectopically expresses *XIST* (*Xa-XIST*<sup>+</sup>); and t11-4Aaz5 cells containing a human inactive X chromosome that normally expresses *XIST* (*Xi*). The blots were scanned, and the density of signal from the *Xi* and *Xa-XIST*<sup>+</sup> clones were essentially identical, demonstrating that the RNA is produced in similar quantities from endogenously expressing and demethylated *XIST* genes.

pattern of *XIST* RNA in the hybrid cells suggested that it was not closely associating with the X chromosome, it is possible that the human X chromosome in a mouse background also occupies a broad, more disperse territory. To examine the specific relationship of the *XIST* RNA to the X chromosome in the AHA hybrid cells, the RNA signal was directly compared with that from a whole X chromosome library hybridization in 100 cells in three separate experiments. In >90% of the AHA reactivants, the RNA signal was obviously not strictly associated with the X chromosome (Fig. 1 *E*). While there generally was some density of *XIST* RNA signal over the chromosome, a large quantity of signal was usually seen spreading out away from the X chromosome.

To determine if the lack of localization is due to artifacts created by reactivating *XIST* expression from the active X chromosome (e.g., the active X chromosome may be refractory to *XIST* binding), we examined the distribution of *XIST* RNA in t86-B1maz1b-3a mouse-human hybrid cells that retain a normal human inactive X chromosome (Willard et al., 1993). In over 100 cells examined in four separate experiments, the distribution of the *XIST* RNA in the hybrid cells was identical to that in the AHA-A5 hybrids; the *XIST* RNA was disperse, and spread into the nucleoplasm with no strict limitation to the X chromosome library signal (Fig. 1 *F*). This result indicates that the lack of localization is not an artifact of global demethylation, as these cells were not treated with demethylating agents, and still show no detectable localization of the *XIST* RNA to the X chromosome. It also strongly suggests that the aberrant localization is not due solely to inherent differences between active and inactive X chromosomes, as the human X chromosome in this cell line is an inactive X chromosome known to remain transcriptionally inert (Brown et al., 1997).

Since the signal intensities in the different cell lines seemed equivalent (Fig. 1, *A*, *D*, and *F*), it appeared that hybrid cells that ectopically express *XIST* produce similar amounts of RNA, as do diploid cells that express *XIST* endogenously. However, we wished to confirm that aberrations in expression from the reactivated *XIST* gene do not cause the mislocalization. For example, demethylation could produce an overabundance of *XIST* RNA, causing the transcripts to spill from the site of transcription into the surrounding nuclear space. Alternatively, reactivation could produce quantities of RNA that are insufficient to


paint the chromosome. This possibility was addressed through a slot blot hybridization assay (Fig. 2). *XIST* RNA from a hybrid cell line containing an active X (*Xa*), reactivated X (*Xa-XIST*<sup>+</sup>), and normally expressing inactive X chromosome (*Xi*) were compared. No *XIST* RNA was detected from the active X chromosome, while a similar abundance of RNA was observed in the *Xa-XIST*<sup>+</sup> and *Xi* cell lines. This result confirms the *in situ* hybridization data that suggested similar amounts in the different somatic cells examined, and clearly shows that *XIST* reactivation by demethylation produces essentially the same levels of *XIST* RNA as the *XIST* gene on the naturally inactivated X chromosome. This result further suggests that lack of localization is not due to artifacts created by demethylation.

### *Stability and Localization of XIST RNA are Uncoupled*

Next we examined the stability of the *XIST* RNA in hybrid cells, as previous results indicate that this nuclear RNA is highly stable (Clemson et al., 1996), and that nuclear stability is an important developmentally regulated feature of this RNA (Panning et al., 1997; Sheardown et al., 1997). In XX somatic cells, only the inactive X chromosome expresses *XIST* (Brown et al., 1992; Clemson et al., 1996; Lee et al., 1996), and the transcripts that coat the chromosome form a nuclear accumulation much larger than other mRNAs examined (Clemson et al., 1996). This large *XIST* RNA accumulation was shown to be composed of two populations: a small focus of short-lived nascent transcripts found at the inactive X chromosome simply because it is transcribed there, and a large spliced and stable population that represents the mature, likely functional RNA (Clemson et al., 1996). Before X inactivation in female cells or undifferentiated ES cells, *XIST* RNA is observed as tiny dots on both X chromosomes, or as a single dot in male cells (Lee et al., 1996; Panning and Jaenisch, 1996; Panning et al., 1997; Sheardown et al., 1997). Very recently it has been shown that during early development, *XIST* RNA is initially expressed in an unstable form from both active and inactive X chromosomes, and becomes stabilized from the inactive X as development proceeds. The transition from single dot to large accumulation is not regulated by increased transcription, but by stabilization of the *XIST* transcripts in *cis* from the inactive X chromosome (Panning et al., 1997; Sheardown et al., 1997).

It is not known whether deposition and stabilization of *XIST* RNA arise from a single mechanism, or if separate mechanisms are involved in stabilizing and positioning the RNA. Analysis of *XIST* RNA in hybrids provides important insights in this regard. In all six hybrid lines examined, it was clear that the *XIST* RNA comprised a large accumulation, with the quantity of signal indistinguishable from that observed in normal cells (Figs. 1 and 2), suggesting that the *XIST* RNA represents a unique *XIST* intermediate, one in which the transcripts are expressed and accumulating, but not localizing to the chromosome. These results are the first to indicate that association of *XIST* RNA with the X chromosome does not automatically arise from the increased stability that occurs after differentiation.

To examine stability by another approach, we compared



**Figure 3.** Stability of the XIST RNA after treatment of cells with actinomycin. Cells were harvested after 0–8 h of actinomycin treatment (A) or control DMSO treatment (D) as the actinomycin is dissolved in DMSO. RT-PCR was performed on the RNA isolated from these cells with primers for *XIST* (A and B) and actin (not shown). The RT-PCR products for a female lymphoblast (GM07350; A) and an inactive X-containing human/mouse somatic cell hybrid (t75-2maz34-1a; B) are shown, and the ratio of *XIST* product to actin product (normalized to the ratio of *XIST* to actin for the control DMSO treatments) is plotted in C. The black line shows the ratio for the female cells, while the grey line shows the ratio for the hybrid cell line.



the half-life of XIST RNA after actinomycin D treatment in unsynchronized human female lymphoblasts (GM07350), where the transcripts are expected to be stable, and in the mouse/human hybrid cell line t75-2maz-34-1B that expresses *XIST* from the inactive human X chromosome. Fig. 3 shows the results of RT-PCR with primers for human *XIST* from RNA isolated from cells after up to 8 h of actinomycin treatment. By normalizing to RT-PCR products for actin mRNA (which has a cytoplasmic half-life of ~25 h [Sympson and Geoghan, 1990]) the half-life of the RNA from the two cell lines could be determined. Both the hybrids and the lymphoblasts showed a half-life for XIST RNA of ~5 h (Fig. 3 B), which is similar to the half-life reported for stable XIST transcripts in diploid fibroblasts (Sheardown et al., 1997). The similarity of the half-life in the two cell lines confirms that the transcripts in the hybrids are stable. The similar half-lives, but very different patterns of XIST RNA localization in the nucleus, strongly suggest that stabilization of XIST RNA does not arise from chromosomal localization, and vice versa. Therefore, some additional mechanism other than stability is responsible for localizing the XIST transcripts to the X chromosome.

We speculate that localization of this RNA to the X chromatin is brought about by specific autosomal factors that allow deposition of the human XIST transcripts along the chromosome. As there is ~30% sequence divergence between mouse and human *Xist/XIST* sequences (Brockdorff et al., 1992; Brown et al., 1992), it is not surprising if the homologues required for localization of Xist RNA are unable to substitute completely for the human factors. In the hopes of finding a human autosome that would allow for XIST localization, and ultimately to identify potential localization factors, we examined the distribution of XIST RNA in six different hybrids containing a human inactive X chromosome. As these hybrids contained various other

human chromosomes, we hoped to identify a hybrid that contained the necessary human factor to allow localization of human XIST. None of the hybrids showed localized XIST RNA, although all exhibited large apparently stable XIST RNA accumulations. The consistent failure of hybrid cells to localize the RNA correctly suggests that either the necessary locus was missing from these hybrids, or that more than one factor is required concurrently, and that these hybrids did not have the right combination of human chromosomes. We favor the latter possibility, as these hybrids have been shown by PCR or karyotyping to include collectively most of the human chromosome complement.

#### **Reactivated Mouse *Xist* RNA Localizes to the Active X Chromosome**

If the failure of the human XIST RNA to localize is the result of mouse/human species differences, then it follows that if murine *Xist* were reactivated, the RNA would be properly localized as the full complement of murine factors would be present. To test this hypothesis, we isolated clones that expressed not only human XIST RNA, but also mouse Xist RNA after additional rounds of demethylation. We studied the nuclear Xist/XIST RNA expression pattern in AHA-A5-2b-g cells, AHA hybrids expressing human XIST that had been treated with additional rounds of demethylating agents to induce mouse *Xist* expression, providing an opportunity to compare directly mouse and human XIST RNA in the same cell. The mouse cells used in the generation of these hybrids (A9) do not express Xist RNA without treatment; and these cells maintained ectopic murine *Xist* expression less consistently than human XIST expression (see below). In our analysis of 80 cells expressing both human and mouse *Xist*, >70% showed the



following pattern of expression: the human XIST RNA showed a small focal concentration, with particles apparently drifting away from the site of transcription into the rest of the nucleoplasm; in contrast, the murine Xist RNA signal was a large but discrete pattern that was much more confined in the nucleus than its human counterpart (Fig. 4, *A* and *B*). We directly compared the distribution of the murine Xist signal to the mouse chromosome library signal in 50 different cells. Like the human XIST RNA in its normal human diploid environment, the murine transcripts formed a punctate accumulation very similar in size and shape to the mouse X chromosome paint, revealing a close association with the active murine X chromosome (Fig. 4, *C–E*). Clearly, murine Xist RNA can properly localize in the mouse background, while in the same cell the human XIST RNA cannot.

These results support the hypothesis that the aberrant human XIST RNA location is due to a lack of species-specific factors necessary for localization. Interestingly, it is apparent that the factors required to localize the Xist RNA are present in terminally differentiated cells that normally do not express Xist RNA. This important result is informative on several other levels. First, it shows that the lack of localization of the human XIST was not due to 5azadC treatment, as the localization of the murine Xist was observed after 5azadC treatment; hence, demethylation does not disallow localization of the transcripts. Second, the mislocalization is not due to general disruption of nuclear structure in hybrid cells. Perhaps most importantly, it shows that the XIST RNA does not discriminate between active and inactive X chromosomes. In these mouse cells the murine X chromosome has already been selected in early embryogenesis to remain active, but is

able to stably express and bind the Xist RNA. Hence, active X chromosomes are not recalcitrant to Xist RNA binding. The fact that the reactivated murine Xist localizes unequivocally shows that the proper deposition of this RNA at the X chromosome can occur outside the realm of early development, and therefore, that no developmental factors are required to set up a pattern of Xist chromosome binding. Apparently all the requirements for expression, stability, and localization of Xist RNA are available in somatic cells.

### *The Mouse X Chromosome Painted by Xist RNA Does Not Inactivate*

Studies in transgenic mice have definitively demonstrated that *Xist* is not only necessary (Penny et al., 1996; Marahens et al., 1997), but also sufficient (Herzing et al., 1997; Lee and Jaenisch, 1997) to initiate X inactivation in developing mouse embryos. In our analysis, we have circumvented development by recreating, in terminally differentiated cells, several of the steps known to correlate with X inactivation in embryonic cells: expression of Xist RNA (Kay et al., 1993; Kay et al., 1994); transition from a single dot of expression to stabilization of a large nuclear RNA accumulation (Panning et al., 1997; Sheardown et al., 1997); and deposition of these stable Xist transcripts along the entire length of the chromosome (Clemson et al., 1996). The obvious next question is, does inactivation of the chromatin result from properly expressed, stabilized, and localized Xist RNA?

While we were able to isolate stable long-term human XIST reactivants after three rounds of treatment with 5azadC, we were not able to isolate a clone that expressed

*Figure 1.* Fluorescence in situ hybridization detection of XIST RNA in normal human and mouse/human hybrid cells. Digoxigenin and biotinylated probes were hybridized in situ and detected with fluorochrome-conjugated avidin or antidigoxigenin antibody. Fluorochromes used were FITC (*green*), rhodamine (*red*), and DAPI (*blue*). (*A*) In normal human diploid fibroblasts (46, XX WI-38 cells), the XIST RNA (*green*) occupies a discrete location in the nucleus. (*B*) The XIST RNA has a similar shape to the X chromosome territory defined by the whole X chromosome library signal (*red*); the weak red signal in the middle of this cell is the active X chromosome which is out of the plane of focus). (*C*) The XIST RNA essentially paints the X chromosome in these normal nuclei (overlap is yellow). (*D*) In mouse/human hybrids containing a single active human X chromosome (AHA-A5) that expresses XIST through treatment with the demethylating agent 5azadC, the XIST RNA (*red*) is dispersed, and spreads throughout the nucleus. (*E*) The XIST RNA (*red*) does not associate or strictly localize to the human whole X chromosome library signal (*green*) in the hybrid cells. (*F*) In mouse/human hybrid cells (t86-B1maz1b-3a) that contain a single human inactive X that expresses XIST endogenously (no 5azadC treatment), the human XIST RNA (*red*) also shows aberrant localization to the whole X chromosome library signal (*green*).

*Figure 4.* In situ RNA and murine X whole chromosome library detection in AS-2B mouse/human hybrid cells that contain both an active human and mouse X chromosome. Both human and murine Xist RNA is expressed from these chromosomes as a result of 5azadC treatment. (*A* and *B*) The difference between the reactivated murine Xist RNA and human XIST RNA is illustrated in these cells, with the murine RNA appearing highly localized (*red*) and the human XIST RNA (*green*) showing a focus of expression (most likely representing the site of transcription) with particles that drift away into the nucleoplasm. Reactivated murine Xist RNA localizes to the active X chromosome (*C–E*). (*C*) The murine Xist RNA (*red*) is discretely contained in the hybrid cell nucleus (*D*) in a pattern similar to the murine X chromosome signal (*green*; the mouse X chromosome library cross-reacts slightly with the human X chromosome, giving a second green signal apparent in the middle cell in *D*); (*E*) the murine Xist RNA colocalizes with the active murine X chromosome (overlap is yellow).

*Figure 5.* The mouse X chromosome painted by the Xist RNA does not inactivate. Codetection of murine Xist RNA (*green*) and Zfx RNA or Pgl1 RNA (*red*) in undenatured 1B-5C-3B cells. (*A* and *B*) Zfx RNA (*red*) is expressed from cells that also express mouse Xist RNA (*green*); (*C*) Pgl1 RNA expression (*red*) is also expressed from chromosomes that express Xist RNA (*green*), suggesting that the active X chromosome is not inactivated as a result of Xist localization. Expression of mouse genes appears to be unaffected in cells that express mouse Xist. (*A*) There is a similar pattern and intensity of expression of Zfx RNA in cells that express mouse Xist (*top*), and those that do not (*bottom*). (*C*). Similarly, there is comparable Pgl1 expression in cells that express mouse Xist (*left*) and cells that do not (*far right*).

mouse *Xist* long-term, despite multiple serial treatments with demethylating agents. After selection and culture of single cell isolates, the level of *Xist* RNA declined over a period of several weeks as detected by RT-PCR. By in situ techniques we saw that initially almost every cell in the population expressed *Xist*. Gradually, the number of cells that expressed *Xist* declined (Fig. 4 B), until eventually no more cells expressed *Xist* RNA.

While we initially thought that this reduction was due to some selective disadvantage conferred by mouse *Xist* expression (for example, the cells could be dying because the mouse *Xist* RNA inactivated the sole murine X chromosome), several considerations make this possibility less likely. First, we detected no change in cell growth rate or evidence of large scale cell death in the reactivated hybrids, both of which would be expected if *Xist* expression was causing large-scale toxicity. Second, we have isolated clones that continued to express mouse *Xist* in >30% of the population for as long as 2 mo after the initial demethylation. Additionally, we were able to subclone pure populations of *Xist* RNA expressers from clones that had reduced expression of *Xist* to only a few cells, which would not be expected if the cells were gradually dying. While it is possible that the reactivants were selected against, we consider it more likely that *Xist* was being remethylated gradually, much like what was seen for the unstable human reactivants (Tinker and Brown, 1998). Our failure to obtain a clone that expressed mouse *Xist* longer than several months, in contrast to the human gene, may reflect differences between the methylation of the two genes; for example, the murine gene may have more methylation which must be completely removed before remethylation occurs, causing loss of *Xist* expression. Ultimately, the difficulty we had in creating long-term mouse *Xist* reactivants did not cause a problem for our experiments, as any cells that showed detectable mouse *Xist* always showed the same large intense pattern, allowing us to perform analysis of individual cells, even when as little as 10% of the population expressed *Xist* RNA.

The definitive way to assess whether the mouse X chromosome painted by reactivated *Xist* RNA is inactivated is to examine expression of individual genes in single cells. Because of the heterogeneity of *Xist* expression in our cell population, RT-PCR analysis could not address the critical question of whether individual genes were still expressed. We therefore used dual in situ detection of murine *Xist* with X-linked genes to examine directly X inactivation in individual cells. *Pgk1* is subject to X inactivation in humans and mice (Disteche, 1995), while *Zfx* is an X-linked gene that escapes inactivation in humans, but is subject to X inactivation in the mouse (Adler et al., 1991). *Zfx* and *Pgk1* nuclear RNA expression were examined in single cells that expressed *Xist* RNA (Fig. 5, A–C), using cDNA clones. In multiple experiments encompassing over 200 hybrid cells, these two genes were clearly transcribed in cells that were also expressing and localizing *Xist* RNA. While we were also able to detect the human RNA with the *Pgk-1* probe, we included in our analysis only those signals that were closely associated with the mouse *Xist* RNA signal, confirming that it was transcribed from the mouse X.

To evaluate whether the *Xist* RNA localization resulted in a more subtle transcriptional downregulation of the

genes vs. a complete silencing of the X chromosome, we compared the distribution of both *Pgk1* and *Zfx* RNA from cells either expressing or not expressing murine *Xist* RNA. As exemplified in Fig. 5, A and C, the *Pgk1* and *Zfx* RNA signals were not discernibly different in cells devoid of *Xist* expression vs. those in which the mouse X is painted by *Xist* RNA, suggesting that the murine X chromosome has not undergone detectable transcriptional downregulation as a result of *Xist* expression and localization. While we cannot rule out the possibility that, over the long term, progressive inactivation will occur, expression of *Zfx* and *Pgk-1* was analyzed in subclones that expressed and localized murine *Xist* RNA for over 2 mo. Therefore, our results show that *Xist* RNA can localize to the mouse X chromosome without inactivation of specific genes for many generations. We conclude that the association of *Xist* RNA with chromatin is not strictly correlated with transcriptional silencing. Furthermore, these results indicate that expression, stabilization, and proper localization of *Xist* RNA in somatic cells is not sufficient to cause inactivation.

## Discussion

Understanding the relationship of XIST RNA localization to X inactivation and the mechanisms governing localization to the chromosome is likely key to understanding the process of X inactivation itself. Results presented here show that despite a nuclear RNA accumulation of normal abundance and stability, XIST RNA does not localize to reactivated or inactive human X chromosomes in mouse/human hybrid cells. To determine if the improper location of RNA in these hybrids explained the lack of subsequent inactivation, we created clones that stably expressed not only the human *XIST*, but mouse *Xist* as well. Distribution of the mouse *Xist* RNA in these hybrids clearly showed a tight association with the active murine X chromosome, in marked contrast to the nonlocalized human *XIST* RNA in the same cells. This important result also allows several conclusions: (a) demethylation with 5azadC does not affect the ability of *Xist* RNA to localize, since the murine *Xist* RNA does localize properly in the same cells; (b) expression of this RNA does not have to begin in early development to allow localization; and (c) *Xist* RNA does not discriminate between the developmentally selected active and the inactive X chromosome. Collectively, these results indicate that the most likely reason for the lack of localization to the human X chromosome is mouse/human species differences. Since completion of this work, it was shown that inactive human X chromosomes, when inserted into mouse embryonal carcinoma cells, activate expression of many genes, yet fail to stop expressing XIST RNA (Yoshida et al., 1997). Results presented here indicate that the XIST RNA may not closely associate with the chromosome in the mouse environment.

As many pre-mRNAs are most concentrated near their site of transcription (Xing and Lawrence, 1993; Xing et al., 1995), it is not clear to what extent limited localization of the XIST RNA with the X chromosome in the hybrids is simply a consequence of its transcription, or if it reflects a weak affinity for the chromosome. For example, the XIST RNA may be transcribed but unable to localize to the chromosome at all, causing the transcripts to drift away



gradually into the nucleoplasm. Alternatively, the murine factors may be able to localize the human transcripts, but because of either reduced affinity or availability, allow only a small fraction of the transcripts to associate with the X chromosome.

### ***Localization of XIST RNA Does Not Arise from Stability***

Recent results showed the transition of Xist RNA from a small dot to an abundant accumulation in early development, and suggested that the increased stability allowed the transcripts to localize to the inactive X chromosome (Panning et al., 1997; Sheardown et al., 1997). Our examination of the stability of the human RNA in human/rodent hybrids indicates that the transcripts are fully stabilized despite the inability to localize to the X chromosome. Hence, localization of XIST RNA does not result from increased stability. Our results make it clear that the process required for transcript stabilization can occur after development as the half-life of XIST RNA in these hybrids is similar to that recently reported in normal diploid fibroblasts (Sheardown et al., 1997).

### ***Differing Impact of Xist Expression Throughout Development***

The observation that the chromosome is completely painted by the Xist RNA, but still shows normal expression of both Zfx and Pcg1 RNA in the same cell shows that the chromosome is still transcriptionally active, and that there is no strict correlation between Xist RNA chromosomal localization and inactivation of chromatin. We conclude that expression, stabilization, and localization of Xist RNA is not sufficient for X inactivation. Importantly, these results also suggest that there is a difference between the role of *Xist*/*XIST* in immature vs. fully differentiated cell types. In creating a system where a single copy of the *Xist* gene produces RNA that is stably localized in fully differentiated cells, we are able to compare our results with those that show the absolute requirement for *Xist* in X inactivation (Penny et al., 1996; Marahens et al., 1997) and the sufficiency of *Xist* with limited surrounding sequences to cause silencing (Herzing et al., 1997; Lee and Jaenisch, 1997). The critical difference is that in these previous studies, *Xist* is ectopically expressed in embryonic cells before differentiation, while in experiments reported here, *Xist* expression is forced in fully differentiated somatic cells. The contrasting results suggest that XIST RNA is fully competent to enact chromatin inactivation during embryogenesis, but later in development loses this ability either through lack of contributing factors or a change in the RNA itself.

Although we have demonstrated in this paper that association of Xist RNA with the X chromosome is not sufficient for transcriptional silencing, the supposition that localization is required for X inactivation is most likely still correct. The process of X inactivation likely involves several steps, including the expression, stabilization, and localization of Xist RNA. Our results not only demonstrate that the localization of XIST RNA with the chromosome does not necessarily culminate in transcriptional silencing,

but also suggest that an additional step or mechanism is required to complete the process of X inactivation.

### ***Potential Role of XIST RNA Localization During Development***

Based on our observations, we postulate a model in which the XIST RNA is involved in the process of X inactivation, but is not directly responsible for transcriptional downregulation. In this model, we speculate that the Xist RNA serves to mark or potentiate the chromosome for a subsequent step or process. For example, the RNA, by virtue of its binding along the length of the chromosome and its affinity for certain factors, could recruit processes that ultimately result in transcriptional silencing. While there are many examples of processes that could act in concert with Xist RNA, we consider two for discussion: methylation and histone deacetylation. Both are established mechanisms of epigenetic imprinting strongly linked with inactive X chromatin; they are stably maintained and transferred to daughter cells, and are involved with sites along the entire inactive X chromosome. Furthermore, methylation (Lock et al., 1987; Keohane et al., 1996) and overall deacetylation of the inactive X (Keohane et al., 1996) occur after *Xist* expression early in embryogenesis. Additionally, this idea may explain why *Xist* is not necessary for maintaining X inactivation in somatic cells, while other results show that it is required during development. XIST RNA could be absolutely required during embryogenesis to establish the pattern of methylation or histone deacetylation by recruiting factors like methyltransferases or deacetylases to the inactive X chromosome; however, once such a pattern was established on the inactive X, then the XIST RNA would no longer be rigorously required. The constitutive expression and localization of XIST RNA (Clemson et al., 1996) in adult cells may serve as a backup mechanism to ensure that epigenetic silencing is rigorously maintained. Ultimately, this model could explain why the *XIST* gene could be deleted in vitro from somatic cells with no obvious effect (Brown and Willard, 1994); i.e., if the primary source of inactivation maintenance was a process prone to little error, (such as maintenance methyltransferase which has an estimated efficiency of 99.9% [Pfeifer et al., 1990]), then as a backup mechanism XIST RNA could be deleted without obvious phenotype in cultured cells. In contrast, in vivo, where even a slight failure of X inactivation would be deleterious, XIST RNA would be needed as a constant fail-safe mechanism.

Of course, if *XIST*/*Xist* initiates subsequent events that ultimately silence the chromosome, then why didn't our reactivated mouse/human hybrid system succumb to transcriptional silencing? It is possible that cells that did inactivate were selected against, allowing us only to detect cells that managed to circumvent X inactivation. This possibility appears less likely because we did not observe any evidence of cell death or change in growth rate, combined with the fact that these cells retained an active human X chromosome, which could compensate for the loss of the single mouse X. Rather, we favor the possibility that the resistance to X inactivation is due to the fact that the Xist RNA was not expressed before differentiation of these cells; e.g., it may be very difficult or impossible to establish

de novo inactivation in cells after development as the required enzymes may not be available. For example, the levels and location of de novo and maintenance methyltransferases differ in early embryonic vs. terminally differentiated cells (Razin and Shemer, 1995; Turker and Bestor, 1997).

To reactivate expression of *XIST/Xist*, the cells were treated with a demethylating agent that can impact cell viability as well as replication timing and condensation of the inactive X chromosome (Gregory et al., 1985; Haaf et al., 1988). Additionally, the new methylation patterns may be slowly incorporated into DNA (Toth et al., 1989). Although we were able to produce clones that expressed human *XIST* indefinitely, we could not isolate clones that expressed murine *Xist* longer than several months. Hence, we cannot rule out the possibility that this was not enough time to set up adequate patterns of genomic imprinting that result in silencing of the previously active X-linked genes like *Zfx* and *Pgk1*. However, the cells were only cultured in 5azadC for a short period of time (24 h) to cause expression of the mouse or human *Xist*, and were subsequently cultured without additional 5azadC for as long as several months with a significant fraction (>30%) still expressing mouse *Xist*. Expression of the X-linked genes *Zfx* and *Pgk1* was analyzed as much as 8 wk after 5azadC treatment, making it less likely that their continued expression is the result of demethylation. Reactivation of normally inactivated genes by 5azadC is extremely rare in hybrids ( $\sim 10^{-4}$ ) or normal ( $10^{-7}$ ) cells (Gartler and Goldman, 1994). It is interesting to speculate that higher susceptibility to reactivation in hybrids is related to mislocalization of *XIST* RNA. Since X inactivation is well-maintained in hybrids, it is unlikely that the lack of de novo X inactivation is due merely to repression of X inactivation in the mouse-human hybrid system studied here. It is clear that our reactivated hybrid system reconstitutes many steps known to be involved in normal X inactivation; namely expression, stabilization, and localization of *Xist* RNA. Therefore, it seems that we have created an intermediate to chromosomal inactivation that would not have been seen otherwise; one that makes it possible to analyze and dissect the separate steps involved in X inactivation.

The authors wish to thank John McNeil for his expert computer assistance, Meg Byron and Sarah Baldry for their technical assistance, and Patrick Brown for his help on several hybrid in situs.

This work was supported by National Institutes of Health grants GM49254 and GM53234 to J.B. Lawrence and Medical Research Council MT13690 (Canada) to C.J. Brown.

Received for publication 11 November 1997 and in revised form 4 May 1998.

## References

- Adler, D.A., S.L. Bressler, V.M. Chapman, D.C. Page, and C.M. Disteche. 1991. Inactivation of the *Zfx* gene on the mouse X chromosome. *Proc. Natl. Acad. Sci. USA*. 88:4592–4595.
- Brockdorff, N., A. Ashworth, G.F. Kay, V.M. McCabe, D.P. Norris, P.J. Cooper, S. Swift, and S. Rastan. 1992. The product of the mouse *Xist* gene is a 15 kb inactive X-specific transcript containing no conserved ORF and located in the nucleus. *Cell*. 71:515–526.
- Brown, C.J., A. Ballabio, J.L. Rupert, R.G. Lafreniere, M. Grompe, R. Tonlorenzi, and H.F. Willard. 1991. A gene from the region of the human X inactivation centre is expressed exclusively from the inactive X chromosome. *Nature*. 349:38–44.
- Brown, C.J., L. Carrel, and H.F. Willard. 1997. Expression of genes from the

- human active and inactive X chromosomes. *Am. J. Hum. Genet.* 60:1333–1343.
- Brown, C.J., A.M. Flenniken, B. Williams, and H.F. Willard. 1990. X chromosome inactivation of the human *TIMP* gene. *Nucl. Acids Res.* 18:4191–4195.
- Brown, C.J., B.D. Hendrich, J.L. Rupert, R.G. Lafreniere, Y. Xing, J.B. Lawrence, and H.F. Willard. 1992. The Human *XIST* gene: analysis of a 17 kb inactive X-specific RNA that contains conserved repeats and is highly localized within the nucleus. *Cell*. 71:527–542.
- Brown, C.J., and H.F. Willard. 1994. The human X inactivation center is not required for maintenance of X chromosome inactivation. *Nature*. 368:154–156.
- Chariot, A., and V. Castronovo. 1996. Detection of *HOXA1* expression in human breast cancer. *Biochem. Biophys. Res. Comm.* 222:292–297.
- Chomczynski, P., and N. Sacchi. 1987. Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal. Biochem.* 162:156–159.
- Clemson, C.M., J.A. McNeil, H.F. Willard, and J.B. Lawrence. 1996. *XIST* RNA paints the inactive X chromosome at interphase: evidence for a novel RNA involved in nuclear/chromosome structure. *J. Cell Biol.* 132:259–275.
- Disteche, C.M. 1995. Escape from X inactivation in human and mouse. *Trends Genet.* 11(1):17–22.
- Duncan, A.M., A. Macdonald, C.J. Brown, D. Wolff, H.F. Willard, and B. Sutton. 1993. Characterization of a small supernumerary ring X chromosome by fluorescence in situ hybridization. *Am. J. Med. Genet.* 47:1153–1156.
- Fey, E.G., G. Krochmalnic, and S. Penman. 1986. The non-chromatin substructures of the nucleus: the ribonucleoprotein RNP-containing and RNP-depleted matrices analyzed by sequential fractionation and resinless section electron microscopy. *J. Cell Biol.* 102:1654–1665.
- Gartler, S.M., and M.A. Goldman. 1994. Reactivation of inactive X-linked genes. *Dev. Genet.* 15:504–514.
- Gregory, P., C. Greene, E. Shapira, and E. Wang. 1985. Alterations in the time of X chromosome replication induced by 5-azacytidine in a patient with 48,XXXYY/47,XXY. *Cytogenet. Cell Genet.* 39:234–236.
- Haaf, T., G. Ott, and M. Schmid. 1988. Inhibition of condensation in the late-replicating X chromosome induced by 5-azadeoxycytidine in human lymphocyte cultures. *Hum. Genet.* 79:18–23.
- Hansen, R.S., T. Canfield, A. Stanek, E. Keitges, and S. Gartler. 1998. Reactivation of *XIST* in normal fibroblasts and a somatic cell hybrid: Abnormal localization of *XIST* RNA in hybrid cells. *Proc. Natl. Acad. Sci. USA*. 95:5133–5138.
- Herzing, L.B., J.T. Romer, J.M. Horn, and A. Ashworth. 1997. *Xist* has properties of the X-chromosome inactivation centre. *Nature*. 386:272–275.
- Johnson, C.V., R.H. Singer, and J.B. Lawrence. 1991. Fluorescent detection of nuclear RNA and DNA: Implication for genome organization. *Methods Cell Biol.* 35:73–99.
- Kay, G.F., S.C. Barton, M.A. Surani, and S. Rastan. 1994. Imprinting and X chromosome counting mechanisms determine *Xist* expression in early mouse development. *Cell*. 77:639–650.
- Kay, G.F., G.D. Penny, D. Patel, A. Ashworth, N. Brockdorff, and S. Rastan. 1993. Expression of *Xist* during mouse development suggests a role in the initiation of X chromosome inactivation. *Cell*. 72:171–182.
- Keohane, A.M., L.P. O'Neill, N.D. Belyaev, J.S. Lavender, and B.M. Turner. 1996. X-inactivation and histone H4 acetylation in embryonic stem cells. *Dev. Biol.* 180:618–630.
- Lawrence, J.B., R.H. Singer, and L.M. Marselle. 1989. Highly localized tracks of specific transcripts within interphase nuclei visualized by in situ hybridization. *Cell*. 57:493–502.
- Lawrence, J.B., C.A. Villnave, and R.H. Singer. 1988. Sensitive high-resolution chromatin and chromosome mapping in situ: presence and orientation of two closely integrated copies of EBV in a lymphoma line. *Cell*. 52:51–61.
- Lee, J.T., and R. Jaenisch. 1997. Long-range cis effects of ectopic X-inactivation centres on a mouse autosome. *Nature*. 386:275–279.
- Lee, J.T., W.M. Strauss, J.A. Dausman, and R. Jaenisch. 1996. A 450 kb transgene displays properties of the mammalian X-inactivation center. *Cell*. 86:83–94.
- Lock, L.F., N. Takagi, and G.R. Martin. 1987. Methylation of the *Hprt* gene on the inactive X occurs after chromosome inactivation. *Cell*. 48:39–46.
- Lyon, M. 1961. Gene action in the X-chromosome of the mouse (*Mus musculus* L.). *Nature*. 190:372–373.
- Marahens, Y., B. Panning, J. Dausman, W. Strauss, and R. Jaenisch. 1997. *Xist* deficient mice are defective in dosage compensation but not spermatogenesis. *Genes Dev.* 11:156–166.
- Panning, B., J. Dausman, and R. Jaenisch. 1997. X chromosome inactivation is mediated by *Xist* RNA stabilization. *Cell*. 90:907–916.
- Panning, B., and R. Jaenisch. 1996. DNA hypomethylation can activate *Xist* expression and silence X-linked genes. *Genes Dev.* 10:1991–2002.
- Penny, G.D., G.F. Kay, S.A. Sheardown, S. Rastan, and N. Brockdorff. 1996. Requirement for *XIST* in X chromosome inactivation. *Nature*. 379:131–137.
- Pfeifer, G.P., S.D. Steigerwald, R.S. Hansen, S.M. Gartler, and A.D. Riggs. 1990. Polymerase chain reaction-aided genomic sequencing of an X chromosome-linked CpG island: methylation patterns suggest clonal inheritance, CpG site autonomy, and an explanation of activity state stability. *Proc. Natl. Acad. Sci. USA*. 87:8252–8256.
- Rack, K.A., J. Chelly, R.J. Gibbons, S. Rider, D. Benjamin, R.G. Lafreniere, D. Osciur, R.W. Hendriks, I.W. Craig, H.F. Willard, et al. 1994. Absence of the *XIST* gene from late-replicating isodicentric X chromosomes in leukaemia. *Hum. Mol. Genet.* 3:1053–1059.
- Razin, A., and R. Shemer. 1995. DNA methylation patterns in the mammalian

- genome. *Hum. Mol. Genet.* 4:1751–1755.
- Sheardown, S.A., S.M. Duthie, C.M. Johnston, A.E.T. Newall, E.J. Formstone, R.M. Arkell, T.B. Nesterova, G.C. Alghisi, S. Rastan, and N. Brockdorff. 1997. Stabilization of Xist RNA mediates initiation of X chromosome inactivation. *Cell.* 91:99–107.
- Sympton, C.J., and T.E. Geoghan. 1990. Actin gene expression in murine erythroleukemia cells treated with cytochalasin D. *Exp. Cell Res.* 189:28–32.
- Tinker, A.V., and C.J. Brown. 1998. Induction of XIST expression from the human active X chromosome in mouse/human somatic cell hybrids by disruption of methylation. *Nucl. Acids Res.* 26:2935–2940.
- Toth, M., U. Lichtenberg, and W. Doerfler. 1989. Genomic sequencing reveals a 5-methylcytosine-free domain in active promoters and the spreading of preimposed methylation patterns. *Proc. Natl. Acad. Sci. USA.* 86:3728–3732.
- Turker, M.S., and T.H. Bestor. 1997. Formation of methylation patterns in the mammalian genome. *Mutat. Res.* 386:119–130.
- Willard, H.F., C.J. Brown, L. Carrel, B. Hendrich, and A.P. Miller. 1993. Epigenetic and chromosomal control of gene expression: molecular and genetic analysis of X chromosome inactivation. *Cold Spring Harbor Symp. Quant. Biol.* 58:315–322.
- Xing, Y., C.V. Johnson, P.R. Dobner, and J.B. Lawrence. 1993. Higher level organization of individual gene transcription and RNA splicing. *Science.* 259:1326–1330.
- Xing, Y., C.V. Johnson, P.T. Moen, J.A. McNeil, and J.B. Lawrence. 1995. Non-random gene organization: Structural arrangements of specific pre-mRNA transcription and splicing with SC-35 domains. *J. Cell Biol.* 131:1635–1647.
- Xing, Y., and J.B. Lawrence. 1993. Nuclear RNA tracks: structural basis for transcription and splicing? *Trends Cell Biol.* 3:346–353.
- Yoshida, I., Y. Nishita, T.K. Mohandas, and N. Takagi. 1997. Reactivation of an inactive human X chromosome introduced into mouse embryonal carcinoma cells by microcell fusion with persistent expression of XIST. *Exp. Cell. Res.* 230:208–219.