

6-28-2013

Breastfeeding During Early Infancy is Associated with Higher Weight-Based World Health Organization Anthropometry

Daniel H. Libraty

University of Massachusetts Medical School

Rosario Z. Capeding

Research Institute for Tropical Medicine

Anamae Obcena

Research Institute for Tropical Medicine

See next page for additional authors

Follow this and additional works at: http://escholarship.umassmed.edu/faculty_pubs

 Part of the [Clinical Epidemiology Commons](#), [Epidemiology Commons](#), [Infectious Disease Commons](#), [Maternal and Child Health Commons](#), and the [Pediatrics Commons](#)

Repository Citation

Libraty, Daniel H.; Capeding, Rosario Z.; Obcena, Anamae; Brion, Job D.; and Tallo, Veronica, "Breastfeeding During Early Infancy is Associated with Higher Weight-Based World Health Organization Anthropometry" (2013). *University of Massachusetts Medical School Faculty Publications*. 391.

http://escholarship.umassmed.edu/faculty_pubs/391

Breastfeeding During Early Infancy is Associated with Higher Weight-Based World Health Organization Anthropometry

Authors

Daniel H. Libraty, Rosario Z. Capeding, Anamae Obcena, Job D. Brion, and Veronica Tallo

Keywords

Breastfeeding, infant, anthropometry, weight, nutrition, bottle-feeding

Comments

Copyright Libraty et al.; Licensee Bentham Open. This is an open access article licensed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/bync/3.0/>) which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided the work is properly cited.

Rights and Permissions

Citation: Libraty DH, Capeding RZ, Obcena A, Brion JD, Tallo V. Breastfeeding During Early Infancy is Associated with Higher Weight-Based World Health Organization Anthropometry. *Open Pediatr Med Journal*. 2013 Jun 28;7. doi:10.2174/1874309920130610001. [Link to article on publisher's site](#)

Breastfeeding During Early Infancy is Associated with Higher Weight-Based World Health Organization Anthropometry

Daniel H. Libraty^{*1}, Rosario Z. Capeding^{2,3}, AnaMae Obcena², Job D. Brion⁴ and Veronica Tallo⁵

¹Division of Infectious Disease and Immunology, Department of Medicine, University of Massachusetts Medical School, Worcester, MA, USA

²Department of Medicine, Research Institute for Tropical Medicine, Manila, Philippines

³Department of Microbiology, Research Institute for Tropical Medicine, Manila, Philippines

⁴San Pablo City Health Office, San Pablo, Philippines

⁵Department of Epidemiology, Research Institute for Tropical Medicine, Manila, Philippines

Abstract: The World Health Organization (WHO) Expert Committee on Physical Status: The Use and Interpretation of Anthropometry established reference anthropometric standards for the growth of healthy infants and children. As part of a prospective clinical study of dengue virus infections in infants, we measured the length and weight of healthy infants in San Pablo, Laguna, Philippines at two scheduled study visits. We examined the correlation between breastfeeding and WHO anthropometric z scores during early infancy in San Pablo, Laguna, Philippines. We found that breastfeeding status and the frequency of breastfeeding during early infancy positively correlated with weight-based WHO anthropometric z scores.

Keywords: Breastfeeding, infant, anthropometry, weight, nutrition, bottle-feeding.

INTRODUCTION

The World Health Organization (WHO) Expert Committee on Physical Status: The Use and Interpretation of Anthropometry established reference anthropometric standards for the growth of healthy infants and children [1]. These standards were designed to reflect the growth of healthy infants and children worldwide, in both developed and developing countries. We examined the correlation between breastfeeding and WHO anthropometric z scores during early infancy in San Pablo, Laguna, Philippines. We found that breastfeeding status and the frequency of breastfeeding during early infancy positively correlated with weight-based WHO anthropometric z scores.

METHODS

Infant Clinical Study. The infant clinical study was approved by the institutional review boards of the Research Institute for Tropical Medicine, Philippines, and the University of Massachusetts Medical School. Mothers and their healthy infants were recruited and enrolled after providing written informed consent. Study enrollment began in October 2006 in San Pablo, Philippines. Healthy infants and their mothers were enrolled when the infant was between 6-18 weeks old. Additional details about the study protocol have been previously reported [2].

Anthropometry. At scheduled study visits, infant weight was measured to the nearest tenth of a kilogram. Infant length was measured to the nearest centimeter. WHO length-for-age, body mass index (BMI)-for-age, weight-for-age, and weight-for-length z scores for study infants were determined using the SPSS macro provided by WHO [3]. Infants with missing values or biologically implausible anthropometric z scores were excluded from analyses. Biologically implausible z scores were length-for-age z score <-6 or >6, BMI-for-age z score <-6 or >6, weight-for-age z score <-6 or >5, or weight-for-length z score <-6 or >6 [3].

Statistical Analysis. The SPSS software package (version 20.0) was used for statistical analyses. Ages are shown as median [95% confidence interval]. Spearman rank-order correlation coefficients and 95% confidence intervals were calculated. P<0.05 was considered significant.

RESULTS AND DISCUSSION

Clinical study. As part of a prospective clinical study of dengue virus infections in infants, we measured the length and weight of healthy infants in San Pablo, Laguna, Philippines at two scheduled study visits. We then calculated WHO anthropometric indices that use these measurements-length-for-age, BMI-for-age, weight-for-age, and weight-for-length z scores.

Breastfeeding is associated with higher weight-based WHO anthropometric z scores compared to bottle-feeding early in infancy. Breastfeeding (exclusive or supplemental) at the first study visit (ages 2.1 [2.1-2.2] mos) was associated with higher BMI-for-age, weight-for-age, and weight-for-

*Address correspondence to this author at the Rm S6-862, Division of ID, UMMS, 55 Lake Ave N, Worcester, MA 01655, USA; Tel: 508-856-4182; Fax: 508-856-4890; E-mail: daniel.libraty@umassmed.edu

Table 1. Correlations Between Breastfeeding and WHO Anthropometric z Scores During Early Infancy (Ages 2.1 [2.1-2.2] Mos, Median [95% CI])

	BMI-for-Age z Score	Weight-for-Age z Score	weight-for-Length z Score	length-for-Age z Score
Spearman <i>r</i> values [95% confidence interval] for breastfeeding (exclusive or supplemental) vs bottle-feeding (all infants) ^a , p-value	0.13 [0.11-0.15], p<0.001 (n=9,072)	0.12 [0.10-0.14], p<0.001 (n=9,079)	0.11 [0.09-0.13], p<0.001 (n=9,072)	0.02 [0.00-0.04], p=0.14 (n=9,075)
Spearman <i>r</i> values [95% confidence interval] for breastfeeding frequencies (among breastfed infants) ^b , p-value	0.10 [0.08-0.12], p<0.001 (n=7,355)	0.07 [0.05-0.09], p<0.001 (n=7,359)	0.09 [0.07-0.11], p<0.001 (n=7,353)	-0.02 [-0.04-0.002], p=0.04 (n=7,355)

^aBottle-feeding (exclusive) was coded as 0, breastfeeding (exclusive or supplemental) was coded as 1.

^bDaily frequencies of breastfeeding (among breastfed infants) were coded as follows:
 ≤ 2 times per 24 h period = 0; ≥ 3 and ≤ 6 times per 24 h period = 1; > 6 times per 24 h period = 2.

length WHO z scores compared to bottle-feeding (Table 1). There was no significant difference in length-for-age z scores between breastfed and bottlefed infants at this time point. Previous studies have reported little difference in growth patterns between breastfed and bottle-fed infants through early infancy [4]. Although we saw a significant positive correlation between breastfeeding and weight-based anthropometric z scores, the degree of correlation was small. There were no differences in all anthropometric z scores between breastfed and bottle-fed infants at the later second study visit (ages 4.4 [4.3-4.4] mos, data not shown). There was a lower percentage of breastfed infants at the second study visit compared to the first (75% breastfed infants at second study visit vs 81% breastfed infants at first study visit). This likely contributed to the absence of a significant association, as the degree of correlation between breastfeeding and weight-based anthropometric z scores was small at the first study visit.

A higher breastfeeding frequency during early infancy is associated with higher weight-based WHO anthropometric z scores. Among breastfed infants at the first study visit, BMI-for-age, weight-for-age, and weight-for-length WHO z scores positively correlated with breastfeeding frequencies (Table 1). This positive association was not seen with length-for-age z scores at the first study visit or with all the anthropometric measures at the second study visit during infancy (ages 4.3 [4.3-4.3] mos, data not shown). The positive association seen between breastfeeding frequencies

and weight-based anthropometric z scores during early infancy cannot establish causation. However, the data likely provide further support for the benefits of frequent breastfeeding early in infancy [5, 6].

CONFLICT OF INTEREST

The authors confirm that this article content has no conflict of interest.

ACKNOWLEDGEMENTS

This work was supported by National Institutes of Health grant NIH/NIAID R01 AI091820.

REFERENCES

- [1] WHO Child Growth Standards based on length/height, weight and age. Acta Paediatr Suppl. 2006; 450: 76-85.
- [2] Libraty DH, Acosta LP, Tallo V, *et al.* A prospective nested case-control study of Dengue in infants: rethinking and refining the antibody-dependent enhancement dengue hemorrhagic fever model. PLoS Med. 2009; 6(10): e1000171.
- [3] WHO Anthro for personal computers, version 3.2.2.: Software for assessing growth and development of the world's children. 2011. Available at: <http://www.who.int/childgrowth/software/en/>
- [4] Nommsen-Rivers LA, Dewey KG. Growth of breastfed infants. Breastfeed Med 2009; 4(1): 45-9.
- [5] Vis HL, Ruchababisha M, Hennart P. Breast-feeding and the growth and development of the infant. Int J Gynaecol Obstet 1987; 25: 239-47.
- [6] Fulhan J, Collier S, Duggan C. Update on pediatric nutrition: breastfeeding, infant nutrition, and growth. Curr Opin Pediatr 2003; 15(3): 323-32.

Received: March 30, 2013

Revised: May 17, 2013

Accepted: June 3, 2013

© Libraty *et al.*; Licensee Bentham Open.

This is an open access article licensed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted, non-commercial use, distribution and reproduction in any medium, provided the work is properly cited.