

May 20th, 12:30 PM

A Pilot Randomized Controlled Trial of a Videoconferencing Smoking Cessation Intervention for Korean American Women: Preliminary Findings

Sun S. Kim

University of Massachusetts Boston, sun.kim@umb.edu

Somporn Sitthisongkram

University of Massachusetts Boston

Kunsook Bernstein

CUNY Hunter College

See next page for additional authors

Follow this and additional works at: http://escholarship.umassmed.edu/cts_retreat

 Part of the [Behavior and Behavior Mechanisms Commons](#), [Substance Abuse and Addiction Commons](#), [Telemedicine Commons](#), and the [Women's Health Commons](#)

Sun S. Kim, Somporn Sitthisongkram, Kunsook Bernstein, Hua (Julia) Fang, Won S. Choi, and Douglas M. Ziedonis, "A Pilot Randomized Controlled Trial of a Videoconferencing Smoking Cessation Intervention for Korean American Women: Preliminary Findings" (May 20, 2016). *UMass Center for Clinical and Translational Science Research Retreat*. Paper 47. http://escholarship.umassmed.edu/cts_retreat/2016/posters/47

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.

Presenter Information

Sun S. Kim, Somporn Sitthisongkram, Kunsook Bernstein, Hua (Julia) Fang, Won S. Choi, and Douglas M. Ziedonis

Keywords

Korean women, smoking cessation, quit smoking, videoconferencing

Creative Commons License

This work is licensed under a [Creative Commons Attribution-Noncommercial-Share Alike 3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/).

A Pilot Randomized Controlled Trial of a Videoconferencing Smoking Cessation Intervention for Korean American Women: Preliminary Findings

Sun S Kim, PhD, APRN, University of Massachusetts Boston-College of Nursing and Sciences
Somporn Sitthisongkram, MS, University of Massachusetts Boston-College of Nursing and Sciences

Kunsook Bernstein, PhD, APN, Hunter College City of New York-School of Nursing
Hua Fang, PhD, University of Massachusetts Medical School-Department of Quantitative Sciences

Won S. Choi, PhD, MPH, University of Kansas Medical Center-Department of Preventive Medicine & Public Health

Douglas Ziedonis, MD, MPH, University of Massachusetts Medical School-Department of Psychiatry

ABSTRACT

Introduction: Korean American women prefer online or telephone smoking cessation interventions that can be remotely accessed from home. However, these interventions have been found ineffective for the group.

Methods: This study is a pilot clinical trial testing the feasibility and acceptability of a videoconferencing smoking cessation intervention for Korean American women and compared its preliminary efficacy with a telephone-based smoking cessation intervention. Korean women in the United States were recruited nationwide and randomly assigned at a ratio of 1:1 to either a video arm or a telephone arm. Participants in both arms received eight 30-minute weekly individualized counseling sessions of a culturally adapted smoking cessation intervention and nicotine patches for 8 weeks. They were followed up at post-quit 1, 2, and 3 months.

Results: A total of 168 Korean Americans were assessed for eligibility, 77 were determined to be eligible and 49 participated in the study. The videoconferencing intervention was acceptable and feasible for women under 50 years, whereas it was not for older women. The videoconferencing intervention produced abstinence rates of 67% at post-quit 1 month and 42% at post-quit 3 months based on self-report. The rate at post-quit 3 months dropped to 33% when those women whose abstinence could not be validated with salivary cotinine tests were treated as smoking. Abstinence rates in the telephone arm did not differ from those in the video arm.

Conclusion: Findings suggest that videoconferencing smoking cessation intervention may be feasible and acceptable for Korean American women under 50 years. However, for older Korean American women, the intervention may not be feasible and telephone-based intervention seems to be just as effective if smoking cessation intervention components are adapted at a deep structural level of Korean culture by integrating its core cultural values and addressing psychosocial, social and environmental forces affecting the behavior.

Contact:

Sun S. Kim

Sun.Kim@umb.edu

UMass Boston

