

May 20th, 12:30 PM

Angina Characteristics as Predictors of Trajectories of Quality of Life Following Acute Coronary Syndrome in the Transitions, Risks and Actions in Coronary Events-Center for Outcomes Research and Education cohort (TRACE-CORE)

Lisa Nobel

University of Massachusetts Medical School, Lisa.Nobel@umassmed.edu

Jennifer Tjia


University of Massachusetts Medical School, jennifer.tjia@umassmed.edu

Jane S. Saczynski

University of Massachusetts Medical School, jane.saczynski@umassmed.edu

See next page for additional authors

Follow this and additional works at: http://escholarship.umassmed.edu/cts_retreat

 Part of the [Cardiology Commons](#), [Cardiovascular Diseases Commons](#), [Clinical Epidemiology Commons](#), [Epidemiology Commons](#), and the [Psychiatric and Mental Health Commons](#)

Lisa Nobel, Jennifer Tjia, Jane S. Saczynski, Molly E. Waring, Milena D. Anatchkova, Arlene Ash, Catarina I. Kiefe, and Jeroan Allison, "Angina Characteristics as Predictors of Trajectories of Quality of Life Following Acute Coronary Syndrome in the Transitions, Risks and Actions in Coronary Events-Center for Outcomes Research and Education cohort (TRACE-CORE)" (May 20, 2016). *UMass Center for Clinical and Translational Science Research Retreat*. Paper 62.
http://escholarship.umassmed.edu/cts_retreat/2016/posters/62

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.

Presenter Information

Lisa Nobel, Jennifer Tjia, Jane S. Saczynski, Molly E. Waring, Milena D. Anatchkova, Arlene Ash, Catarina I. Kiefe, and Jeroan Allison

Keywords

acute coronary syndrome, quality of life, angina

Creative Commons License

This work is licensed under a [Creative Commons Attribution-Noncommercial-Share Alike 3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/).

Angina Characteristics as Predictors of Trajectories of Quality of Life Following Acute Coronary Syndrome in the Transitions, Risks and Actions in Coronary Events-Center for Outcomes Research and Education cohort (TRACE-CORE)

Lisa Nobel, University of Massachusetts Worcester, Worcester, MA, United States; Jennifer Tjia, University of Massachusetts Worcester, Worcester, MA, United States; Jane Saczynski, University of Massachusetts Worcester, Worcester, MA, United States; Molly E. Waring, University of Massachusetts Worcester, Worcester, MA, United States; Milena D. Anatchkova, PhD, University of Massachusetts Medical School, Evidera, Lexington, MA, United States; Arlene Ash, University of Massachusetts Worcester, Worcester, MA, United States; Catarina Kiefe, University of Massachusetts Worcester, Worcester, MA, United States; Jeroan Allison, University of Massachusetts Worcester, Worcester, MA, United States

BACKGROUND: To describe longitudinal trajectories of health-related quality of life (HRQoL) after hospitalization with an acute coronary syndrome (ACS), their associations with baseline angina characteristics, and associations with anxiety, depression, and cognitive impairment.

METHODS: TRACE-CORE participants (N=1,613) completed the SF-36 during hospitalization for ACS and 1, 3, & 6 months post-discharge. Latent growth curves identified trajectories of physical and mental components of HRQoL (MCS and PCS) and sequential multiple logistic regression estimated associations between trajectories and angina characteristics.

RESULTS: Participants (N=1613) had mean age 63.3 (SD 11.4) years, 33.0% female, and 78.2% non-Hispanic white. We identified 2 MCS trajectories: AVERAGE and IMPAIRED HRQoL. The majority of participants (81.0%) had AVERAGE MCS at baseline (mean MCS 53.6) and slight improvement in scores over time. A minority (19.0%) had IMPAIRED HRQoL at baseline (mean MCS 36.7) and slight improvement in scores over time. We identified 2 similar PCS trajectories with similar patterns of scores over time: AVERAGE (71.1%) and IMPAIRED (28.9%) HRQoL at baseline. Adjusting for demographics & comorbidities, patients with less severe baseline angina were more likely to have AVERAGE MCS (odds ratio [OR]/10 unit change in severity 1.1) and PCS (OR 1.1) trajectories, and similarly for less frequent angina (MCS OR 1.2; PCS OR 1.3). The associations of MCS trajectory with severity and frequency lost significance after adjusting for psychosocial factors, whereas the PCS associations remained significant [All $p < 0.05$ unless noted].

CONCLUSIONS: About 1/3 of patients exhibited impaired 6-month HRQoL trajectories, which can be predicted by angina characteristics. Psychosocial factors may explain the prediction of mental, not physical, trajectories. Interventions to improve HRQoL after ACS should consider psychosocial factors and angina.

Lisa Nobel, MD/PhD Candidate
Department of Quantitative Health Sciences
University of Massachusetts Medical School
368 Plantation Street
The Albert Sherman Center
Worcester, MA 01605
E-mail: lisa.nobel@umassmed.edu