3-2010

Correlation of Measures of Psychotherapy Competency in Psychiatry Residents

Julia Matthews-Bellinger
University of Massachusetts Medical School, Julia.Matthews-Bellinger@umassmed.edu

Sheldon Benjamin
University of Massachusetts Medical School, sheldon.benjamin@umassmed.edu

Mohit Chauhan
Austin Medical Center

See next page for additional authors

Follow this and additional works at: http://escholarship.umassmed.edu/psych_pp

Part of the Psychiatry Commons

Correlation of Measures of Psychotherapy Competency in Psychiatry Residents

Authors

Julia Matthews-Bellinger, Sheldon Benjamin, Mohit Chauhan, and Mary L. Zanetti

Correlation of Measures of Psychotherapy Competency in Psychiatry Residents

Julia Matthews, PhD, MD; Sheldon Benjamin, MD; Mohit Chauhan, MD[a]; Mary Zanetti, EdD

UMass Medical School, UMass Memorial Healthcare, Worcester, MA   [a]Austin Medical Center, Mayo Health System, Austin, MN

INTRODUCTION: The ACGME Residency Review Committee in Psychiatry has stipulated that general psychiatry trainees develop "competency in applying supportive, psychodynamic, and cognitive-behavioral psychotherapies to both brief and long-term individual practice [1]." Residency programs are required to demonstrate and faculty are required to verify that trainees have attained the required competencies. Yet no generally accepted reliable and valid measure of psychotherapy competency has yet been developed.

Several measures are widely used to assess resident competence in psychotherapy [2,7].

- Evaluation by individual psychotherapy supervisors [3,4]
- Rating of resident's conduct of psychotherapy in live, recorded, or transcribed performance
- Training portfolios
- Performance on a multiple choice examination such as The Columbia Psychodynamic Psychotherapy Competence Test [5]
- Resident self-assessment such as the Counseling Self-Estimate Inventory (COSE), a self-report instrument designed to assess confidence and self-efficacy [6]

There have been few studies examining the reliability and validity of available competency assessment measures, and little empirical research examining the success of training programs in developing trainee competence in psychotherapy.

The current study had two goals:

1. To examine the changes in various performance measures as residents progress through training
2. To explore correlations among various measures of psychotherapy competence

METHODS: We conducted a retrospective analysis of all available data for residents in our program from July 2000 through July 2009 (63 residents), including:

1) Psychiatry Resident in Training Examination (PRITE) — Global Psychiatry score and Psychosocial Therapies subscore

2) Columbia Psychodynamic Psychotherapy Competency Test (CPPCT) - Scores (given as percentile rank among all examinees)

3) Counseling Self Estimate Inventory (COSE) — Scores on the COSE were collected for all PGY1 or PGY2 residents; however, COSE assessments were not routinely collected for PGY3 and PGY4 residents in the early years of this study. Our analysis used the first 34 of 37 items on the COSE due to inadvertent omission on some forms.

4) Supervisor ratings - Averaged global supervisor ratings of resident performance (5 point scale) in the following 9 areas: formulation skills; psychotherapeutic interventions; tolerance of uncertainty; transference, countertransference, boundaries; ability to utilize different conceptual models; humanistic qualities; practice-based learning (self evaluation, integration of feedback into practice); communication and interpersonal skills; working with difficult patients.

TABLE 1: Number of Residents with Available Data

	PGY-1	PGY-2	PGY-3	PGY-4
PRITE	49	47	42	33
CPPCT	49	48	40	21
COSE	49	48	26	17
Supervisor Rating	NA	39	35	24

Statistical analyses were performed with the Statistical Package for Social Sciences, SPSS. Analyses comparing performance in different training years were done using unpaired t tests (two tail) on pooled data for each training year. Correlations were done using paired data for individual residents to derive Pearson two tailed correlation coefficients. Due to missing data, the N for correlations was often less than the total N.

RESULTS: Figures 1-3 show changes in COSE, PRITE, and CPPCT by training year.

Fig 1: COSE Mean Item Score

*denotes p < 0.01 compared to PGY-1 and PGY-2

Fig 2: CPPCT Mean Percentile Rank

Fig 3: PRITE Standard Score

*denotes p < 0.01 compared to PGY-1

Changes in Performance Measures during Residency

- **PRITE:** Global Psychiatry scores changed significantly from PGY-2 - PGY-4. Psychosocial Therapy subscore did not change significantly from PGY-1 to PGY-2, but increased significantly in PGY-3 & PGY-4.

- **COSE (resident confidence):** COSE score unchanged (both total score & all individual COSE items) between PGY-1 and PGY-2. COSE increased dramatically by early PGY-3 after one year of classes and conducting supervised psychotherapy with 2 patients (24 of 34 COSE items showed significant change at p < 0.05 level or greater). COSE increased further in PGY-4 year (COSE N for PGY-4 too small for meaningful analysis). COSE item score range = 1-7.

- **CPPCT:** CPPCT showed trend toward improved performance (average percentile rank) each training year (changes did not reach significance).

- **Mean supervisor ratings** consistently high & did not change significantly between PGY 2 & PGY-4 (either group means or paired analysis).

Correlated Measures
- Individual resident's scores on COSE throughout training
- Individual resident's scores on PRITE in PGY-2 & PGY-3
- Individual resident's scores on PRITE & CPPCT in PGY-2
- Total PGY-1 & 2 COSE score (before psychotherapy experience) with PGY-4 CPPCT score

Non-Correlated Measures
- COSE score with overall PRITE psychiatry score or PRITE psychosocial subscore
- COSE score with CPPCT in same training year
- PGY-2 supervisor rating with PGY-2 COSE score
- PGY-3 supervisor rating with PRITE psychiatry or psychosocial score or CPPCT score

Negatively Correlated Measures
- PGY-2 & 3 COSE scores with PGY-3 psychotherapy supervisor rating

STUDY LIMITATIONS
- Small N (6 residents/yr max) and incomplete data
- Supervisor evaluations non-standardized & without measures of validity or reliability
- Performance measures collected at different times during training years
- Retrospective analysis of data not systematically collected for research

Table 2: Correlations Among Measures of Competence[§]

	Total COSE PGY 1	Total COSE PGY 2	Total COSE PGY 3	Rank PRITE Psych PGY 2	Rank PRITE Psysoc PGY 2	Rank PRITE Psych PGY 3	Rank PRITE Psysoc PGY 3	CPPCT Nat. Rank PGY 1	CPPCT Nat. Rank PGY 2	CPPCT Nat. Rank PGY 3	CPPCT Nat. Rank PGY 4	Sup. Eval. PGY 2	Sup. Eval. PGY 3
Total COSE PGY 1	1 N=49	.805** p<.000 N=36	.448* p=.042 N=21	-.134 p=.456 N=33	.031 p=.863 N=33	-.029 p=.881 N=29	.036 p=.855 N=29	.141 p=.400 N=38	.013 p=.942 N=34	.034 p=.868 N=27	.594** p=.032 N=13	.014 p=.944 N=29	-.291 p=.159 N=25
Total COSE PGY 2		1 N=48	.464* p=.026 N=23	-.019 p=.905 N=41	-.006 p=.972 N=41	.118 p=.492 N=36	.269 p=.113 N=36	.111 p=.540 N=20	.099 p=.534 N=34	.207 p=.241 N=26	.616** p=.008 N=17	-.149 p=.399 N=34	-.370* p=.037 N=32
Total COSE PGY 3			1 N=26	-.319 p=.121 N=25	.222 p=.286 N=25	-.154 p=.483 N=23	-.248 p=.253 N=23	-.085 p=.722 N=20	-.146 p=.478 N=26	.202 p=.379 N=21	.218 p=.546 N=10	.002 p=.994 N=23	-.485* p=.030 N=20
Rank PRITE Psychiatry PGY 2				1 N=47	.403** p=.005 N=47	.648** p=.000 N=41	.495** p=.001 N=41	.454* p=.012 N=30	.359* p=.030 N=47	-.027 p=.868 N=39	-.157 p=.546 N=21	.448** p=.007 N=39	.063 p=.715 N=36
Rank PRITE Psychosocial PGY 2					1 N=47	.404** p=.009 N=41	.367* p=.018 N=41	.397* p=.030 N=30	.031 p=.834 N=47	-.083 p=.615 N=39	.120 p=.604 N=21	.269 p=.098 N=39	-.147 p=.389 N=36
Rank PRITE Psychiatry PGY 3						1 N=42	.595** p=.000 N=42	.089 p=.673 N=25	.166 p=.294 N=42	-.047 p=.775 N=25	.159 p=.491 N=33	-.004 p=.984 N=36	-.010 p=.956 N=33
Rank PRITE Psychosocial PGY 3							1 N=42	.049 p=.814 N=25	.141 p=.374 N=42	.136 p=.403 N=25	.062 p=.790 N=12	-.120 p=.508 N=36	-.240 p=.158 N=33
CPPCT Rank PGY 1								1 N=39	.536** p=.002 N=30	.353 p=.099 N=23	.278 p=.382 N=12	.508** p=.007 N=27	.114 p=.615 N=22
CPPCT Rank PGY 2									1 N=48	.219 p=.175 N=40	-.023 p=.922 N=21	.206 p=.209 N=39	-.022 p=.897 N=36
CPPCT Rank PGY 3										1 N=40	.392 p=.344 N=8	.344 p=.058 N=30	.163 p=.350 N=31
CPPCT Rank PGY 4											1 N=21	.390 p=.122 N=17	.260 p=.297 N=18
Supervisor Evaluation PGY 2												1 N=39	.192 p=.329 N=28
Supervisor Evaluation PGY 3													1 N=36

[§]Pearson correlation coefficients, two-tailed, paired analysis

DISCUSSION
- Resident confidence increases with first experiences conducting supervised psychodynamic psychotherapy, paralleled by significant improvement in PRITE psychosocial subscores.
- The early COSE-late CPPCT score correlation may suggest that greater psychotherapy interest or experience on entry predicts greater learning.
- The negative correlation between supervisor rating and COSE scores suggests a possible supervisory reaction to overconfidence and/or inflation of ratings to boost confidence.
- Uniformly above average supervisor scores that do not change year to year may reflect reluctance to give lower ratings, rating by PGY-expectation rather than competence, or lack of dependability of supervisor ratings.
- The relative lack of correlations may mean measures assess different aspects of performance, lack validity, or merely reflect the small N's in our study.

CONCLUSIONS:
Resident self-assessment does not appear to reflect competence as assessed by supervisors, or knowledge as assessed by objective tests.

Global supervisor evaluations do not appear to reliably measure competence.

Standardized supervisor assessment instruments are needed.

Further work is needed to develop reliable, valid measures of competency.

REFERENCES:

1. ACGME Program Requirements for Graduate Medical Education in Psychiatry (Effective: July 1, 2007)

2. Psychotherapy Competency Assessment Tools: User-Friendly Summary of Available Assessment Tools and Annotations Vers. 2-14-05. http://www.aadprt.org/training/assesment_tools/

3. Mohl P, Lomax J, Tasman A, et al, (1990) Psychotherapy training for the psychiatrist of the future. Am J Psychiatry 145:7-13

4. Holloway E, Neufeldt S, (1995) Supervision: its contribution to treatment efficacy. J Consult Clin Psychol 63:207-213

5. Mullen, LS, Rieder RO, Glick, RA, Luber, B, Rosen, PJ, (2004)Testing Psychodynamic Psychotherapy Skills among Psychiatric Residents: the Psychodynamic Psychotherapy Competency Test. Am J Psychiatry 161:1658-1664

6. Larson, LM, Suzuki, LA, Gillespie, KN, Potenza, MT, Bechtel, MA, and Toulouse, AL, (1992) Development and validation of the Counseling Self-Estimate Inventory. J Counseling Psychology 39:105-120

7. Giordano FL, Briones DF. (2003) Assessing Residents' Competence in Psychotherapy Acad Psychiatry 27:145–147