

6-22-2011

Temporal trends in serum concentrations of polychlorinated dioxins, furans, and PCBs among adult women living in Chapaevsk, Russia: a longitudinal study from 2000 to 2009


Olivier Humblet
Harvard School of Public Health

Oleg Sergeyevev
Samara State Medical University

Larisa Altshul
Environmental Health and Engineering, Inc.

See next page for additional authors

Follow this and additional works at: <http://escholarship.umassmed.edu/oapubs>

 Part of the [Cell Biology Commons](#), [Endocrinology, Diabetes, and Metabolism Commons](#), [Environmental Public Health Commons](#), and the [Pediatrics Commons](#)

Repository Citation

Humblet, Olivier; Sergeyevev, Oleg; Altshul, Larisa; Korrnick, Susan A.; Williams, Paige L.; Emond, Claude; Birnbaum, Linda S.; Burns, Jane S.; Lee, Mary M.; Revich, Boris; Shelepchikov, Andrey; Feshin, Denis; and Hauser, Russ, "Temporal trends in serum concentrations of polychlorinated dioxins, furans, and PCBs among adult women living in Chapaevsk, Russia: a longitudinal study from 2000 to 2009" (2011). *Open Access Articles*. 2258.
<http://escholarship.umassmed.edu/oapubs/2258>

Temporal trends in serum concentrations of polychlorinated dioxins, furans, and PCBs among adult women living in Chapaevsk, Russia: a longitudinal study from 2000 to 2009

Authors

Olivier Humblet, Oleg Sergeev, Larisa Altshul, Susan A. Korrick, Paige L. Williams, Claude Emond, Linda S. Birnbaum, Jane S. Burns, Mary M. Lee, Boris Revich, Andrey Shelepchikov, Denis Feshin, and Russ Hauser


Comments

© 2011 Humblet et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Rights and Permissions

Citation: Environ Health. 2011 Jun 22;10:62. [Link to article on publisher's site](#)



RESEARCH

Open Access

Temporal trends in serum concentrations of polychlorinated dioxins, furans, and PCBs among adult women living in Chapaevsk, Russia: a longitudinal study from 2000 to 2009

Olivier Humblet¹, Oleg Sergeev^{2,3}, Larisa Altshul^{4,5}, Susan A Korrick^{1,6}, Paige L Williams⁷, Claude Emond^{8,9}, Linda S Birnbaum¹⁰, Jane S Burns¹, Mary M Lee¹¹, Boris Revich¹², Andrey Shelepchikov¹³, Denis Feshin¹³ and Russ Hauser^{1*}

Abstract

Background: The present study assessed the temporal trend in serum concentrations of polychlorinated dibenzo-*p*-dioxins, dibenzofurans, and biphenyls (PCBs) among residents of a Russian town where levels of these chemicals are elevated due to prior industrial activity.

Methods: Two serum samples were collected from eight adult women (in 2000 and 2009), and analyzed with gas chromatography-high-resolution mass spectrometry.

Results: The average total toxic equivalency (TEQ) decreased by 30% (from 36 to 25 pg/g lipid), and the average sum of PCB congeners decreased by 19% (from 291 to 211 ng/g lipid). Total TEQs decreased for seven of the eight women, and the sum of PCBs decreased for six of eight women. During this nine year period, larger decreases in serum TEQs and PCBs were found in women with greater increases in body mass index.

Conclusions: This study provides suggestive evidence that average serum concentrations of dioxins, furans, and PCBs are decreasing over time among residents of this town.

Introduction

Polychlorinated dibenzo-*p*-dioxins (PCDDs), dibenzofurans (PCDFs), and biphenyls (PCBs) are classes of toxic organochlorine compounds which are ubiquitous in the environment [1,2]. They are lipophilic and persistent, and their elimination from the body can take years or even decades [3]. Current human exposure is mostly through diet [4], although some populations may have continuing environmental exposures due to past or current industrial activity [5-8].

Numerous studies in the general population have reported that serum concentrations of these chemicals correlate positively with age [9,10], but also that average

population levels are declining over time [11]. The objective of this study was to characterize temporal trends in serum concentrations of PCDD/PCDF/PCBs among reproductive age women living in Chapaevsk, Russia. Elevated levels of these compounds have been detected in serum samples from Chapaevsk residents [12,13], as well as in environmental samples and food [14]. For over 50 years this town was the site of chlorinated chemical production at the Middle Volga Chemical Plant [7], also known as SVZH. Plant activity started decreasing in 1991 [15], and ceased in 2003. The plant produced chemical blister agents (mustard gas and lewisite) prior to 1949, hexachlorocyclohexane (lindane) and its derivatives from 1967 to 1987, then crop protection chemicals (liquid chlorine, acids, methyl chloroform, and vinyl chloride) until its closure [7]. Dioxins were generated as unwanted by-products of chemical manufacturing [7], whereas PCBs were typically used in electrical capacitors and

* Correspondence: rhauser@hohp.harvard.edu

¹Environmental and Occupational Medicine and Epidemiology Program, Department of Environmental Health, Harvard School of Public Health, Boston, Massachusetts, USA

Full list of author information is available at the end of the article

transformers, although their specific use at SVZH is unknown. Environmental release of these chemicals most likely occurred during incineration and disposal of hazardous waste from the plant. We collected serum samples nine years apart (in 2000 and 2009) from eight non-occupationally exposed women living in Chapaevsk, and in 2009 all samples were sent for analysis of PCDD/PCDF/PCBs.

Methods

Study population

Ten healthy women were recruited in 2000 for a pilot feasibility study on early pregnancy loss, at which time a serum sample was collected and archived, and a health and lifestyle questionnaire was completed. Newly married women in Chapaevsk were invited to join, and 10 enrolled in the pilot project. To be included in the original study women needed to be married, age 20-34 years, and trying to conceive. To investigate temporal trends in PCDD/PCDF/PCB exposure, participants were re-contacted in 2009, and a second serum sample and lifestyle questionnaire was collected from eight of the ten women (of the two others, one had left Chapaevsk, and one declined to participate).

The research protocol was approved by the Human Studies Institutional Review Boards of the Chapaevsk Medical Association, UMass Medical School, and Harvard School of Public Health. All participants gave their consent before participating in each of the two assessments.

Measurement of dioxin, furan, and PCB serum concentrations

Samples were stored at -20 degrees C until analysis. All 16 serum samples (10-15 ml each) were analyzed in 2009 using isotope dilution gas chromatography-high-resolution mass spectrometry (GC-HRMS-ID) at the Laboratory of Analytical Toxicology of the Severtsov Institute of Ecology and Evolution in Moscow, Russia. The list of dioxin-like congeners (called dioxins in this manuscript) included 7 PCDDs, 10 PCDFs, 4 co-planar PCBs, and 6 mono-ortho PCBs. The list of non-dioxin like PCBs (NDL-PCBs) included congeners 18, 28, 44, 49, 52, 66, 74, 87, 99, 101, 110, 128, 138, 146, 149, 151, 153, 170, 172, 177, 178, 180, 183, 187, 190, 194, 195, 196, 199, 203, 206, 209. Congeners are identified according to the International Union for Pure and Applied Chemistry (IUPAC) nomenclature.

Prior to analysis of study samples the laboratory carefully cleaned and checked all reagents, solvents, and glassware, and analyzed a method blank to confirm that it was free from interference. Then each set of four samples was extracted and analyzed with one method blank, which was treated identically to the serum samples.

For the analysis of dioxins, furans and co-planar PCBs the serum samples were spiked with a mixture of $^{13}\text{C}_{12}$ -

labeled PCDDs/PCDFs and PCBs as internal standards. Methods for sample preparation, clean up and analysis follow US Environmental Protection Agency (EPA) methods 1613 [16] and 1668 [17]. Analysis was performed using a Thermo Finnigan MAT 95XP. The analytes were separated on a capillary column (SGE-BPX5, 30 meters, 0.25 mm, 0.25 microns) and quantified using selected-ion-monitoring high resolution (10,000 resolving power) mass spectrometry (HRGC-ID/HRMS). Quantification was by isotope dilution mass spectrometry using calibration standards containing ^{13}C labeled and unlabeled analytes. Mono-ortho (M-PCBs) and non-dioxin-like PCBs were separated on a capillary column (SGE-HT8, 30 meters, 0.25 mm, 0.25 microns).

LODs were calculated as three times chromatographic noise. The LODs for dioxins and furans ranged from 0.01 to 0.04 pg/g, and for C-PCBs from 0.1 to 0.6 pg/g serum. The levels of target analytes in 4 blanks analyzed with study samples were below the LODs. The mean (SD) recoveries were 90% (5.5) for TCDD and 84% (3.1) for HxCDD.

Serum total cholesterol and triglycerides were measured with an enzymatic colorimetric assay using the COBAS INTEGRA 400 plus system, and the serum total lipid content was calculated using the Phillips equation [18]. All PCDD/PCDF/PCB serum concentrations were lipid-adjusted. Individual congener concentrations below the limit of detection (LOD) were assigned a value equal to the LOD divided by the square root of 2 [19,20].

The serum concentrations of PCDD/PCDF/PCBs were summarized with two measures: the total sum of toxic equivalencies (i.e. total TEQ [21]) for PCDDs, PCDFs, coplanar PCBs (C-PCBs) and mono-ortho PCBs (M-PCBs); and the sum of all 36 assayed non-coplanar PCB concentrations (i.e. Σ PCBs).

Statistical methods

Pearson correlation coefficients were calculated for the log₁₀-transformed serum concentrations in 2000 and 2009; 95% confidence intervals (CIs) for the correlation coefficients were calculated using the method of Shen and Lu [22].

Results

Demographics

The women were aged 21 to 27 years at initial enrollment, and 30 to 35 years at the second follow-up in 2009. All had lived exclusively in Chapaevsk since 2000. One woman moved to Chapaevsk in 2000; the others had spent a maximum of 2 years living outside of Chapaevsk. None were employed at SVZH, the local chemical plant. BMIs ranged from 17 to 30 at enrollment in 2000, and from 20 to 31 in 2009. Seven of the eight women had a body mass index (BMI) <25 kg/m² in 2000; five did so in 2009. Six of eight women breastfed (either 1 or 2

children) between 2000 and 2009, with total durations ranging from 3 to 30 months (Table 1). One woman was currently breastfeeding; all others had last breastfed in 2007 or earlier. The two women who did not breastfeed were nulliparous.

Serum concentrations of total TEQs and ΣPCBs

The total TEQ and ΣPCB serum concentrations in both 2000 and 2009 are shown in Table 1 (results for all measured congeners are shown in Additional File 1, Table S1). The mean serum concentrations in 2000 and 2009 were, for total TEQs, 36 and 25 pg/g lipid, and for ΣPCBs, 291 and 211 ng/g lipid, respectively. Excluding the one woman who moved to Chapaevsk in 2000 (and therefore does not reflect the exposure levels of long-term residents), the respective 2000 and 2009 mean serum concentrations were, for total TEQs, 39 and 26 pg/g lipid, and for ΣPCBs, 312 and 210 ng/g lipid.

Trends in total TEQs and PCBs between 2000 and 2009 for each individual are shown graphically in Figure 1. Decreasing concentrations were found for seven of the eight women for TEQs, and for six of the eight women for ΣPCBs. The mean percent change was -30% for total TEQs, and -19% for ΣPCBs (Table 1). The mean decreases in the TEQs of the subtypes of dioxin-like compounds were -42% for PCDDs, -25% for PCDFs, -24% for M-PCBs, and -14% for C-PCBs. Similar mean decreases were seen when concentrations were expressed as the sum of measured congener values instead of in TEQs: -37% for PCDDs, -35% for PCDFs, -24% for M-PCBs, and -1% for C-PCBs. The slightly smaller decrease in C-PCB concentrations than for C-PCB TEQs (i.e. -1% vs. -14%) is because concentrations of PCB77 increased for 4 of the 8 women, but the small TEF of this congener (i.e. 0.0001) resulted in the increase having little effect on the mean C-PCB TEQs.

Several individual characteristics were associated with the magnitude of the changes in serum concentration between 2000 and 2009 (although the small sample size precluded formal statistical analyses on these predictors). First, a greater decrease in both total TEQs and ΣPCBs was observed among the four women with the largest increases in BMI during this nine year period, as compared to the other four women. TEQs decreased by an average of 39% among women whose BMIs increased by 10% or more, compared to 22% among women whose BMIs increased <10% or decreased. Similarly, PCBs decreased by an average of 42% among the former group of women, but increased by 3.8% among the latter. The latter group also includes the only woman's whose BMI was >30 kg/m² at either assessment. Second, the four women who breastfed longest (i.e. ≥12 months) had larger average decreases in TEQs than the women who breastfed less (-37% vs. -23% for TEQs, respectively), but there was not a greater decrease in PCBs by breastfeeding duration (-20% vs -19%, respectively).

Correlations between serum concentrations in 2000 and 2009

The Pearson correlations between the log₁₀-transformed serum concentrations in 2000 and 2009 were 0.3 (95% CI: -0.5; 0.8) for total TEQs (after excluding the highest value, which had inflated the correlation to 0.8), and 0.1 (95% CI: -0.6; 0.7) for ΣPCBs. However, the interpretability of these correlations was limited by the extremely wide confidence intervals.

Discussion

Summary

This study consisted of eight women from a town in Russia with high exposure to dioxins and PCBs from non-occupational sources. Each woman's serum

Table 1 Individual characteristics and serum concentrations of total TEQs and ΣPCBs, in 2000 and 2009, among 8 Russian women

	BMI percent change, 2000 to 2009	BMI in 2009	Months of breastfeeding, 2000 to 2009	Total TEQs (pg/g lipid)			Σ PCBs (ng/g lipid)		
				2000	2009	Percent change, 2000 to 2009	2000	2009	Percent change, 2000 to 2009
1	-3.0%	19.7	>12	16.7	14.7	-12%	139	215	+54%
2	0.4%	22.7	>12	39.5	19.7	-50%	327	185	-43%
3	3.7%	31.2	<6	29.2	31.2	+7%	238	290	+22%
4	8.4%	24.5	0	89.7	60.9	-32%	210	173	-18%
5	11.3%	19.7	>12	27.5	19.6	-29%	408	232	-43%
6	13.9%	27.0	6-12	28.3	19.9	-30%	388	266	-31%
7	22.9%	21.5	0	25.1	15.5	-38%	336	175	-48%
8	28.6%	25.6	>12	32.6	13.6	-58%	280	149	-47%
Mean (SD):	11% (11)	24 (4)	11 (10)	36 (23)	25 (16)	-30% (21)	291 (92)	211 (49)	-19% (38)

All percent changes are calculated using the 2000 values as the denominator.




Figure 1 Serum concentrations of serum total TEQs and Σ PCBs among eight Russian women, in 2000 and 2009. Figure 1A shows the serum total TEQ concentrations (pg/g lipid). Figure 1B shows the serum Σ PCB concentrations (ng/g lipid). Participants are numbered as in Table 1.

concentration of dioxins and PCBs was assessed in both 2000 and 2009. All of the women's serum concentrations of both total TEQs and Σ PCBs were greater than the 90th percentile values for US women 20 to 39 years of age in 2003-2004 (i.e. 16.2 pg/g lipid for total TEQs, and 157.8 ng/g lipid for Σ PCBs [10]) For almost all of the women, serum concentrations of total TEQs and Σ PCBs decreased substantially between 2000 and 2009

and, as expected, larger decreases were seen among women with greater BMI increases [23].

Although one must exercise caution in making inferences to the entire population of Chapaevsk (pop. 72,000) from the decreases observed among these eight women, concordant results were seen in our recent cross-sectional study among Chapaevsk women [13]. 446 adult women's serum concentrations of PCDD/PCDF/

PCBs were collected between 2003 and 2005, with average serum concentrations similar to those seen here. Levels were lower among women sampled later in the study period, adjusting for all other predictors. In multivariate linear regressions a one-year later blood draw was associated with 9.8% lower mean serum concentrations of Σ PCBs, and 10.5% lower mean total TEQs. While these age-adjusted regression results should not be interpreted as predictive of longitudinal changes in serum concentrations, they do suggest a trend similar to what was observed among the eight women.

Strengths and limitations

This study is one of the few to measure repeated serum concentrations of PCDD/PCDF/PCBs in a non-occupationally and non-accidentally exposed cohort. A particular strength of this study is that serum samples from both 2000 and 2009 were analyzed at the same time, precluding any potential bias from laboratory drift. Dioxins and PCBs are highly stable in stored serum [24], thus little degradation would be expected in the samples stored since 2000. However, the small sample size of eight women warrants that these findings be interpreted with caution. Furthermore, no longitudinal data on consumption of local meat or produce was available to assess the contribution of changing dietary patterns to the temporal changes in TEQ and PCB serum concentrations observed here.

Conclusions

This study reports the serum dioxin and PCB concentrations of adult women from a Russian town with high levels of non-occupational exposure to these compounds. All the women's levels were higher than typically seen in the US. Each woman was sampled twice: in 2000 and again in 2009. Among most of the women, serum concentrations of dioxins and PCBs decreased over this time period. Decreases were seen for all classes of dioxin-like compounds (i.e. PCDDs, PCDFs, C-PCBs and M-PCBs).

These results, while based on a small number of women, are consistent with those from our large cross-sectional study recently conducted among women in the same town. In the latter study, serum concentrations of dioxins and PCBs were lower, on average, among women sampled later in the study period, adjusting for all other predictors. While it is known that a general trend towards decreasing dioxin and PCB levels exists in the general population [25,26], it is significant that this also appears to be true among women living in an environment with a substantial, and relatively recent (SVZH ceased operations in 2003, although production started decreasing in 1991), non-occupational exposure source.

Additional material

Additional file 1: Table s1. The supplementary data table shows the distributions of all measured serum concentrations of polychlorinated dibenzo-*p*-dioxin, polychlorinated dibenzofuran, and polychlorinated biphenyl congeners, in 2000 and 2009, among the 8 study participants.

List of abbreviations

BMI: Body mass index kg/m²; C: Celsius; CI: Confidence interval; C-PCBs: Coplanar PCBs; LOD: limit of detection; M-PCBs: Mono-*ortho* PCBs; ng/g: nanogram per gram, 10⁻⁹ g; PCBs: Polychlorinated biphenyls; PCDDs: Polychlorinated dibenzo-*p*-dioxins; PCDFs: Polychlorinated dibenzofurans; pg/g: picogram per gram, 10⁻¹² g; SD: Standard deviation; SVZH: Middle Volga Chemical Plant; TEQ: Toxic equivalents; WHO: World Health Organization;

Acknowledgements

We gratefully thank the study participants, and the staff of the Chapaevsk Medical Association. The research described in this article has been reviewed by the National Institute of Environmental Health Sciences, and approved for publication. Approval does not signify that the contents necessarily reflect the views of the Agency, nor does the mention of trade names or commercial products constitute endorsement or recommendation for use. This work was funded by U.S. EPA grant R82943701 and NIEHS grants ES014370, ES00002, and 5T32-ES07069-28. OH was supported by 5T32ES016645-02 from NIEHS/NHGRI. MML is a member of the UMass DERC (DK32520). LA is a consultant for Environmental Health and Engineering, Inc. CE is an International Consultant and the President of BioSimulation Consulting Inc.

Author details

¹Environmental and Occupational Medicine and Epidemiology Program, Department of Environmental Health, Harvard School of Public Health, Boston, Massachusetts, USA. ²Department of Physical Education and Health, Samara State Medical University, Samara, Russia. ³Chapaevsk Medical Association, Chapaevsk, Russia. ⁴Department of Environmental Health, Harvard School of Public Health, Boston, Massachusetts. ⁵Environmental Health and Engineering, Inc., Needham, Massachusetts, USA. ⁶Channing Laboratory, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts, USA. ⁷Department of Biostatistics, Harvard School of Public Health, Boston, Massachusetts, USA. ⁸Department of Environmental and Occupational Health, Faculty of Medicine, University of Montreal, Montreal, Quebec, Canada. ⁹BioSimulation Consulting Inc., Newark, Delaware, USA. ¹⁰National Cancer Institute, National Institutes of Health, Department of Health and Human Services, Bethesda, Maryland, USA. ¹¹Pediatric Endocrinology Division, Departments of Pediatrics and Cell Biology, University of Massachusetts Medical School, Worcester, Massachusetts, USA. ¹²Center for Demography and Human Ecology, Institute for Forecasting, Russian Academy of Sciences, Moscow, Russia. ¹³Institute of Ecology and Evolution of Russian Academy of Sciences, Moscow, Russia.

Authors' contributions

OH participated in the study design, conducted the data analysis, and drafted the manuscript. OS supervised study recruitment and collection of blood samples and questionnaire data, and contributed to the drafting of the manuscript and the revisions. LA participated in the study design and coordination, and contributed to the manuscript revisions. SK, PW, CE, LB, JB, and ML assisted with the interpretation of the data and contributed to the drafting of the manuscript and the revisions. BR facilitated the data collection and reviewed the manuscript. AS and DF conducted the mass spectroscopy analyses and contributed to the manuscript revisions. RH conceived of the study, participated in its design and coordination, and helped draft and revise the manuscript. All authors read and approved the final version.

Competing interests

The authors declare that they have no competing interests.

Received: 24 November 2010 Accepted: 22 June 2011

Published: 22 June 2011

References

1. ATSDR: Toxicological Profile for Polychlorinated Biphenyls (PCBs). Atlanta, GA; 2000.
2. Schecter A, Birnbaum L, Ryan JJ, Constable JD: Dioxins: an overview. *Environ Res* 2006, **101**:419-428.
3. Milbrath MO, Wenger Y, Chang CW, Emond C, Garabrant D, Gillespie BW, Jolliet O: Apparent half-lives of dioxins, furans, and polychlorinated biphenyls as a function of age, body fat, smoking status, and breast-feeding. *Environ Health Perspect* 2009, **117**:417-425.
4. Duarte-Davidson R, Jones KC: Polychlorinated biphenyls (PCBs) in the UK population: estimated intake, exposure and body burden. *Sci Total Environ* 1994, **151**:131-152.
5. Hansen L, DeCaprio A, Nisbet I: PCB congener comparisons reveal exposure histories for residents of Anniston, Alabama, USA. *Fresenius Environmental Bulletin* 2003, **12**:181-190.
6. Kocan A, Petrik J, Jursa S, Chovanцова J, Drobna B: Environmental contamination with polychlorinated biphenyls in the area of their former manufacture in Slovakia. *Chemosphere* 2001, **43**:595-600.
7. Revich B, Aksel E, Ushakova T, Ivanova I, Zhuchenko N, Klyuev N, Brodsky B, Sotskov Y: Dioxin exposure and public health in Chapaevsk, Russia. *Chemosphere* 2001, **43**:951-966.
8. Zheng GJ, Leung AO, Jiao LP, Wong MH: Polychlorinated dibenzo-p-dioxins and dibenzofurans pollution in China: sources, environmental levels and potential human health impacts. *Environ Int* 2008, **34**:1050-1061.
9. Patterson DG Jr, Turner WE, Caudill SP, Needham LL: Total TEQ reference range (PCDDs, PCDFs, cPCBs, mono-PCBs) for the US population 2001-2002. *Chemosphere* 2008, **73**:S261-277.
10. Patterson DG Jr, Wong LY, Turner WE, Caudill SP, Dipietro ES, McClure PC, Cash TP, Osterloh JD, Pirkle JL, Sampson EJ, Needham LL: Levels in the U.S. population of those persistent organic pollutants (2003-2004) included in the Stockholm Convention or in other long range transboundary air pollution agreements. *Environ Sci Technol* 2009, **43**:1211-1218.
11. Furst P: Dioxins, polychlorinated biphenyls and other organohalogen compounds in human milk. Levels, correlations, trends and exposure through breastfeeding. *Mol Nutr Food Res* 2006, **50**:922-933.
12. Burns JS, Williams PL, Sergeev O, Korrick S, Lee MM, Revich B, Altshul L, Patterson DG Jr, Turner WE, Needham LL, Saharov I, Hauser R: Predictors of serum dioxins and PCBs among peripubertal Russian boys. *Environ Health Perspect* 2009, **117**:1593-1599.
13. Humblet O, Williams PL, Korrick SA, Sergeev O, Emond C, Birnbaum LS, Burns JS, Altshul L, Patterson DG, Turner WE, Lee MM, Revich B, Hauser R: Predictors of serum dioxin, furan, and PCB concentrations among women from Chapaevsk, Russia. *Environ Sci Technol* 2010, **44**:5633-5640.
14. Sergeev O, Saharov I, Shelepchikov AA, Revich B, Sotskov Y, Brodsky E, Denisova T, Feshin D, Zeilert V: Levels of PCDDs/PCDFs in the environment and food after 3 years of full plant inactivity, Chapaevsk, Russia. *Organohalogen Compounds* 2007, **69**:1388-1391.
15. Hauser R, Williams P, Altshul L, Korrick S, Peebles L, Patterson DG Jr, Turner WE, Lee MM, Revich B, Sergeev O: Predictors of serum dioxin levels among adolescent boys in Chapaevsk, Russia: a cross-sectional pilot study. *Environ Health* 2005, **4**:8.
16. US Environmental Protection Agency Method 1613: Tetra- through Octa-Chlorinated Dioxins and Furans by Isotope Dilution HRGC/HRMS. [http://www.epa.gov/region03/1613.pdf].
17. US Environmental Protection Agency Method 1668B: Chlorinated Biphenyl Congeners in Water, Soil, Sediment, Biosolids, and Tissue by HRGC/HRMS. [http://water.epa.gov/scitech/methods/cwa/bioindicators/upload/2009_01_07_methods_method_1668.pdf].
18. Phillips DL, Pirkle JL, Burse WW, Bernert JT Jr, Henderson LO, Needham LL: Chlorinated hydrocarbon levels in human serum: effects of fasting and feeding. *Arch Environ Contam Toxicol* 1989, **18**:495-500.
19. Baccarelli A, Pfeiffer R, Consonni D, Pesatori AC, Bonzini M, Patterson DG Jr, Bertazzi PA, Landi MT: Handling of dioxin measurement data in the presence of non-detectable values: overview of available methods and their application in the Seveso chloracne study. *Chemosphere* 2005, **60**:898-906.
20. Hornung RW, Reed LD: Estimation of average concentration in the presence of non-detectable values. *Applied Occupational Environmental Hygiene* 1990, **5**:48-51.
21. Van den Berg M, Birnbaum LS, Denison M, De Vito M, Farland W, Feeley M, Fiedler H, Hakansson H, Hanberg A, Haws L, Rose M, Safe S, Schrenk D, Tohyama C, Tritscher A, Tuomisto J, Tysklind M, Walker N, Peterson RE: The 2005 World Health Organization reevaluation of human and Mammalian toxic equivalency factors for dioxins and dioxin-like compounds. *Toxicol Sci* 2006, **93**:223-241.
22. Computation of Correlation Coefficient and Its Confidence Interval in SAS. Paper 170-31. [http://www2.sas.com/proceedings/sugi31/170-31.pdf].
23. Wolff MS, Britton JA, Teitelbaum SL, Eng S, Deych E, Ireland K, Liu Z, Neugut AI, Santella RM, Gammon MD: Improving organochlorine biomarker models for cancer research. *Cancer Epidemiol Biomarkers Prev* 2005, **14**:2224-2236.
24. NIOSH Manual of Analytical Methods, POLYCHLORINATED BIPHENYLS in serum: METHOD 8004. [http://www.cdc.gov/niosh/docs/2003-154/pdfs/8004.pdf].
25. Lorber M: A pharmacokinetic model for estimating exposure of Americans to dioxin-like compounds in the past, present, and future. *Sci Total Environ* 2002, **288**:81-95.
26. Lorber M, Patterson D, Huwe J, Kahn H: Evaluation of background exposures of Americans to dioxin-like compounds in the 1990s and the 2000s. *Chemosphere* 2009, **77**:640-651.

doi:10.1186/1476-069X-10-62

Cite this article as: Humblet et al.: Temporal trends in serum concentrations of polychlorinated dioxins, furans, and PCBs among adult women living in Chapaevsk, Russia: a longitudinal study from 2000 to 2009. *Environmental Health* 2011 **10**:62.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

