

5-12-2006

# Hypothermia with J (Osborne) waves

David H. Spodick

*University of Massachusetts Medical School*

Follow this and additional works at: <http://escholarship.umassmed.edu/oapubs>

 Part of the [Cardiology Commons](#)

---

## Repository Citation

Spodick, David H., "Hypothermia with J (Osborne) waves" (2006). *Open Access Articles*. 15.  
<http://escholarship.umassmed.edu/oapubs/15>

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in Open Access Articles by an authorized administrator of eScholarship@UMMS. For more information, please contact [Lisa.Palmer@umassmed.edu](mailto:Lisa.Palmer@umassmed.edu).

# Hypothermia With J (Osborne) Waves

David H. Spodick, MD, DSc

[click an author to search for more articles by that author](#)

A 40-year-old woman was found unconscious outside on a bitterly cold day; her body temperature was 83°F. She was never able to provide a coherent history. J (Osborne) waves are seen at the QRS-T junctions, especially in leads II, III, aVF, and V<sub>2</sub>–V<sub>6</sub> (arrows).

Osborne waves are characteristic of deep hypothermia, whether accidental (as seen here) or medically induced, in both humans and animals. A similar configuration may occur in severe hypercalcemia, nervous system disorders, and other conditions. Hypothermia affects both depolarization and repolarization, and the QT interval is prolonged. The mechanism of production of J waves is poorly understood, but they may be related to the transmural potential gradient at the onset of repolarization, which has been seen in vitro.

J-wave size has been related to the degree of hypothermia.

*The American Heart Hospital Journal. 4;2:156 (Spring 2006)*

