

May 22nd, 4:30 PM - 6:00 PM

Mood Disorders and Trauma: What are the Associations?

Yael Dvir

University of Massachusetts Medical School

Michael Hill

University of Massachusetts Medical School

Steven M. Hodge

University of Massachusetts Medical School

See next page for additional authors

Follow this and additional works at: https://escholarship.umassmed.edu/cts_retreat

Part of the [Mental Disorders Commons](#), [Pediatrics Commons](#), [Psychiatry Commons](#), and the [Trauma Commons](#)

Dvir, Yael; Hill, Michael; Hodge, Steven M.; and Frazier, Jean A., "Mood Disorders and Trauma: What are the Associations?" (2012). *UMass Center for Clinical and Translational Science Research Retreat*. 16.
https://escholarship.umassmed.edu/cts_retreat/2012/posters/16

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.

Presenter Information

Yael Dvir, Michael Hill, Steven M. Hodge, and Jean A. Frazier

Creative Commons License

Creative

Commons

This work is licensed under a [Creative Commons Attribution-Noncommercial-Share Alike 3.0 License](#).

Attribution-

Noncommercial-

Share

Alike

3.0

License

MOOD DISORDERS AND TRAUMA - WHAT ARE THE ASSOCIATIONS?

Yael Dvir M.D., Michael Hill B.S, Steven M Hodge M.A., Jean A Frazier M.D.

University of Massachusetts Medical School, Community Health Link

Contact: Email: Yael.Dvir@umassmed.edu

ABSTRACT

Objectives: Mood dysregulation in traumatized children may be misdiagnosed as bipolar disorder (BD) and conversely, the diagnosis of BD overlooked. Our aim is to characterize the relationship between trauma and mood dysregulation and pediatric BD.

Methods: We are assessing youths ages 8-18 who present with mood symptoms and past trauma divided into two groups: 1. Trauma+Unmodified DSM-IV-TR BD (T+BD) and 2. Trauma+Mood Disorder NOS (T+MD). Differences in clinical variables between groups are analyzed using t-tests for continuous and chi-square tests for categorical variables ($\alpha=0.05$).

Results: Age at onset of trauma for youth with T+BD (n=10) compared with T+MD (n=10) was similar (2.6 ± 1.8 versus 3.3 ± 1.9 years; $p=0.4$) as were types of trauma and number of incidents, and age at onset of mood symptoms (T+BD 7 ± 2.5 versus T+MD 7.8 ± 1.8 $p=0.4$). The T+BD group had higher scores on the sexual abuse subscale of the Childhood Trauma Questionnaire ($p=0.04$) and BPRS mania subscale ($p=0.02$), and higher total number of major depressive episodes ($p=0.04$) and manic episodes ($p=0.03$) per the KSCID. Youth with T+BD reported a trend toward higher rates of ideation to self-harm compared to youth with T+MD ($p=0.08$). Both groups had similar PTSD and ADHD symptoms, and similar number of psychotropic medications (BD 3.6 ± 2.9 MD 2.7 ± 2.1 $p=0.4$). Finally, family history findings suggest a trend towards higher rates of any Axis I disorder in the T+BD families ($p=0.07$), and significantly higher rates of anxiety disorders ($p=0.05$), BD ($p=0.04$), and schizophrenia ($p=0.02$).

Conclusions: Results suggest differences in clinical presentation and higher rates of BD and schizophrenia in the T+BD families. Taken together, these preliminary results suggest potential biological and genetic vulnerabilities which may predispose children to develop specific mood disorders under certain circumstances; the ability to identify these children early on could change their prognostic trajectory.