

2013

# Mindfulness-Based Stress Reduction and Change in Health-Related Behaviors

Elena Salmoirago-Blotcher  
*University of Massachusetts Medical School*

Matthew Hunsinger  
*University of Rochester*

Lucas Morgan  
*University of Massachusetts Boston*

*See next page for additional authors*

Follow this and additional works at: [https://escholarship.umassmed.edu/cardio\\_pp](https://escholarship.umassmed.edu/cardio_pp)

 Part of the [Behavior and Behavior Mechanisms Commons](#), [Community Health and Preventive Medicine Commons](#), [Movement and Mind-Body Therapies Commons](#), and the [Preventive Medicine Commons](#)

## Repository Citation

Salmoirago-Blotcher, Elena; Hunsinger, Matthew; Morgan, Lucas; Fischer, Daniel; and Carmody, James, "Mindfulness-Based Stress Reduction and Change in Health-Related Behaviors" (2013). *Cardiovascular Medicine Publications and Presentations*. 87.  
[https://escholarship.umassmed.edu/cardio\\_pp/87](https://escholarship.umassmed.edu/cardio_pp/87)

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in Cardiovascular Medicine Publications and Presentations by an authorized administrator of eScholarship@UMMS. For more information, please contact [Lisa.Palmer@umassmed.edu](mailto:Lisa.Palmer@umassmed.edu).

---

# Mindfulness-Based Stress Reduction and Change in Health-Related Behaviors

**Authors**

Elena Salmoirago-Blotcher, Matthew Hunsinger, Lucas Morgan, Daniel Fischer, and James Carmody

**Keywords**

mindfulness, mindfulness-based stress reduction, health behaviors

**Rights and Permissions**

Copyright The Author(s). This is the authors' initial version of the paper before peer review as submitted to: Journal of Evidence-Based Complementary and Alternative Medicine.

**Mindfulness-Based Stress Reduction and Change in Health-Related Behaviors**

Journal:	<i>Journal of Evidence-Based Complementary &amp; Alternative Medicine</i>
Manuscript ID:	Draft
Manuscript Type:	Original Manuscript
Date Submitted by the Author:	n/a
Complete List of Authors:	Salmoirago-Blotcher, Elena; University of Massachusetts Medical School, Medicine Hunsinger, Matthew; University of Rochester School of Medicine and Dentistry, Anesthesiology Morgan, Lucas; University of Massachusetts, Clinical Psychology Fischer, Daniel; University of Massachusetts Medical School, Medicine Carmody, James; University of Massachusetts Medical School, Medicine
Keywords:	mindfulness, mindfulness-based stress reduction, health behaviors
Abstract:	<p>How best to support change in health-related behaviors is an important public health challenge. The role of mindfulness training in this process has received limited attention.</p> <p>We sought to explore whether mindfulness training is associated with changes in health-related behaviors. The Health Behaviors Questionnaire was used to obtain self-reported dietary behaviors, drinking, smoking, physical activity and sleep quality before and after attendance at an eight-week Mindfulness-Based Stress Reduction program. T-test for paired data and chi-square were used to compare pre-post intervention means and proportions of relevant variables with <math>p = .05</math> as level of significance. Participants (<math>n = 174</math>; mean age 47 years, range: 19-68; 61 % female) reported significant improvements in dietary behaviors and sleep quality. Partial changes were seen in drinking and physical activity, and no change in smoking. In conclusion, mindfulness training promotes favorable changes in selected health-related behaviors deserving further study through randomized controlled trials.</p>

1  
2  
3 Mindfulness-Based Stress Reduction and Change in Health-Related Behaviors  
4  
5  
6  
7

8 Elena Salmoirago-Blotcher, MD, PhD<sup>a)</sup>; Matthew Hunsinger, PhD<sup>b)</sup>; Lucas Morgan,  
9 MA<sup>c)</sup>, Daniel Fischer, BA<sup>a)</sup>; and James Carmody, PhD<sup>d)</sup>  
10  
11

12 <sup>a)</sup> Division of Cardiovascular Medicine, University of Massachusetts Medical School,  
13 Worcester, MA; <sup>b)</sup> Department of Anesthesiology, University of Rochester School of  
14 Medicine and Dentistry; <sup>c)</sup> Department of Clinical Psychology, University of  
15 Massachusetts, Boston; <sup>d)</sup> Division of Preventive and Behavioral Medicine, University of  
16 Massachusetts Medical School, Worcester, MA  
17  
18  
19  
20  
21  
22  
23

24  
25  
26  
27 *Corresponding author:*  
28

29 Elena Salmoirago-Blotcher, MD, PhD  
30

31 Division of Cardiovascular Medicine  
32

33 University of Massachusetts Medical School  
34

35 55 Lake Avenue North, room S3-855  
36

37 Worcester , MA 01655 - USA  
38

39 Phone: 508-856-4413; Fax: 508-856-4571  
40

41 Email: [Elena.Salmoirago-Blotcher@umassmed.edu](mailto:Elena.Salmoirago-Blotcher@umassmed.edu)  
42  
43  
44  
45  
46  
47

48 *Word count: 2,258*  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

### Abstract

How best to support change in health-related behaviors is an important public health challenge. The role of mindfulness training in this process has received limited attention. We sought to explore whether mindfulness training is associated with changes in health-related behaviors. The Health Behaviors Questionnaire was used to obtain self-reported dietary behaviors, drinking, smoking, physical activity and sleep quality before and after attendance at an eight-week Mindfulness-Based Stress Reduction program. T-test for paired data and chi-square were used to compare pre-post intervention means and proportions of relevant variables with  $p = .05$  as level of significance. Participants ( $n = 174$ ; mean age 47 years, range: 19-68; 61 % female) reported significant improvements in dietary behaviors and sleep quality. Partial changes were seen in drinking and physical activity, and no change in smoking. In conclusion, mindfulness training promotes favorable changes in selected health-related behaviors deserving further study through randomized controlled trials.

*Key words:* mindfulness; mindfulness-based stress reduction; health behaviors

## Introduction

Health-related behaviors such as poor diets, physical inactivity, and smoking are major contributors to a range of disorders including obesity, the metabolic syndrome, type 2 diabetes, and cardiovascular disease.<sup>1,2</sup> Promoting sustainable change in these behaviors is however a seemingly intractable public health problem.

The capacity for self-regulation is central to making and sustaining behavioral change, and programs that support this capacity represent an opportunity to improve behavioral outcomes.<sup>3</sup> Mindfulness programs, including the widely disseminated Mindfulness-Based Stress Reduction Program,<sup>4</sup> have been shown to enhance emotion self-regulation.<sup>5,6</sup> However, the possible effect of mindfulness training on health-related behaviors has received limited attention. The aim of this observational study was to examine whether attendance in a Mindfulness-Based Stress Reduction program would be associated with changes in health-related behaviors (diet, smoking, drinking, physical activity) and sleep quality compared to baseline and whether any changes were associated with the duration of individual practice and improvement in mindfulness skills.

## Methods

### Population

Study participants belonged to several cohorts of individuals who enrolled in the University of Massachusetts Mindfulness-Based Stress Reduction program in Worcester, MA during 2006. Participants were adults with a wide range of health-related problems including chronic pain, anxiety, depression, and personal or employment-related stress. Each class included approximately 20–25 participants; health-care practitioners referred about half and others were self-referred. Participation in the program was on a self-pay

1  
2  
3 basis. Prospective participants attend an orientation session prior to the beginning of the  
4  
5 program during which they receive detailed information about the program format and  
6  
7 requirements. All participants are formally asked for permission to use their de-identified  
8  
9 information for research purposes.  
10  
11

### 12 **The Mindfulness Stress Reduction Program**

13  
14  
15 This program (described in detail elsewhere)<sup>4</sup> consists of seven weekly classes of  
16  
17 two and half hours, and an all-day weekend class during the sixth week. Mindfulness is  
18  
19 taught through three formal exercises: the body scan, in which attention is moved  
20  
21 systematically throughout the body with instructions to notice whatever sensations are  
22  
23 present; gentle stretching exercises to support mindfulness during movement; and sitting  
24  
25 meditation, during which the arising of cognitions is noticed. During each of these  
26  
27 modalities trainees are encouraged to bring a non-judgmental and non-reactive  
28  
29 observational stance to whatever is arising in awareness. Up to 45 minutes/day of  
30  
31 individual practice of these exercises is also prescribed, using instructions contained in  
32  
33 CD recordings. Participants are also encouraged to integrate mindfulness into their  
34  
35 everyday activities.  
36  
37  
38  
39

### 40 **Assessments**

41  
42  
43 Baseline assessments were completed immediately prior to the orientation  
44  
45 sessions, while post-program questionnaires were completed during the final program  
46  
47 session. Home practice data were collected by means of a mindfulness practice log in  
48  
49 which participants recorded the number of minutes of formal and informal mindfulness  
50  
51 practice they did each day. Whenever a log was missing, participants were asked to  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 retrospectively complete a log for the previous week. When data on individual practice  
4  
5 were missing we assumed that the minutes of practice for that day were zero.  
6  
7

### 8 *Primary outcomes*

9  
10 Health behaviors were self-reported through the Health Behaviors Questionnaire,<sup>7</sup>  
11  
12 an instrument developed as a pragmatic measure for use in intervention-based research  
13  
14 conducted in primary care settings. It consists of 22 items selected from known health  
15  
16 behavior scales identified through literature reviews and previous multi-site studies: nine  
17  
18 items relate to physical activity and are derived from the Rapid Assessment of Physical  
19  
20 Activity (RAPA) questionnaire,<sup>8</sup> three relate to alcohol consumption (from the  
21  
22 Behavioral Risk Factor Surveillance System Survey Questionnaire),<sup>9</sup> three to cigarette  
23  
24 smoking,<sup>9</sup> and seven (derived from the “Starting the Conversation” questionnaire)<sup>10</sup> to  
25  
26 dietary patterns. The Health Behaviors Questionnaire’s authors added a final item  
27  
28 assessing sleep quality during the previous week (using a six-point Likert response scale  
29  
30 ranging from 'very poor' to 'very good').  
31  
32  
33  
34  
35

36 Demographic characteristics were self-reported and included age, gender, marital  
37  
38 status, use of prescription medications and prior psychotherapy. Mindfulness was  
39  
40 measured using the Five Facets of Mindfulness questionnaire,<sup>11</sup> an instrument derived  
41  
42 from a factor analysis of questionnaires measuring a trait-like general tendency to be  
43  
44 mindful in daily life. Items are rated on a Likert scale ranging from 1 (never or very  
45  
46 rarely true) to 5 (very often or always true). This instrument has been shown to have good  
47  
48 internal consistency and significant relationships in the predicted directions with a variety  
49  
50 of constructs related to mindfulness.<sup>12</sup>  
51  
52  
53  
54

### 55 **Data analysis**



1  
2  
3 For the three drinking questions, a score of 0 was assigned to the second and third  
4 questions for anyone who answered 'no' to the first question. A sum variable was created  
5 by summing responses across the three questions. For changes in smoking behaviors, the  
6 first question "Have you smoked at least 100 cigarettes in your entire life" was excluded  
7 because it was not relevant to the assessment of behavioral change. For the second  
8 question, a score of 0 was assigned to 'no' responses and a score of 1 to a 'yes' response.  
9 The third smoking question was treated as a continuous score. To assess dietary  
10 behaviors, responses were re-coded so that the 'none' response was at the beginning of the  
11 scale; some of the questions were also reverse-coded such that, for most of the  
12 continuous indicators of health behaviors, higher scores would indicate worse (i.e., less  
13 healthy) behaviors. For physical activity, responses were computed according to the  
14 original scoring method.<sup>8</sup> Participants were assigned to a category of physical activity  
15 (sedentary, underactive, underactive-regular, active) based on their responses to questions  
16 1 through 7. Responses for items 8 and 9 were summed to create a single flexibility score,  
17 where higher scores reflected greater strength and flexibility. Sleep quality was assessed  
18 with a single question that was treated as a continuous variable.

19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41 Exploratory analyses aimed at the identification of possible mechanisms of the  
42 association between mindfulness training and changes in behaviors. Since mindfulness  
43 scores and behaviors were collected at the same time, a formal mediation analysis was  
44 not conducted and only associations were considered.

45  
46  
47  
48  
49  
50  
51 Two-tailed t-test for paired data was used to compare pre-post intervention means  
52 of relevant continuous variables and chi-square to compare proportions. Spearman rho  
53 was used to examine associations between change in behaviors and change in  
54  
55  
56  
57  
58  
59  
60

mindfulness scores and duration of individual practice, respectively. Results are reported as mean pre-post training change with 95 % confidence intervals ( $p = 0.05$  level of significance). All data analyses were conducted using SPSS statistical software version 20.

## Results

Complete baseline and post-training data were available for 174 participants (Figure 1). The mean age was 47 (SD = 10.3 years; range: 19 - 68); 125 participants (61%) were female, and 76% were married or living as married. Most participants reported being in white collar/professional occupations. Over 60% of participants reported current or previous psychotherapy and 70% reported the use of prescription medications. Overall, participants reported engaging in healthy behaviors at baseline, as indicated by the low average number of sweetened beverages, snacks, and dessert servings per week and by the low prevalence of current smokers.

No significant baseline differences were found between the participants who failed to provide post-treatment data ( $n = 32$ ) and the rest of the sample ( $n = 174$ ) for demographic variables or for mindfulness scores.

### *Program attendance*

Of the 174 participants with complete data, 168 (97%) attended six or more of the eight weekly sessions, whereas five (4%) participants attended five or fewer sessions.

### *Home mindfulness practice*

About 70% (121/174) of study participants provided some or their entire home practice data. The average number of returned logs was 6.16 (SD = 1.34) out of 7 logs.

### **Change in health-related behaviors**

### *Drinking*

There were no changes in the overall drinking score (Table 1). When examining specific drinking behaviors, participants reported a reduction in the number of days they drank during the previous month, but no changes in the number of drinks they had on those drinking days or in the number of binge drinking episodes.

### *Smoking*

Participation in mindfulness training was neither associated with pre-post intervention changes in the proportion of participants who smoked at least part of a cigarette in the previous week, nor with significant changes in the number of cigarettes participants typically smoked in the previous week.

### *Dietary behaviors*

The overall dietary score significantly improved after completion of the program, indicating an improvement in overall dietary behaviors. When looking at individual responses, participants reported a significant reduction in the number of desserts consumed, as well as an increase in the intake of fruits or vegetables. We also observed marginally significant reductions in the number of sweetened beverages, fast food meals, and in the use of fats.

### *Physical activity*

The proportion of participants in the “sedentary” category decreased after the completion of the program. Participants also reported an increase in strength/flexibility scores by the end of the training.

### *Sleep quality*

1  
2  
3 Examination of sleep quality scores suggested that the quality of sleep improved  
4 across the training (table 1).  
5  
6

### 7 8 **Exploratory analyses** 9

#### 10 *Change in mindfulness skills and health-related behaviors* 11

12 We did not detect correlations between changes in overall mindfulness score or  
13 the various facets of mindfulness and changes in health behaviors, with the exception of  
14 sleep quality, which was positively associated with change in overall mindfulness scores  
15 (rho = 0.30,  $p < 0.01$ ) as well as with changes in three of the subscales: Observe (rho =  
16 0.28,  $p = 0.01$ ), Acting with Awareness (rho = 0.19,  $p = 0.01$ ), and React (rho = 0.21,  $p <$   
17 0.01).  
18  
19  
20  
21  
22  
23  
24  
25

#### 26 *Associations between individual mindfulness practice and changes in behaviors* 27

28 Higher smoking scores (indicating worse smoking behavior) were inversely  
29 associated with the number of days spent in sitting meditation (rho = - 0.23,  $p = 0.04$ ),  
30 yoga practice (rho = - 0.30,  $p = 0.01$ ), and informal meditation (rho = - 0.25,  $p = 0.03$ ).  
31  
32 Individual home practice was not associated with changes in dietary behavior, drinking,  
33 or physical activity. The improvement in sleep quality was related to the amount of time  
34 participants spent in informal practice (rho = 0.23,  $p = 0.01$ ), but not with the duration of  
35 the other individual practices.  
36  
37  
38  
39  
40  
41  
42  
43  
44

### 45 46 **Discussion** 47

48 Participation in a standard Mindfulness Based Stress Reduction program was  
49 associated with improved dietary behaviors (specifically, decreased consumption of  
50 desserts and increased intake of fruits and vegetables), modest changes in physical  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 activity and drinking habits, and better sleep quality compared to baseline. No changes  
4  
5 were observed in smoking behaviors.  
6  
7

8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

There are only a limited number of studies examining the association between mindfulness and health behaviors. Large survey-based studies in college populations have shown that higher dispositional mindfulness (i.e., the capacity that individuals have to be mindful prior to training)<sup>13, 14</sup> is associated with increased physical activity, better sleep quality<sup>15</sup> and healthier dietary habits.<sup>16</sup> While preliminary studies investigating the effect of mindfulness interventions alone on dietary habits in adults are either not conclusive<sup>17</sup> or did not show an effect on the dietary intake of several nutrients,<sup>18</sup> mindfulness training delivered as part of a vegetable-based dietary intervention decreased the consumption of saturated fat and animal protein, increased the intake of vegetable protein, and improved physical activity in men with recurrent prostate cancer; changes that were self-maintained three months post-intervention.<sup>19-21</sup> Of note, the current study sample had overall good dietary behaviors at baseline, with an average of just over one fast food meal and unhealthy snack per week, resulting in a possible floor effect. A floor effect may also explain the lack of improvement in smoking behaviors, as the prevalence of smokers in this study was low. In fact, pilot studies of mindfulness interventions for smoking cessation have shown promising results on point prevalence abstinence rate.<sup>22,23</sup> Interestingly, one of these studies detected a positive association between compliance with mindfulness practice and smoking abstinence.<sup>23</sup>

There is no conclusive agreement in the literature in regard to the effects of mindfulness training on sleep. While a review of the literature has suggested that the positive effects of mindfulness training on sleep quality and duration have yet to be

1  
2  
3 demonstrated,<sup>24</sup> this study confirms more recent findings suggesting an effect of  
4 mindfulness training on sleep quality among breast cancer patients (although not  
5 maintained over time)<sup>25</sup> and in menopausal women with hot flashes.<sup>26</sup>  
6  
7

8  
9  
10 The change in the proportion of sedentary individuals observed in this study was  
11 driven by an increase in activities that promote flexibility. Since such activities (yoga) are  
12 taught as part of the Mindfulness-Based Stress Reduction curriculum, no conclusions can  
13 be drawn as to whether the training improved overall physical activity.  
14  
15

16  
17 This study has several limitations. Due to its observational design and the lack of  
18 a control group, it cannot be ruled out that the observed changes are due to unspecific  
19 effects (i.e. generic support deriving from participation in a group-based program;  
20 participation in the program being part of a more general decision to improve health) that  
21 are unrelated to mindfulness training. Second, the study was conducted in individuals  
22 who voluntarily enrolled in a standard Mindfulness-Based Stress Reduction program.  
23 Such individuals differed from the general population in a number of respects, including  
24 that they mostly had white-collar professions and had a high prevalence of healthy  
25 behaviors at baseline. Third, all behaviors were self-reported; and fourth, we cannot  
26 exclude the presence of social desirability bias. Finally, due to the lack of follow-up  
27 assessments, it is unclear whether the observed changes would be maintained over time.  
28  
29

30  
31 In conclusion, this study found that participation in a Mindfulness-Based Stress  
32 Reduction program was associated with changes in health-related behaviors, particularly  
33 in dietary behaviors, and in sleep quality. These encouraging findings deserve further  
34 study in larger randomized controlled trials to explore the possible role of mindfulness  
35 training in initiating and maintaining behavioral change.  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 **Acknowledgements:** none  
4

5 **Authors' contributions:**  
6

7  
8 Elena Salmoirago-Blotcher conceived the study and wrote the manuscript;  
9  
10 Matthew Hunsinger conducted the statistical analysis and revised the manuscript; Lucas  
11  
12 Morgan conducted the statistical analysis; Daniel Fischer conducted literature reviews  
13  
14 and assisted with the manuscript drafting; James Carmody conceived the study, collected  
15  
16 the data and revised the manuscript. All authors approved the final version of the  
17  
18 manuscript for publication. All authors approved the final version of the  
19  
20 manuscript for publication.  
21

22 **Conflicting interests:** none reported  
23

24 **Ethical approval:** All participants gave their permission to use their de-identified  
25  
26 information for research purposes.  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## References

1. Anderson LH, Martinson BC, Crain AL, et al. Health care charges associated with physical inactivity, overweight, and obesity. *Prev Chronic Dis*. 2005; 2:A09.
2. Glasgow RE, Kaplan RM, Ockene JK, Fisher EB, Emmons KM. Patient-reported measures of psychosocial issues and health behavior should be added to electronic health records. *Health Aff (Millwood)*. 2012; 31:497-504.
3. Williams PG, Thayer JF: Executive functioning and health: introduction to the special series. *Ann Behav Med*. 2009; 37:101-105.
4. Kabat-Zinn J. Full catastrophe living: using the wisdom of your body and mind to face stress, pain and illness. New York: Delacorte, 1990.
5. Goldin P, Gross J. Effect of mindfulness meditation training on the neural bases of emotion regulation in social anxiety disorder. *Emotion*. 2010;10:83-91.
6. Chambers R, Gullone E, Allen NB. Mindful emotion regulation: An integrative review. *Clinical Psychology Review*. 2009; 29:560-572.
7. Glasgow RE, Ory MG, Klesges LM, et al. Practical and relevant self-report measures of patient health behaviors for primary care research. *Ann Fam Med*. 2005; 3:73-81.
8. Topolski TD, LoGerfo J, Patrick DL, et al.: The Rapid Assessment of Physical Activity (RAPA) among older adults. *Prev Chronic Dis*. 2006; 3:A118.
9. Center for Disease Control and Prevention. *Behavioral Risk Factor Surveillance System Survey Questionnaire*. Atlanta, Georgia: U.S. Department of Health and Human Services, 2011.



10. Paxton AE, Strycker LA, Toobert DJ, Ammerman AS, Glasgow RE. Starting the conversation: Performance of a brief dietary assessment and intervention tool for health professionals. *American Journal of Preventive Medicine*. 2011; 40:67-71.
11. Baer RA, Smith GT, Hopkins J, Krietemeyer J, Toney L. Using self-report assessment methods to explore facets of mindfulness. *Assessment*. 2006; 13:27-45.
12. Baer RA, Smith GT, Lykins E, et al. Construct validity of the Five Facet Mindfulness Questionnaire in meditating and nonmeditating samples. *Assessment*. 2008; 15:329-342.
13. Brown KW, Ryan RM. The benefits of being present: mindfulness and its role in psychological well-being. *J Pers Soc Psychol*. 2003; 84:822-848.
14. Kabat-Zinn J. Mindfulness-based interventions in context: past, present, and future. *Clinical Psychology: Science and Practice*. 2003;10:144-156.
15. Roberts KC, Danoff-Burg S. Mindfulness and health behaviors: is paying attention good for you? *J Am Coll Health*. 2011; 59:165-173.
16. Gilbert D, Waltz J. Mindfulness and Health Behaviors. *Mindfulness*. 2010; 1:227-234.
17. Daubenmier J, Kristeller J, Hecht FM, et al. Mindfulness intervention for stress eating to reduce cortisol and abdominal fat among overweight and obese women: an exploratory randomized controlled study. *J Obes*. 2011; 2011:651936.
18. Kearney DJ, Milton ML, Malte CA, et al. Participation in mindfulness-based stress reduction is not associated with reductions in emotional eating or uncontrolled eating. *Nutr Res*. 2012;32:413-420.

19. Hebert JR, Hurley TG, Harmon BE, et al. A diet, physical activity, and stress reduction intervention in men with rising prostate-specific antigen after treatment for prostate cancer. *Cancer Epidemiology*. 2012; 36:e128-e136.
20. Carmody J, Olendzki B, Reed G, Andersen V, Rosenzweig P. A dietary intervention for recurrent prostate cancer after definitive primary treatment: results of a randomized pilot trial. *Urology*. 2008; 72:1324-1328.
21. Carmody JF, Olendzki BC, Merriam PA, et al. A novel measure of dietary change in a prostate cancer dietary program incorporating mindfulness training. *J Acad Nutr Diet*. 2012; 112:1822-1827.
22. Brewer JA, Mallik S, Babuscio TA, et al. Mindfulness training for smoking cessation: results from a randomized controlled trial. *Drug Alcohol Depend*. 2011; 119:72-80.
23. Davis JM, Fleming MF, Bonus KA, Baker TB. A pilot study on mindfulness based stress reduction for smokers. *BMC Complement Altern Med*. 2007; 7:2.
24. Winbush NY, Gross CR, Kreitzer MJ. The effects of mindfulness-based stress reduction on sleep disturbance: a systematic review. *Explore (NY)*. 2007, 3:585-591.
25. Andersen SR, Wurtzen H, Steding-Jessen M, et al. Effect of mindfulness-based stress reduction on sleep quality: Results of a randomized trial among Danish breast cancer patients. *Acta Oncol*. 2013; 52:336-344.
26. Carmody JF, Crawford S, Salmoirago-Blotcher E, et al. Mindfulness training for coping with hot flashes: results of a randomized trial. *Menopause*. 2011; 18:611-620.

Table 1. Pre-post program change in health behaviors

Behavior	Pre	Post	Change (mean, CI)	P*
<b>Drinking</b>				
<i>Overall drinking behavior score</i>	12.45 (11.50)	12.10 (11.29)	0.349 (- 0.76, 1.45)	0.540
Days drank at least 1 drink during past month	8.84 (9.30)	8.01 (8.81)	0.82 (0.10, 1.55)	0.026
Number of drinks on drinking days	1.29 (1.04)	1.34 (1.07)	- 0.05 (- 0.19, 0.10)	0.543
Binge drinking† episodes over past month	2.40 (4.17)	3.00 (4.60)	- 0.61 (- 1.42, .20)	0.141
<b>Smoking</b>				
Smoked at least part of a cigarette over past week	15 (23)	13 (24)		0.705‡
Number of cigarettes during past week	1.48 (17.31)	2.52 (13.72)	- 1.03 (- 2.38, 0.31)	0.13
<b>Dietary behaviors</b>				
<i>Overall dietary score</i>	7.99 (3.60)	7.24 (3.58)	0.75 (0.35, 1.15)	0.001
Fast food meals or snacks (times/week)	1.27 (1.16)	1.10 (1.01)	0.16 (- 0.02, 0.34)	0.082
Fruit/veggies servings (servings/day)	0.70 (.85)	0.55 (.71)	0.16 (0.04, 0.27)	0.006

Sodas or sweetened beverages (servings/day)	1.02 (1.28)	0.89 (1.16)	0.12 (- 0.05, 0.30)	0.156
Chicken, beans or fish (times/week)	0.59 (.97)	0.55 (.93)	0.04 (- 0.13, 0.20)	0.673
Snacks-chips or crackers (times/week)	1.14 (1.07)	1.15 (1.11)	- 0.01 (- 0.16, 0.13)	0.876
Desserts or other sweets (times/week)	2.15 (1.08)	1.96 (1.11)	0.18 (0.06, 0.31)	0.005
Use of margarine, butter or meat fat for seasoning (times/week)	1.25 (.60)	1.18 (.54)	0.07 (- 0.02, 0.15)	0.115
<b>Physical activity</b>				
‘Sedentary’	9 (5)	2 (1)		0.035‡
Rarely or no physical activity	19 (10)	11 (6)		0.021‡
‘Underactive’	104 (60)	94 (61)		0.477‡
Some light or moderate physical activity, but not every week	153 (76)	145 (83)		0.659‡
‘Underactive-regular’	25 (15)	27 (18)		0.782‡
Moderate physical activity every week, < 5	103 (51)	102 (59)		0.726‡

days per week or < 30 minutes at a time				
Vigorous physical activity every week, < 3 days per week or < 20 minutes at a time	41 (21)	38 (22)		0.547‡
'Active' Category	34 (20)	30 (20)		0.617‡
Moderate physical activity 30 minutes or more per day, 5 or more days/week	58 (28)	55 (32)		0.573‡
Moderate physical activity 20 or more minutes per day, 3 or more days/week	47 (24)	44 (26)		0.783‡
Overall strength/flexibility score	1.65 (1.30)	2.29 (.91)	- 0.64 (- 0.83, - 0.44)	0.001
Activities to increase muscle strength, such as lifting weights or calisthenics, once a week or more	0.47 (.99)	0.53 (.91)	- 0.03 (- 0.10, 0.03)	0.356
Activities to improve	0.53 (.24)	0.89 (.57)	- 0.63(- 0.80, -0.47)	0.001

flexibility, such as stretching or yoga, once a week or more				
<b>Sleep quality</b>				
Sleep quality scores	3.41 (0.10)	4.10 (0.10)	0.68 (0.47, 0.90)	0.01

Numbers are n (%) or means (SD)

Dietary and drinking score = sum of scores on individual answers, where a higher score indicates unhealthier dietary or drinking behaviors

\* Two-tailed T test for paired data unless otherwise specified

† Binge drinking defined as  $\geq 5$  drinks (men) or  $\geq 4$  drinks (women)

‡ Chi-square

Figure 1 - Flow of patients through the study

