

UMass Chan Medical School

eScholarship@UMassChan

UMass Center for Clinical and Translational
Science Research Retreat

2013 UMass Center for Clinical and
Translational Science Research Retreat

May 8th, 12:30 PM - 1:30 PM

Setting Proper Matrices for Accelerometry-based Physical Activity Measures in Older Women

Rachel Duckham

University of Massachusetts Medical School

Et al.

Let us know how access to this document benefits you.

Follow this and additional works at: https://escholarship.umassmed.edu/cts_retreat



Part of the Behavior and Behavior Mechanisms Commons, Community Health and Preventive Medicine Commons, Exercise Physiology Commons, Translational Medical Research Commons, and the Women's Health Commons

Repository Citation

Duckham R, Crouter S, Churchill LC, Kane KJ, Franklin PD, Li W. (2013). Setting Proper Matrices for Accelerometry-based Physical Activity Measures in Older Women. UMass Center for Clinical and Translational Science Research Retreat. <https://doi.org/10.13028/xjw0-k024>. Retrieved from https://escholarship.umassmed.edu/cts_retreat/2013/posters/62

Creative Commons License



This work is licensed under a [Creative Commons Attribution-NonCommercial-Share Alike 3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/). This material is brought to you by eScholarship@UMassChan. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMassChan. For more information, please contact Lisa.Palmer@umassmed.edu.

Setting Proper Matrices for Accelerometry-based Physical Activity Measures in Older Women

Rachel L Duckham,¹ Scott Crouter,² Linda Churchill¹, Kevin Kane¹, Patricia Franklin³, Wenjun Li¹

¹ Health Geography Lab, Division of Preventive & Behavioral Medicine, UMass Medical School

² College of Nursing and Health Sciences, UMass Boston

³ Department of Orthopedics and Physical Rehabilitation, UMass Medical School

Contact information: Wenjun.Li@umassmed.edu; 508-856-6574

BACKGROUND: Accelerometers to objectively measure physical activity (PA) among adults and youth are widespread, with few studies in the older population. Optimal PA cut-points in older adults who are often frail and/or have limited function still remain unclear.

Purpose: To compare the level of PA in older women with lower extremity problems using existing PA cut-points.

METHODS: Women (N=89) aged 65 to 85 years were instructed to wear an Actigraph accelerometer for seven consecutive days. Wear-time and activities away from the home were recorded daily. Questionnaires were used to assess participant sociodemographic attributes, lower extremity and comorbid conditions. PA was defined using cut-points identified in the literature (Freedson, Copeland (one axis) and Carr (vector)). Weekly minutes of sedentary, light, moderate and vigorous activity was compared among age groups, and those with or without lower extremity problems affecting walking.

RESULTS: Preliminary data of 72 women with valid accelerometry measures (≥ 6 consecutive days of ≥ 8 hours of daily wearing time). According to the Freedson cut-points, average weekly minutes of light, moderate and vigorous PA were 4,674 (612, 8,520), 24.0 (range 0, 1,050) and 6.6 (range: 0, 330), respectively. Application of Copeland and Carr cut-points yielded similar patterns. Such patterns were persistent when the analysis was stratified by age group (<75 vs. ≥ 75 y) and whether having any lower extremity problem affecting walking.

CONCLUSION: The existing cut-points for PA are ineffective in differentiation of active and inactive older. Age-appropriate cut-points for PA need to be established in older adults. Due to the lack of time spent in moderate/vigorous PA, health promotions may consider focusing on decreasing sedentary time and increasing time spent in light PA rather than promoting 150 minutes per week of moderate/vigorous PA in older adults.

Word count: 285