

University of Massachusetts Medical School

eScholarship@UMMS

UMass Center for Clinical and Translational
Science Research Retreat

2013 UMass Center for Clinical and
Translational Science Research Retreat

May 8th, 12:30 PM - 1:30 PM

Frequency and Impact of Intensive Care Unit Complications on Moderate-Severe Traumatic Brain Injury – Early Results of the Outcome Prognostication in Traumatic Brain Injury (OPTIMISM) Study

Susanne Muehlschlegel

University of Massachusetts Medical School

Et al.

Let us know how access to this document benefits you.

Follow this and additional works at: https://escholarship.umassmed.edu/cts_retreat



Part of the [Critical Care Commons](#), [Health Services Research Commons](#), [Nervous System Diseases Commons](#), [Neurology Commons](#), [Surgery Commons](#), [Translational Medical Research Commons](#), and the [Trauma Commons](#)

Repository Citation

Muehlschlegel S, Carandang RA, Ouillette C, Hall WR, Anderson F, Goldberg RJ. (2013). Frequency and Impact of Intensive Care Unit Complications on Moderate-Severe Traumatic Brain Injury – Early Results of the Outcome Prognostication in Traumatic Brain Injury (OPTIMISM) Study. UMass Center for Clinical and Translational Science Research Retreat. <https://doi.org/10.13028/478w-4h16>. Retrieved from https://escholarship.umassmed.edu/cts_retreat/2013/posters/24

Creative Commons License



This work is licensed under a [Creative Commons Attribution-NonCommercial-Share Alike 3.0 License](#).

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.

Frequency and Impact of Intensive Care Unit Complications on Moderate-Severe Traumatic Brain Injury – Early Results of the Outcome Prognostication in Traumatic Brain Injury (OPTIMISM) Study

Susanne Muehlschlegel, MD, MPH^{1,2,3}; Raphael Carandang, MD^{1,3}; Cynthia Ouillette, RN¹, Wiley Hall, MD^{1,3}; Fred Anderson, PhD^{3,4}; Robert Goldberg, PhD⁵

¹Departments of Neurology, Division of Neurocritical Care, ²Anesthesia/Critical Care and ³Surgery, ⁴Center for Outcomes Research, ⁵Department of Quantitative Health Sciences; University of Massachusetts Medical School, Worcester, MA

Contact information:

Susanne Muehlschlegel, MD, MPH

Assistant Professor of Neurocritical Care

Departments of Neurology (Division of Neurocritical Care), Anesthesiology/Critical Care and Surgery

University of Massachusetts Medical School

55 Lake Ave North

Worcester, MA 01655

Phone: 508-334-4651

FAX: 508-856-6778

Email: susanne.muehlschlegel@umassmemorial.org

ABSTRACT

Background: Known predictors of adverse outcomes in patients with moderate-severe TBI (msTBI) explain only a relatively small proportion of patient-related outcomes. The frequency and impact of intensive care unit complications (ICU-COMPL) on msTBI-associated outcomes is poorly understood.

Methods: In 213 consecutive msTBI patients admitted to a Level-I-Trauma-Center neuro-trauma-ICU, twenty-eight ICU-COMPL (21 medical and 7 neurological) were prospectively collected and adjudicated by group consensus, using pre-defined criteria. We determined frequencies, and explored associations of ICU-COMPL and functional neurological outcomes measured by Glasgow Outcome Scale (GOS) at hospital discharge using multivariable logistic regression.

Results: The average age of the study sample was 53 years, and the median presenting Glasgow Coma Scale and Injury Severity Scores were 5 and 27, respectively. Hyperglycemia (79%), fever (62%), systemic inflammatory response syndrome (60%), and hypotension requiring vasopressors (42%) were the four most common medical ICU-COMPL. Herniation (39%), intracranial rebleed (39%), and brain edema requiring osmotherapy (37%) were the three most common neurological ICU-COMPL. After adjusting for admission variables, duration of ventilation, and ICU length-of-stay, patients with brain edema (OR 5.8; 95% CI 2,16.7) had a significantly increased odds for dying during hospitalization whereas patients with hospital-acquired urinary tract infection (UTI) had a decreased odds (OR 0.05; 95% CI 0.005,0.6). Sensitivity-analysis revealed that UTI occurred later, suggesting a non-causal association with survival. Brain herniation (OR 15.7; 95% CI 2.6,95.4) was associated with an unfavorable functional status (GOS 1-3).

Conclusion: ICU-COMPL are very common after msTBI, have a considerable impact on short-term outcomes, and should be considered in the prognostication of these high-risk patients. Survival associations of time-dependent complications warrant cautious interpretation.

