

May 20th, 12:30 PM

Adipose Tissue Inflammation in Diabetic versus Non-diabetic Pregnancies


Tiffany A. Moore Simas
University of Massachusetts Medical School

Raziel Rojas-Rodriguez
University of Massachusetts Medical School

Jacqueline Tessa Draper
University of Massachusetts Medical School

See next page for additional authors

Follow this and additional works at: https://escholarship.umassmed.edu/cts_retreat

 Part of the [Cellular and Molecular Physiology Commons](#), [Endocrine System Diseases Commons](#), [Endocrinology Commons](#), [Endocrinology, Diabetes, and Metabolism Commons](#), [Maternal and Child Health Commons](#), [Nutritional and Metabolic Diseases Commons](#), [Obstetrics and Gynecology Commons](#), [Reproductive and Urinary Physiology Commons](#), [Translational Medical Research Commons](#), and the [Women's Health Commons](#)

Moore Simas, Tiffany A.; Rojas-Rodriguez, Raziel; Draper, Jacqueline Tessa; Sert, Aylin; Kruse, Maxwell; Boeras, Crina; Rosenthal, Brittany; Gaelikman, Olga; Min, So-Yun; Leung, Katherine G.; and Corvera, Silvia, "Adipose Tissue Inflammation in Diabetic versus Non-diabetic Pregnancies" (2014). *UMass Center for Clinical and Translational Science Research Retreat*. 69.
https://escholarship.umassmed.edu/cts_retreat/2014/posters/69

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.

Presenter Information

Tiffany A. Moore Simas, Raziell Rojas-Rodriguez, Jacqueline Tessa Draper, Aylin Sert, Maxwell Kruse, Crina Boeras, Brittany Rosenthal, Olga Gaelikman, So-Yun Min, Katherine G. Leung, and Silvia Corvera

Comments

Abstract of poster presented at the 2014 UMass Center for Clinical and Translational Science Research Retreat, held on May 20, 2014 at the University of Massachusetts Medical School, Worcester, Mass.

Jacqueline Draper participated in this study as a medical student in the Senior Scholars research program at the University of Massachusetts Medical School.

Creative Commons License

This work is licensed under a [Creative Commons Attribution-Noncommercial-Share Alike 3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/).

Adipose Tissue Inflammation in Diabetic versus Non-diabetic Pregnancies

Tiffany A. Moore Simas, MD, MPH, MEd
UMass Medical School/UMMHC
508-334-6678
Tiffany.mooresimas@umassmemorial.org

Tiffany A. Moore Simas, MD, MPH, MEd^{1a}; Raziell Rojas-Rodriguez, BS^{1b}; Jacqueline Tessa Draper, BS¹; Aylin Sert, MEd¹; Maxwell Kruse^{1b}; Crina Boeras, MD^{1a}; Brittany Rosenthal^{1b}; Olga Gealekman, PhD^{1b}; So Yun Min, MSc^{1b}; Kathy Leung, MPH^{1a}; Silvia Corvera, MD^{1b}

¹University of Massachusetts Medical School; ^a Obstetrics & Gynecology, ^b Program in Molecular Medicine

Objective: Beyond weight associated with fetus, placenta, amniotic fluid and increased blood volume, adipose tissue (AT) expansion is an accepted and expected component of pregnancy weight gain. Normal pregnancy is associated with relative insulin resistance (IR). In non-pregnant humans, AT expansion has been associated with IR and AT inflammation. However, it is not known whether AT expansion and IR in pregnancy are also associated with AT inflammation. This pilot study examined relationships between AT expansion and inflammation in control versus diabetic pregnancies.

Methods: Eligible subjects undergoing scheduled Cesarean delivery for obstetric indications were prospectively enrolled. Subjects provided demographic and anthropometric data, and biologic specimens. Immunofluorescence microscopy was performed on subcutaneous (SQ) and omental (OM) AT samples to evaluate macrophage infiltration. Included gravidas had paired AT samples and either negative glucola screening (controls) or gestational or pre-gestational Type 2 diabetes mellitus (T2DM).

Results: 13 subjects with SQ and OM AT samples were evaluated (10-controls, 3-diabetics (2-T2DM and 1-GDM)). Mean BMI and gestational weight gain of controls was 27.8 kg/m² (range 19.5-42) and 27.6 pounds (range 15-36) and of diabetics was 30.6 kg/m² (range 30-33) and 19 pounds (range -3-30), respectively. Macrophage infiltration was seen in OM AT from 2/ 3 diabetics and 0/ 10 controls (see figure).

Conclusions: These results indicate that AT expansion in non-diabetic pregnancies is not accompanied by macrophage infiltration. Thus, the IR of normal pregnancy is unlikely to be related to AT inflammation, and AT expansion per se does not lead to AT inflammation. However, as has been reported for T2DM in non-pregnant humans, the presence (T2DM) or development (GDM) of diabetes in pregnancy is associated with macrophage infiltration of AT. Despite the small sample size, the observed large differences in macrophage infiltration between controls and diabetics suggest that these findings will persist in a larger cohort.

