

UMass Chan Medical School

eScholarship@UMassChan

---

UMass Center for Clinical and Translational  
Science Research Retreat

2014 UMass Center for Clinical and  
Translational Science Research Retreat

---

May 20th, 12:30 PM

## Evening Snacking in Relation to Self-reported Declines in Sleep Quality during Pregnancy: Preliminary Results from the Decision-Making, Eating, and Weight Gain During Pregnancy (DEW) Study

Wendy McCallum

*University of Massachusetts Medical School*

*Et al.*

Let us know how access to this document benefits you.

Follow this and additional works at: [https://escholarship.umassmed.edu/cts\\_retreat](https://escholarship.umassmed.edu/cts_retreat)



Part of the [Behavior and Behavior Mechanisms Commons](#), [Dietetics and Clinical Nutrition Commons](#), [Obstetrics and Gynecology Commons](#), [Translational Medical Research Commons](#), and the [Women's Health Commons](#)

---

### Repository Citation

McCallum W, Appelhans BM, Moore Simas TA, Xiao R, Alikhan S, Pierre-Louis IC, Olendzki BC, Pagoto SL, Waring ME. (2014). Evening Snacking in Relation to Self-reported Declines in Sleep Quality during Pregnancy: Preliminary Results from the Decision-Making, Eating, and Weight Gain During Pregnancy (DEW) Study. UMass Center for Clinical and Translational Science Research Retreat. <https://doi.org/10.13028/dx30-2q40>. Retrieved from [https://escholarship.umassmed.edu/cts\\_retreat/2014/posters/98](https://escholarship.umassmed.edu/cts_retreat/2014/posters/98)

Creative Commons License



This work is licensed under a [Creative Commons Attribution-NonCommercial-Share Alike 3.0 License](#).

This material is brought to you by eScholarship@UMassChan. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMassChan. For more information, please contact [Lisa.Palmer@umassmed.edu](mailto:Lisa.Palmer@umassmed.edu).

## **Evening snacking in relation to self-reported declines in sleep quality during pregnancy: preliminary results from the Decision-Making, Eating, and Weight Gain during pregnancy (DEW) Study**

Wendy McCallum, BA<sup>1</sup>, Bradley M. Appelhans, PhD<sup>2</sup>, Tiffany A. Moore Simas, MD MPH MEd<sup>3,4,5</sup>, Sherry R. Xiao, MPH<sup>1</sup>, Shums Alikhan, BS<sup>1</sup>, Isabelle C. Pierre-Louis, BS<sup>1</sup>, Barbara C. Olendzki, RD MPH LDN<sup>6</sup>, Sherry L. Pagoto, PhD<sup>6</sup>, Molly E. Waring, PhD<sup>1</sup>

<sup>1</sup> Division of Epidemiology of Chronic Diseases and Vulnerable Populations, Department of Quantitative Health Sciences, University of Massachusetts Medical School

<sup>2</sup> Department of Preventive Medicine, Rush University Medical Center, Chicago, IL

<sup>3</sup> Division of Research, Department of Obstetrics and Gynecology and Department of Pediatrics, University of Massachusetts Medical School

<sup>4</sup> Department of Pediatrics, University of Massachusetts Medical School

<sup>5</sup> Department of Obstetrics and Gynecology, UMass Memorial Health Care

<sup>6</sup> Division of Behavioral and Preventive Medicine, Department of Medicine, University of Massachusetts Medical School

Contact info: Molly E. Waring, PhD; Department of Quantitative Health Sciences, University of Massachusetts Medical School, 368 Plantation Street, AS8-1077, Worcester, MA, 01605; email [molly.waring@umassmed.edu](mailto:molly.waring@umassmed.edu); phone (508) 856-3504

**Background:** Poor sleep in non-pregnant adults has been associated with increased evening snacking, which may contribute to weight gain. Sleep disturbances are common during pregnancy.

**Objective:** To examine the association between changes in sleep quality from pre-pregnancy and evening snacking.

**Methods:** In an ongoing prospective cohort study, pregnant women were recruited from UMMHC obstetric practices and the community. Participants are 18+ years, with singleton gestation <36 weeks, pre-pregnancy BMI 18.5-40 kg/m<sup>2</sup>, English-speaking, and with plans to deliver at UMMHC. Participants were asked “compared to the three months before you became pregnant, how is your sleep quality now?”; we combined responses of “about the same”/“a little better”/“a lot better” versus “a little worse”/“much worse”. Participants completed three 24-hour dietary recalls (2 weekdays, 1 weekend day). Evening snacks were defined as eating occasions after dinner but before bedtime during which food items other than water was consumed. Fisher’s Exact tests and t-tests provided comparisons for evening snacking (yes/no), number of snacks, and energy intake.

**Results:** Women with complete data (n=55) were 58% non-Hispanic White and aged 30.0 (SD:4.3) years; gestational age at study visit was 23.0 (SD:5.9) weeks. Of 866 meals reported, 94 were evening snacks. 71% (n=39) reported that their current sleep quality was worse than before pregnancy. Evening snacks were reported by 90% of women reporting worse sleep and 69% same/better (p=0.1028). While the number of snacks among snackers did not differ by change in sleep quality (M[SD]: 2.2[1.2] versus 1.6[0.8], p=0.2372), energy intake from these snacks was higher among women whose sleep quality had declined (M[SD]: 630[488] versus 309[331] kcal, p=0.0480).

**Conclusions:** Declines in sleep quality during pregnancy may be linked to evening snacking. More research is needed to understand the role of sleep quality, eating behavior, and weight gain during pregnancy.