

May 20th, 12:30 PM

Contemporary Analysis of Malignancies in Women of Child-Bearing Age: An NSQIP Analysis

Eva Rouanet

University of Massachusetts Medical School

Ann-Kristin U. Friedrich

University of Massachusetts Medical School

Kate Dinh

University of Massachusetts Medical School

See next page for additional authors

Follow this and additional works at: https://escholarship.umassmed.edu/cts_retreat

Part of the [Neoplasms Commons](#), [Oncology Commons](#), [Surgery Commons](#), and the [Women's Health Commons](#)

Rouanet, Eva; Friedrich, Ann-Kristin U.; Dinh, Kate; Baratta, Kevin P.; Whalen, Giles F.; Santry, Heena; and LaFemina, Jennifer, "Contemporary Analysis of Malignancies in Women of Child-Bearing Age: An NSQIP Analysis" (2016). *UMass Center for Clinical and Translational Science Research Retreat*. 71.

https://escholarship.umassmed.edu/cts_retreat/2016/posters/71

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.

Presenter Information

Eva Rouanet, Ann-Kristin U. Friedrich, Kate Dinh, Kevin P. Baratta, Giles F. Whalen, Heena Santry, and Jennifer LaFemina

Keywords

cancer, pregnancy, nonpregnany, malignancies

Creative Commons License

Creative

Commons

This work is licensed under a [Creative Commons Attribution-Noncommercial-Share Alike 3.0 License](#).

Attribution-

Noncommercial-

Share

Alike

3.0

License

Contemporary Analysis of Malignancies in Women of Child-Bearing Age: An NSQIP Analysis

Eva Rouanet¹ MSII, Ann-Kristin Friedrich, MD¹, Kate Dinh, MD¹, Kevin P. Baratta, MD MPH¹, Giles F. Whalen, MD FACS¹, Heena Santry MD FACS¹, Jennifer LaFemina MD FACS¹

¹University of Massachusetts Medical School

Background: Recent evidence suggests that cancer incidence among pregnant women is increasing. The pattern of malignancies in pregnant women and how these compare to their nonpregnant counterparts has not been explored. Here we describe the differences in the proportion of resected malignancies in this population.

Methods: The American College of Surgeons National Surgical Quality Improvement Program (ACS-NSQIP) database was used to identify women aged 18-49 who underwent an operation for malignancy from 2007-2012. Age-adjusted distribution of specific surgical interventions for malignancy based on ICD-9 codes were compared among pregnant and non-pregnant women using logistic regression analysis.

Results: 42,732 subjects with malignancies surgically treated during child-bearing age were identified. 0.33% (n=143) were pregnant. The most common tumors requiring resection were breast (51%), thyroid (17%), and colorectal (9%). The distribution for most cancers was similar between groups. The age-adjusted proportion was significantly increased in breast, major salivary gland and oropharyngeal malignancies (p<0.05). The proportion of resected colorectal cancers was significantly lower in pregnant women (p<0.05; Table 1).

Conclusion: This study serves as the first comprehensive and contemporary overview of malignancies resected in women of childbearing age. This study demonstrates that the proportion of resections among pregnant women was significantly greater in breast, major salivary gland and oropharyngeal cancers and lower for colorectal cancers. While these data might represent true differences in cancer incidence, further work is necessary to demonstrate if these are true differences in incidence versus differences in detection and treatment of the pregnant patient.

Contact:

Eva Rouanet

M.D. Candidate, Class of 2018

University of Massachusetts Medical School

Eva.rouanet@umassmed.edu