

10-1-2016

Dignity in mental health practice and research: Time to unite on innovation, outreach and education

Douglas M. Ziedonis

University of Massachusetts Medical School

Celine Larkin

University of Massachusetts Medical School

Raghu Appasani

University of Massachusetts Medical School

Follow this and additional works at: https://escholarship.umassmed.edu/psych_pp

 Part of the [Mental and Social Health Commons](#), [Psychiatry Commons](#), and the [Psychiatry and Psychology Commons](#)



This work is licensed under a [Creative Commons Attribution-Noncommercial-Share Alike 3.0 License](#).

Repository Citation

Ziedonis, Douglas M.; Larkin, Celine; and Appasani, Raghu, "Dignity in mental health practice and research: Time to unite on innovation, outreach and education" (2016). *Psychiatry Publications and Presentations*. 784.

https://escholarship.umassmed.edu/psych_pp/784

Indian J Med Res. 2016 Oct; 144(4): 491–495.

PMCID: PMC5345292

doi: [10.4103/0971-5916.200885](https://doi.org/10.4103/0971-5916.200885)

Dignity in mental health practice & research: Time to unite on innovation, outreach & education

[Douglas Ziedonis](#),* [Celine Larkin](#), and [Raghu Appasani](#)

Department of Psychiatry, University of Massachusetts Medical School, Worcester, MA, USA

*For correspondence : Douglas.ziedonis@umassmemorial.org

Received 2016 Sep 12

Copyright : © 2017 Indian Journal of Medical Research

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

Dignity has been described as ‘the inherent and inalienable worth of all human beings irrespective of social status such as race, gender, physical or mental state’¹. Dignity is, therefore, at the core of psychological well-being, social connection and humanity. Mental health interventions that explicitly promote and preserve dignity for people experiencing mental distress are growing in clinical practice and research in the USA, India, Europe and elsewhere². However, there is a need for more research and policy supporting the implementation and evaluation of these initiatives. Here, we highlight some of these programmes and make recommendations on how to further integrate dignity in mental health research, practice, outreach and education in India and the world.

Advocating for dignity in the context of mental distress and isolation requires a redoubling of effort to engage a person who may feel disconnected, to communicate even when a person seems uncomprehending and to connect when a person's perceptions and understanding are different from one's own. Dignity should be afforded to all, and evidence is accruing that recovery-oriented support and mental health first aid in the community and in treatment settings make a difference. These efforts not only create a positive experience for the person at the centre of concern, the family and support network, clinicians and the public but also lead to real dialogue, person-centred care, reduced stigma and broader social and recovery capital.

Protecting and promoting dignity plays a central role in the UNESCO Universal Declaration on Bioethics and Human Rights³. This goal must include strategies that support psychological and mental health first aid for all, including awareness of issues faced by people experiencing mental distress and isolation⁴. Dignity is fundamentally a social construct and people require a sense of connection in their society to create meaning, purpose and self-worth. Social isolation and loneliness are often at the core of mental distress, and interventions to increase a person's sense of dignity and belonging buffer against psychological and social stressors that otherwise precipitate or worsen mental distress⁵. Conversely, prejudice, stigma, discrimination and disrespect can thwart a person's sense of belonging and hinder help-seeking from their social network, community or clinical services during times of mental distress⁶. To overcome these attitudinal barriers, interventions need to be implemented flexibly and sustainably such as using gatekeepers, peers, train-the-trainer models, cell phone- and internet-based strategies, community pairing partnerships and other low-cost community-based approaches.

Dignity in global mental health

The challenges and solutions associated with promoting dignity in mental health can vary as a function of culture and geography. For example, rural mental health has become a growing concern throughout the world, especially given the low provision of mental health services in these areas. India, like many countries, is a complex nation of contrasts where geographical barriers to mental health care can be seen in sharp focus. Seventy per cent of India's population lives in rural areas, where there are higher rates of affective disorders, particularly among the elderly⁷ and about four psychiatrists serve every one million Indian citizens, tending to work in urban areas⁸. Due to the lack of resources, stigma and a shortage of education around mental health, people in rural areas remain isolated and at risk for suicide⁹. Such barriers need innovative solutions that incorporate recovery-orientated approaches involving primary care practitioners, caregivers, community members and people with lived experience of mental health problems. In rural areas, there is an additional need to develop a workforce that has the skills to provide evidence-based approaches in a manner that supports dignity, engagement and empowerment.

Psychological and mental health first aid for all

How can we better promote and protect dignity in mental health for all? How can we encourage more dialogue and support in day-to-day clinical practice? How do we encourage tolerance and curiosity to foster lasting connections? As a starting point, humble inquiry increases clinicians' listening abilities and sense of connection and dignity. Mindful communication can also support deep listening with dignity while building personal resilience and compassion for self and others. Through meditation and yoga practices, individuals increase self-awareness to their immediate experience in a purposeful manner (*i.e.*, to their breath, body sensation and feelings) with curiosity and without judgement¹⁰. These skills and way of being are enhanced through formal and informal practices that increase self-awareness and emotional intelligence¹¹.

Specific programmes such as Mental Health First Aid, Emotional CPR, Alcoholics Anonymous (and other 12-Step Fellowships) and Open Dialogue can serve to expand psychological support in ways that enhance dignity and respect for all. Mental Health First Aid is a widely implemented and evidence-based approach that is applied at a community level. Through brief gatekeeper training, members of the general public are educated about mental health and equipped with skills to intervene with those who are distressed or suicidal. A systematic review¹² demonstrated a large effect of the intervention on knowledge and medium-sized effects on both attitudes and help-providing behaviours. Starting with the Mental Health First Aid model as a template, Delphi consensus studies in India¹³, as well as Japan¹⁴ and the Philippines¹⁵, succeeded in generating culturally tailored guidelines of actions for members of the public to use when they encounter a suicidal person. Rural and urban areas in the USA are implementing Mental Health First Aid, and smaller Eastern European countries such as Latvia are using videos of individuals with lived experience of depression, schizophrenia and dementia in an effort to reduce stigma and isolation¹⁶.

There are numerous other examples of programmes designed to reduce stigma which involve people with lived experience. The voices of people with lived experience can mental health services to be more aware and supportive of dignity in mental health. The innovative Emotional CPR intervention, which was developed by an international team of persons with lived experience, teaches specific principles for people to embody when helping someone in emotional distress including full listening, authenticity, non-judging attitude, making connection and empowerment as a means of restoring dignity¹⁷. Twelve-Step Fellowship programmes¹⁸ and Hearing Voices¹⁹ are other examples of peer support that foster hope and recovery.

Open Dialogue prioritizes respect and dignity for people experiencing severe mental distress and seeks to understand the person and his/her family in the broader personal, social and cultural context²⁰. This recovery-oriented approach quickly brings together the person at the centre of concern with the family or support network and multiple clinicians. In network meetings, clinicians engage in natural conversation in an

attempt to understand the multiple perspectives of the current situation by listening to people's stories (over their symptoms) and creating space for all to be heard. The approach is fully transparent, with shared treatment planning discussions occurring only in the presence of the individual at the centre of concern. The individual and support network engage in a dialogue which includes responding to reflections that clinicians express amongst themselves. Initial studies have shown promising results in preventing relapse, improving quality of life, as well as increased satisfaction for patients, families and clinicians^{21,22}.

Having multiple mental health problems such as co-occurring mental illness and addiction (co-occurring disorders) can worsen stigma²³ and requires integrated holistic care that provides hope and empowerment²⁴. Co-occurring disorders become more complex when these occur in the context of homelessness and recent incarceration. Homelessness and incarceration experiences in themselves often involve the degradation of dignity through the loss of privacy and autonomy. The MISSION model targets homeless veterans, often with co-occurring disorders, in the USA through a network of clinicians and peer support specialists in a manner that promotes dignity and has resulted in improved treatment engagement and outcomes²⁵.

There are numerous other community-based recovery-oriented interventions that promote dignity and social integration. The Clubhouse model, implemented in 33 countries, provides integrated therapeutic working communities for people with severe mental health problems and members work (in transitional, supported or independent employment) in an approach that emphasizes individuals' strengths and relationships. A recent systematic review found that the intervention was effective in promoting employment, reducing hospitalizations and improving quality of life²⁶. The broadest application of a community-based approach can be seen in the phenomenon of 'therapeutic towns' such as Geel (Belgium)²⁷ and Sopo (Colombia)²⁸. These towns serve to create a 'foster community' where those with mental health problems are integrated into community and working life.

In rural India, the MINDS Foundation operates in Gujarat using a grassroot approach which implements screening and intervention for mental illness through health worker training, community education and engagement and direct service delivery. Through assessment of the communities' attitudes and knowledge of mental health, MINDS has been able to formulate targeted mental health education programming and successfully train community health workers in identifying mental health symptoms, providing initial screening and being a resource for continued holistic mental health care²⁹. This model has reduced stigma and promoted help-seeking by identifying those in need and connecting them in a respectful and dignified manner to appropriate affordable care.

Recommendations and conclusions

There are many additional opportunities to further develop, implement and evaluate psychological and mental health first aid approaches that support dignity in mental health. However, policymakers and healthcare leaders must prioritize and support this area of work. There is a need for people with lived experience, researchers and clinicians to work together and further develop and study interventions that integrate dignity as a core principle of therapeutic practice. Engaging a broader group of stakeholders would include families, community leaders and educators. Ethical frameworks already aim to protect dignity in research by emphasizing respect for autonomy, beneficence, non-maleficence and justice in research procedures³⁰. To take this aim one step further in mental health research, dignity could be integrated as a measurable process and outcome in intervention development and evaluation, through the adaptation of existing psychometric instruments or through linguistic and behavioural observation coding. Such approaches have previously been applied with constructs of recovery and therapeutic alliance.

Research on how clinicians and people with lived experience engage and connect with others through web-based and mobile mental health interventions also requires further study. There are several well-established web-based interventions for mild depression such as MoodGym³¹ and iFightDepression³², operating in Western countries that could be tailored for Asian settings as needed. In rural India, the MINDS Foundation

collaborates with Medic Mobile, Inc., to deliver a simple 22-question survey through an SMS mobile device in an effort to better identify those in need of mental health care. These approaches reduce barriers of cost, geography and limited trained professionals³³ and promote self-management and patient empowerment that can enhance dignity.

Improved integration of mental health care into primary care will further improve mental health outcomes for all, especially in the context of a supportive and respectful primary care clinician-patient alliance. Primary care programmes can benefit from using change management strategies from implementation science or quality improvement approaches in order to change their treatment culture and integrate mental health evidence based approaches. The aim of dignity and respect for all should be explicitly identified in health providers' mission statements and other policies. More importantly, these should be upheld in daily communication and interactions at all levels. These concepts (and their associated skills such as respectful communication, deep listening and patience) also need to be enshrined in curricula for clinical training and maintained through continuing education.

In broader society, politicians, media, schools and communities all have a role to play in tackling mental health-related stigma and promoting dignity through education, policies and action. The isolation of mental distress can be a very painful experience. With societal and organizational support, dignity and respect can help bring healing and recovery one step closer for all.

Footnotes

This editorial is published on the occasion of World Mental Health Day – October 10, 2016

References

1. Funk M, Drew N, Baudel M. Dignity in mental health. Occoquan, VA: World Federation for Mental Health; 2015. Supporting dignity through mental health legislation; pp. 14–5.
2. Rudnick A. Recovery of people with mental illness: philosophical and related perspectives. Oxford: Oxford University Press; 2012.
3. ten Have HAMJ, Jean MS, editors. The UNESCO Universal declaration on bioethics and human rights: Background, principles and application. Paris: UNESCO; 2009.
4. Saxena S, Hanna F. Dignity – A fundamental principle of mental health care. *Indian J Med Res.* 2015;142:355–8. [PMCID: PMC4683817] [PubMed: 26609024]
5. Thoits PA. Mechanisms linking social ties and support to physical and mental health. *J Health Soc Behav.* 2011;52:145–61. [PubMed: 21673143]
6. Clement S, Schauman O, Graham T, Maggioni F, Evans-Lacko S, Bezborodovs N, et al. What is the impact of mental health-related stigma on help-seeking? A systematic review of quantitative and qualitative studies. *Psychol Med.* 2015;45:11–27. [PubMed: 24569086]
7. Swarnalatha N. The prevalence of depression among the rural elderly in Chittoor district, Andhra Pradesh. *J Clin Diagn Res.* 2013;7:1356–60. [PMCID: PMC3749635] [PubMed: 23998065]
8. Khandelwal SK, Jhingan HP, Ramesh S, Gupta RK, Srivastava VK. India mental health country profile. *Int Rev Psychiatry.* 2004;16:126–41. [PubMed: 15276945]
9. Radhakrishnan R, Andrade C. Suicide: An Indian perspective. *Ind J Psychiatry.* 2012;54:304–19. [PMCID: PMC3554961]
10. Kabat-Zinn J. Full catastrophe living. New York: Bantam Dell; 1990.

11. Good DJ, Lyddy CJ, Glomb TM. Contemplating mindfulness at work: an integrative review. *J Manag.* 2016;42:114–42.
12. Hadlaczky G, Hökby S, Mkrtchian A, Carli V, Wasserman D. Mental Health First Aid is an effective public health intervention for improving knowledge, attitudes, and behaviour: a meta-analysis. *Int Rev Psychiatry.* 2014;26:467–75. [PubMed: 25137113]
13. Colucci E, Kelly CM, Minas H, Jorm AF, Chatterjee S. Mental Health First Aid guidelines for helping a suicidal person: a Delphi consensus study in India. *Int J Ment Health Syst.* 2010;4:4. [PMCID: PMC2847966] [PubMed: 20167125]
14. Colucci E, Kelly CM, Minas H, Jorm AF, Suzuki Y. Mental Health First Aid guidelines for helping a suicidal person: A Delphi consensus study in Japan. *Int J Ment Health Syst.* 2011;5:12. [PMCID: PMC3108941] [PubMed: 21592409]
15. Colucci E, Kelly CM, Minas H, Jorm AF, Nadera D. Mental Health First Aid guidelines for helping a suicidal person: A Delphi consensus study in the Philippines. *Int J Ment Health Syst.* 2010;4:32. [PMCID: PMC3017011] [PubMed: 21167076]
16. Samele C, Frew S, Urquia N. Brussels: European Commission; 2013. [accessed on November 11, 2016]. Mental Health Systems in the European Union Member States, Status of Mental Health in Populations and Benefits to be Expected from Investments into Mental Health (EuroPoPP-MH). European profile of prevention and promotion of mental health (EuroPoPP-MH) Available from: http://ec.europa.eu/health/mental_health/docs/europopp_full_en.pdf.
17. Fisher D, McCarthy D, Sweeney J. The value of using dialogue and self in recovery. In: Jones JS, Fitzpatrick JJ, Rogers VL, editors. *Psychiatric-mental health nursing: an interpersonal approach.* 2nd ed. New York, N.Y: Springer Publishing Company; 2016. pp. 31–47.
18. Schenker MD. *A clinician's guide to 12-step recovery: Integrating 12-step programs into psychotherapy.* New York: W.W Norton & Company, Inc; 2009.
19. Ruddle A, Mason O, Wykes T. A review of hearing voices groups: Evidence and mechanisms of change. *Clin Psychol Rev.* 2011;31:757–66. [PubMed: 21510914]
20. Olson M, Seikkula J, Ziedonis D. *The key elements of dialogic practice in open dialogue.* Worcester, MA: University of Massachusetts Medical School; 2014.
21. Gordon C, Gidugu V, Rogers ES, DeRonck J, Ziedonis D. Adapting open dialogue for early-onset psychosis into the U.S. health care environment: a feasibility study. *Psychiatr Serv.* 2016;67:1166–8. [PubMed: 27417900]
22. Seikkula J, Aaltonen J, Alakare B, Haarakangas K, Keränen J, Lehtinen K. Five-year experience of first-episode nonaffective psychosis in open-dialogue approach: treatment principles, follow-up outcomes, and two case studies. *Psychother Res.* 2006;16:214–28.
23. Glass JE, Williams EC, Bucholz KK. Psychiatric comorbidity and perceived alcohol stigma in a nationally representative sample of individuals with DSM-5 alcohol use disorder. *Alcohol Clin Exp Res.* 2014;38:1697–705. [PMCID: PMC4059348] [PubMed: 24848495]
24. Ziedonis DM. Integrated treatment of co-occurring mental illness and addiction: clinical intervention, program, and system perspectives. *CNS Spectr.* 2004;9:892–904, 925. [PubMed: 15618940]
25. Smelson D, Kalman D, Losonczy MF, Kline A, Sambamoorthi U, Hill LS, et al. A brief treatment engagement intervention for individuals with co-occurring mental illness and substance use disorders: results of a randomized clinical trial. *Community Ment Health J.* 2012;48:127–32. [PubMed: 20859765]

26. McKay C, Nugent KL, Johnsen M, Eaton WW, Lidz CW. A systematic review of evidence for the clubhouse model of psychosocial rehabilitation. *Adm Policy Ment Health*. 2016 In press.
27. van Bilsen HP. Lessons to be learned from the oldest community psychiatric service in the world: Geel in Belgium. *BJPsych Bull*. 2015;40:207–11. [PMCID: PMC4967781]
28. Rodríguez-Frías E. Challenging assumptions: Psychiatric disabilities and grassroots development. *Grassroots Dev*. 2010;31:36–41.
29. Schoonover J, Lipkin S, Javid M, Rosen A, Solanki M, Shah S, et al. Perceptions of traditional healing for mental illness in rural Gujarat. *Ann Glob Health*. 2014;80:96–102. [PubMed: 24976546]
30. Gillon R. Medical ethics: Four principles plus attention to scope. *BMJ*. 1994;309:184–8. [PMCID: PMC2540719] [PubMed: 8044100]
31. Twomey C, O'Reilly G. Effectiveness of a freely available computerised cognitive behavioural therapy programme (MoodGYM) for depression: meta-analysis. *Aust N Z J Psychiatry*. 2016 In press.
32. Arensman E, Koburger N, Larkin C, Karwig G, Coffey C, Maxwell M, et al. Depression awareness and self-management through the internet: protocol for an internationally standardized approach. *JMIR Res Protoc*. 2015;4:e99. [PMCID: PMC4705028] [PubMed: 26251104]
33. Yellowlees P, Chan S. Mobile mental health care – An opportunity for India. *Indian J Med Res*. 2015;142:359–61. [PMCID: PMC4683818] [PubMed: 26609025]

Articles from The Indian Journal of Medical Research are provided here courtesy of **Medknow Publications**