

## EDITORIAL

# Skin Cancer Risk in Gay and Bisexual Men

## A Call to Action

Aaron J. Blashill, PhD; Sherry Pagoto, PhD

**In this issue of *JAMA Dermatology*,** Mansh and colleagues<sup>1</sup> report results from the first known study to examine the prevalence of skin cancer in sexual minorities. The authors leveraged large US-representative and California-representative epidemiological databases to explore self-reported diagnoses of skin cancer and use of indoor tanning among nearly



Related article [page 1308](#)

200 000 adults. The findings revealed significant sexual orientation health disparities. Skin cancer rates among sexual minority men ranged between 4.3% and 6.7%, whereas the rates among heterosexual men ranged between 2.7% and 3.2%, for increased adjusted odds of skin cancer for sexual minority men between 1.56 and 2.13. Nonsignificant differences emerged between heterosexual and sexual minority women. Although Mansh and colleagues<sup>1</sup> did not statistically compare the rates of skin cancer between sexual minority men and women, it is worth noting that sexual minority men reported the highest rates across sex and sexual orientation.

In addition to sexual orientation disparities in skin cancer diagnoses, Mansh and colleagues<sup>1</sup> reported substantial disparities in a key skin cancer risk behavior: indoor tanning. Sexual minority men reported a 12-month prevalence of any indoor tanning between 5.1% and 7.4% compared with 1.5% to 1.6% among heterosexual men, 2.6% to 4.1% among sexual minority women, and 5.0% to 6.5% among heterosexual women. Indoor tanning is a likely factor in the elevated rate of skin cancer in sexual minority men<sup>2</sup> since the International Agency for Research on Cancer<sup>3</sup> assigned indoor tanning as a group 1 carcinogen in 2009.

Previous research<sup>4</sup> on indoor tanning has reported that women are the population engaging in the highest rates and are most in need of intervention. As such, research focused on reducing indoor tanning has been exclusively focused on women.<sup>5</sup> To our knowledge, no trial on interventions for reducing indoor tanning has targeted sexual minority men, and no evidence exists to show that interventions targeting women would resonate with sexual minority men. Public health campaigns for skin cancer prevention, often informed by intervention and health communication research, have been instrumental in raising awareness about skin cancer. To the extent that sexual minority groups continue to be neglected in skin cancer prevention research, they will remain untargeted in public health campaigns, further facilitating this health disparity.

The findings reported by Mansh and colleagues<sup>1</sup> add to the growing list of sexual orientation health disparities that have

been revealed when sexual orientation has been measured.<sup>6</sup> Sexual minority men are at increased risk for developing not only skin cancer but also other forms of cancer,<sup>7</sup> human immunodeficiency virus,<sup>8</sup> substance use disorders,<sup>9</sup> and mental illness.<sup>10</sup> A prevailing theory on why sexual minority men are disproportionately affected by physical and mental health problems is the sexual minority stress model,<sup>11</sup> which suggests that stigma, prejudice, discrimination, and victimization create a hostile social environment for sexual minorities, which subsequently leads to elevated psychological distress. Elevated distress created by macrolevel social forces leads to health risk behaviors, including condomless sex, tobacco use, physical inactivity, and, possibly, indoor tanning. Supporting the role of psychological distress are studies showing that, among women, stress and depression are disproportionately elevated among indoor tanners.<sup>12</sup>

Indoor tanning has also been strongly associated with motives to enhance physical appearance.<sup>13</sup> Sexual minority men report elevated body dissatisfaction compared with heterosexual men, with levels of body image concerns similar to those of heterosexual women.<sup>14</sup> The convergence of elevated stress and body dissatisfaction may make sexual minority men particularly vulnerable to tanning as a way to cope with these issues. Unfortunately, to our knowledge, no published studies have examined predictors of indoor tanning among sexual minority men. The development of theoretically and empirically based skin cancer prevention programs will require first identifying drivers of elevated skin cancer risk in this population.

The results from Mansh and colleagues<sup>1</sup> have implications for both clinical care and research. Assessing sexual orientation as part of the initial patient interview in clinical dermatology settings could identify men at elevated risk for skin cancer and possible indoor tanners. Many physicians do not assess patient sexual orientation as part of routine care<sup>15</sup>; however, when sexual minority patients are informed why this information is important to their care and that it will be kept confidential, many are open to sharing it with their healthcare professionals. Recent data<sup>16</sup> have highlighted significant explicit and implicit negative attitudes toward sexual minorities by physicians and other health care professionals, suggesting that additional training in sexual minority issues is needed in medical school curricula. The expression of explicit or implicit bias toward sexual minorities is destructive to a physician-patient relationship and likely a liability to care. In addition, the Institute of Medicine<sup>6</sup> and Healthy People 2020 have called for routine assessment of sexual orientation in

medical care and in all federally funded epidemiological surveys. The California Health Interview Surveys, the Adult National Health Interview Survey, and the National Longitudinal Study of Adolescent to Adult Health (Add Health) are 3 examples of epidemiological surveys including these data.

Skin cancer risk reduction in sexual minority men is an area in urgent need of research. Studies exploring predictors of indoor tanning and other skin cancer risk behaviors in sexual minority men and how these fit into the sexual minority stress model would further our understanding of why sexual minority men are disproportionately affected by preventable dis-

eases. Further research is also needed to determine where sexual minority men engage in indoor tanning (ie, salon vs non-salon locations) and to describe the nature of their tanning habits. Finally, skin cancer prevention interventions are also needed to reduce risk behavior in this population. Physicians can improve care and lend insights about this population by assessing sexual orientation as part of routine care and entering it into electronic medical records. Ultimately, reducing health disparities in sexual minorities will also require recognizing and eliminating bias and stigma in both research and clinical practice.

#### ARTICLE INFORMATION

**Author Affiliations:** Department of Psychology, San Diego State University, San Diego, California (Blashill); Joint Doctoral Program in Clinical Psychology, San Diego State University/University of California, San Diego (Blashill); Division of Prevention and Behavioral Medicine, Department of Medicine, University of Massachusetts Medical School, Worcester (Pagoto).

**Corresponding Author:** Aaron J. Blashill, PhD, Department of Psychology, San Diego State University, 6363 Alvarado Ct, Ste 103, San Diego, CA 92120 (aaron.blashill@sdsu.edu).

**Published Online:** October 7, 2015.  
doi:10.1001/jamadermatol.2015.3125.

**Funding/Support:** Some investigator time was supported by grant K23MH096647 from the National Institute of Mental Health (Dr Blashill).

**Role of the Funder/Sponsor:** The funding organization had no role in the design and conduct of the work; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

#### REFERENCES

1. Mansh M, Katz KA, Linos E, Chren M-M, Arron S. Association of skin cancer and indoor tanning in sexual minority men and women [published online October 7, 2015]. *JAMA Dermatol*. doi:10.1001/jamadermatol.2015.3126.
2. Blashill AJ, Safren SA. Skin cancer risk behaviors among US men: the role of sexual orientation. *Am J Public Health*. 2014;104(9):1640-1641.
3. El Ghissassi F, Baan R, Straif K, et al; WHO International Agency for Research on Cancer Monograph Working Group. A review of human carcinogens, part D: radiation. *Lancet Oncol*. 2009;10(8):751-752.
4. Guy GP Jr, Tai E, Richardson LC. Use of indoor tanning devices by high school students in the United States, 2009. *Prev Chronic Dis*. 2011;8(5):A116.
5. Heckman CJ, Manne SL, eds. *Shedding Light on Indoor Tanning*. Dordrecht, the Netherlands: Springer; 2012.
6. Institute of Medicine (IOM). *The Health of Lesbian, Gay, Bisexual, and Transgender People: Building a Foundation for Better Understanding*. Washington, DC: National Academies Press; 2011. [http://books.nap.edu/openbook.php?record\\_id=13128&page=R2](http://books.nap.edu/openbook.php?record_id=13128&page=R2). Accessed July 23, 2015.
7. Boehmer U, Miao X, Ozonoff A. Cancer survivorship and sexual orientation. *Cancer*. 2011;117(16):3796-3804.
8. Cochran SD, Mays VM. Physical health complaints among lesbians, gay men, and bisexual and homosexually experienced heterosexual individuals: results from the California Quality of Life Survey. *Am J Public Health*. 2007;97(11):2048-2055.
9. McCabe SE, Hughes TL, Bostwick WB, West BT, Boyd CJ. Sexual orientation, substance use behaviors and substance dependence in the United States. *Addiction*. 2009;104(8):1333-1345.
10. King M, Semlyen J, Tai SS, et al. A systematic review of mental disorder, suicide, and deliberate self harm in lesbian, gay and bisexual people. *BMC Psychiatry*. 2008;8(1):70.
11. Meyer IH. Prejudice, social stress, and mental health in lesbian, gay, and bisexual populations: conceptual issues and research evidence. *Psychol Bull*. 2003;129(5):674-697.
12. Hillhouse J, Stapleton J, Turrissi R. Association of frequent indoor UV tanning with seasonal affective disorder. *Arch Dermatol*. 2005;141(11):1465.
13. Cafri G, Thompson JK, Jacobsen PB, Hillhouse J. Investigating the role of appearance-based factors in predicting sunbathing and tanning salon use. *J Behav Med*. 2009;32(6):532-544.
14. Morrison MA, Morrison TG, Sager CL. Does body satisfaction differ between gay men and lesbian women and heterosexual men and women? a meta-analytic review. *Body Image*. 2004;1(2):127-138.
15. Wimberly YH, Hogben M, Moore-Ruffin J, Moore SE, Fry-Johnson Y. Sexual history-taking among primary care physicians. *J Natl Med Assoc*. 2006;98(12):1924-1929.
16. Sabin JA, Riskind RG, Nosek BA. Health care providers' implicit and explicit attitudes toward lesbian women and gay men. *Am J Public Health*. 2015;105(9):1831-1841.