

11-16-2012

The role of intraflagellar transport proteins in ciliary assembly and function

Gregory J. Pazour

University of Massachusetts Medical School

Follow this and additional works at: <http://escholarship.umassmed.edu/oapubs>



Part of the [Cell and Developmental Biology Commons](#)

Repository Citation

Pazour, Gregory J., "The role of intraflagellar transport proteins in ciliary assembly and function" (2012). *Open Access Articles*. 2452.
<http://escholarship.umassmed.edu/oapubs/2452>

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in Open Access Articles by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.

The role of intraflagellar transport proteins in ciliary assembly and function

GJ Pazour

From First International Cilia in Development and Disease Scientific Conference (2012)
London, UK. 16-18 May 2012

The sensory and motility functions of cilia play critical roles in the development of vertebrates and defects in these organelles lead to a wide range of structural birth defects. The intraflagellar transport (IFT) system is required for building all types of mammalian cilia. IFT particles are composed of about 20 proteins and these proteins are highly conserved across ciliated species. IFT25 and IFT27, however, are absent from certain ciliated organisms like *Caenorhabditis* and *Drosophila* suggesting that they may have a unique role distinct from ciliogenesis. We generated *Ift25* and *Ift27* null mice and show that these proteins are not required for ciliary assembly but are required for proper Hedgehog signaling, which in mammals occurs within cilia. Mutant mice die at birth with multiple phenotypes indicative of Hedgehog signaling dysfunction. Cilia lacking IFT25/27 have defects in the signal-dependent transport of multiple Hedgehog components including Patched-1, Smoothed, and Gli2 and fail to activate the pathway upon stimulation. These are the first examples of null IFT mutations that perturb Hedgehog signaling independent of ciliary architecture. Thus, IFT function is not restricted to assembling cilia where signaling occurs but also plays a direct role in signal transduction events.

Published: 16 November 2012

doi:10.1186/2046-2530-1-S1-O9

Cite this article as: Pazour: The role of intraflagellar transport proteins in ciliary assembly and function. *Cilia* 2012 **1**(Suppl 1):O9.

Correspondence: Gregory.Pazour@umassmed.edu
Program in Molecular Medicine, University of Massachusetts Medical School, USA

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

