

6-18-2018

Editorial: Controversies in the Local Management of Lung Cancer

John M. Varlotto

University of Massachusetts Medical School

Giulia Veronesi

Humanitas Clinical and Research Center

Follow this and additional works at: <https://escholarship.umassmed.edu/oapubs>

 Part of the [Neoplasms Commons](#), [Oncology Commons](#), [Respiratory Tract Diseases Commons](#), and the [Therapeutics Commons](#)



This work is licensed under a [Creative Commons Attribution 4.0 License](#).

Repository Citation

Varlotto, John M. and Veronesi, Giulia, "Editorial: Controversies in the Local Management of Lung Cancer" (2018). *Open Access Articles*. 3509.

<https://escholarship.umassmed.edu/oapubs/3509>

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in Open Access Articles by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.



Editorial: Controversies in the Local Management of Lung Cancer

John M. Varlotto^{1*} and Giulia Veronesi²

¹Department of Radiation Oncology, University of Massachusetts Medical Center, Worcester, MA, United States, ²Division of Thoracic Surgery, Humanitas Clinical and Research Center, Rozzano, Milan, Italy

Keywords: lung cancer, multi-modality treatment, thoracic surgery, radiation therapy, lung cancer treatment

Editorial on the Research Topic

Controversies in the Local Management of Lung Cancer

At the time we agreed to edit this special edition of Frontiers in early 2017, the immunotherapy revolution was already established in the second-line treatment of metastatic non-small cell lung cancer (NSCLC) (1–4) and even to a select group of patients (those whose tumor cells expressed PD-L1 of $\geq 50\%$) (5) in the first-line treatment of metastatic disease. We felt that the summarization and understanding of the strengths and benefits of our local modalities were needed prior to the possible integration of immunotherapy into the treatment paradigms of Stages I–III NSCLC and small cell lung cancer (SCLC).

Three of this issue's 10 articles deal with the controversial topic of neo-adjuvant treatment prior to surgery in the management of Stage III lung cancer from the perspective of a multi-disciplinary team (Lewis et al.), a Radiation Oncologist (Sher), and a Thoracic Surgical group (Van et al.). Although no definitive management strategy was recommended as “the way” to manage Stage III NSCLC, each article does a great job of reviewing the current literature, while offering its own unique perspective. The article by Van et al. additionally discusses the rarely investigated topic of salvage surgery.

Four articles dealt with unconventional and/or innovative uses of radiation. Ohri provides a provocative article concerning dose escalation as well as dose de-escalation. Although the idea of dose de-escalation is unpalatable to most Radiation Oncologists, this idea may be considered to be prescient because of the now described synergism between concomitant cytotoxic therapy and immunotherapy in patients with metastatic non-squamous cell NSCLC (6). Kumar et al. offer a pioneering approach to the treatment of Stage III NSCLC which they have labeled quadmodality therapy (concurrent chemo/radiation followed by a stereotactic boost to residual sites of disease, followed by immunotherapy) for the more aggressive treatment of locally advanced lung cancer (Kumar et al.). Both studies, offer different approaches to the hopefully successful integration of immunotherapy with standard chemo/radiation in the near future. Bergsma et al. discuss the rationale and increasingly strong evidence for administering stereotactic body radiation therapy (SBRT) in patients with oligometastatic NSCLC (Bergsma et al.). The work of these investigators and others have led to the NRG LU002 trial that will be investigating the role of SBRT for patients with three or fewer sites of remaining extracranial disease after chemotherapy. Chun et al. have provided a nice review article on heavy ion therapy in the management of NSCLC. Although the recently published prospective randomized trial of passive scattering proton radiation vs intensity-modulated radiation did not reveal any benefit from proton therapy in terms of local failure, radiation pneumonitis, and lung dosimetry, protons resulted in lower heart doses despite larger gross tumor volumes in that trial arm (7). Because heart dose was a strong determinant of overall survival in RTOG 0167's failure to improve outcomes by radiation dose escalation (8), it is hoped that the Bragg–Peak associated with heavy ion therapy will be proven to offer improved local control while limiting normal tissue damage.

OPEN ACCESS

Edited and Reviewed by:

Timothy James Kinsella,
Warren Alpert Medical School of
Brown University, United States

*Correspondence:

John M. Varlotto
john.varlotto@umassmemorial.org

Specialty section:

This article was submitted to
Radiation Oncology,
a section of the journal
Frontiers in Oncology

Received: 30 May 2018

Accepted: 06 June 2018

Published: 18 June 2018

Citation:

Varlotto JM and Veronesi G (2018)
Editorial: Controversies in the Local
Management of Lung Cancer.
Front. Oncol. 8:233.
doi: 10.3389/fonc.2018.00233

The articles by Bloom et al. and Jeremic et al. remind the readers of *Frontiers* how important local management is to the outcomes of SCLC. Bloom et al. offer a comprehensive review of the literature concerning prophylactic cranial irradiation (PCI) in the management of surgically resectable SCLC. Because the incidence of early-stage SCLC is increasing (9), the review of this topic is very important. Although the authors could not reach a definitive conclusion regarding PCI, it is hoped that NRG CC003 demonstrates that hippocampal sparing allows for effective PCI while limiting harmful neurocognitive sequelae and makes the decision to administer PCI in surgically resectable SCLC an easier one. Jeremic et al. comprehensively review consolidative thoracic radiation in the management of extensive stage SCLC from their pioneering work published in 1999 (10) to the more recent work of Slotman et al. (11). They also discuss the potential merits of different radiation fractionation regimens and the use of sequential vs concomitant chemo/radiation regimens. Furthermore, the authors provide a timely update of the conflicting data (12, 13) in regards to the use of PCI for the treatment of patients with extensive stage SCLC.

Although there are no articles in this edition concerning the benefits and risk of SBRT in comparison to surgical resection for patients with Stage I NSCLC, and past retrospective studies (14, 15) and one small publication of prospective data (16) have suggested equivalency, it is hoped that future studies may shed light on proper patient selection for either of these very effective techniques. Until then, the article by Varlotto et al. sheds needed light on how psychosocial aspects can adversely affect the outcomes of surgically resected NSCLC. These authors demonstrated that not being married and having insurance resulted in lower overall survival/lung cancer-specific survival, increased 90-day mortality and a higher incidence of positive nodes upon resection.

During the last decade surgical treatment of lung cancer changed deeply thanks to the large diffusion of minimally invasive approaches that become the standard approach for early-stage lung cancer (17). Thanks to the introduction of robotic technology, more recently, even selected patients with

more advanced disease, even after induction treatment for N2 involvement, have been approached with minimally invasive robotic surgery to reduce surgical trauma, a very important goal in patients with already increased fragility due to systemic treatment (18). Recent publications showed that robotic approach was safe and effective in this subgroup of patients, paving the way to potential changes in the indications of surgical resection in stage III patients, and a rethinking of the better timing for systemic and local treatment (19) with potential advantages to propose minimally invasive surgery upfront for N2 single station disease.

Since our initial agreement to edit this special edition, immunotherapy has continued to radically change the lung cancer landscape. A press release from Merck on April 9, 2018 concerning Keynote-042 demonstrated a superior OS for pembrolizumab (a PD-1 inhibitor) as compared to platinum-based chemotherapy in the first-line setting. The benefit was shown in a population of patients with PD-L1 of $\geq 1\%$ which means that the majority of patients presenting with metastatic NSCLC can benefit from immunotherapy alone as their initial treatment. Furthermore, Keynote 189 (Van et al.) noted an improved overall survival and progression-free survival using combined immunotherapy (pembrolizumab) and chemotherapy as compared to chemotherapy alone. Additionally, consolidative durvalumab (a PD-L1 inhibitor) has shown an impressive progression-free survival benefit in patients with stage III lung cancer after the completion of concurrent chemo/radiation (20). We feel that the articles in this special edition can help the readers of *Frontiers* better understand the strength and limitations of our existing local therapies so that we can better understand how to incorporate immunotherapy in the management of Stage I–III NSCLC and SCLC.

We are very grateful to the tremendous efforts of the fellow authors, the reviewers, and staff at *Frontiers* for making this special edition possible.

AUTHOR CONTRIBUTIONS

GV and JV both wrote this article.

REFERENCES

- Borghaei H, Paz-Ares L, Horn L, Spigel DR, Steins M, Ready NE, et al. Nivolumab versus docetaxel in advanced nonsquamous non-small-cell lung cancer. *N Engl J Med* (2015) 373:1627–39. doi:10.1056/NEJMoa1507643
- Brahmer J, Reckamp KL, Baas P, Crinò L, Eberhardt WE, Poddubskaya E, et al. Nivolumab versus docetaxel in advanced squamous-cell non-small-cell lung cancer. *N Engl J Med* (2015) 373:123–35. doi:10.1056/NEJMoa1504627
- Herbst RS, Baas P, Kim D-W, Felip E, Pérez-Gracia JL, Han JY, et al. Pembrolizumab versus docetaxel for previously treated, PD-L1-positive, advanced non-small-cell lung cancer (KEYNOTE-010): a randomized controlled trial. *Lancet* (2016) 387:3540–50. doi:10.1016/S0140-6736(15)01281-7
- Rittmeyer A, Barlesi F, Watercamp D, Park K, Ciardiello F, von Pawel J, et al. Atezolizumab versus docetaxel in patients with previously treated non-small-cell lung cancer (OAK): a phase 3, open-label, multicentre randomized controlled trial. *Lancet* (2017) 389:255–65. doi:10.1016/S0140-6736(16)32517-X
- Reck M, Rodríguez-Abreu D, Robinson AG, Hui R, Csósz T, Fülöp A, et al. Pembrolizumab versus chemotherapy for PD-L1-positive non-small-cell lung cancer. *N Engl J Med* (2016) 375:1823–33. doi:10.1056/NEJMoa1606774
- Gandhi L, Rodríguez-Abreu D, Gadgeel S, Esteban E, Felip E, De Angelis F, et al. Pembrolizumab plus chemotherapy in metastatic non-small cell lung cancer. *N Engl J Med* (2018) 378(22):2078–92. doi:10.1056/NEJMoa1801005
- Liao Z, Lee JJ, Komaki R, Gomez DR, O'Reilly MS, Fossella FV, et al. Bayesian adaptive randomization trial of passive scattering proton therapy and intensity-modulated photon radiotherapy for locally advanced non-small-cell lung cancer. *J Clin Oncol* (2018). doi:10.1200/JCO.2017.74.0720
- Bradley JD, Paulus R, Komaki R, Masters G, Blumenschein G, Schild S, et al. Standard-dose versus high-dose conformal radiotherapy with concurrent and consolidation carboplatin plus paclitaxel with or without cetuximab for patients with stage IIIA or IIIB non-small-cell lung cancer (RTOG 0617): a randomised, two-by-two factorial phase 3 study. *Lancet Oncol* (2015) 16:187–99. doi:10.1016/S1470-2045(14)71207-0
- Varlotto JM, Recht A, Flickinger JC, Medford-Davis LN, Dyer A-M, DeCamp MM. Lobectomy leads to optimal survival in early-stage small cell lung cancer: a retrospective analysis. *J Thorac Cardiovasc Surg* (2011) 142:538–46. doi:10.1016/j.jtcvs.2010.11.062
- Jeremic B, Shibamoto Y, Nikolic N, Milicic B, Milisavljevic S, Dagovic A, et al. The role of radiation therapy in the combined modality treatment of patients with extensive disease small-cell lung cancer (ED SCLC): a randomized study. *J Clin Oncol* (1999) 17:2092–9. doi:10.1200/JCO.1999.17.7.2092

11. Slotman BJ, van Tinteren H, Praag JO, Kneigiens JL, El Sharouni SY, Hatton M, et al. Use of thoracic radiotherapy for extensive stage small-cell lung cancer: a phase 3 randomised controlled trial. *Lancet* (2015) 385:36–42. doi:10.1016/S0140-6736(14)61085-0
12. Takahashi T, Yamanaka T, Seto T, Harada H, Nokihara H, Saka H, et al. Prophylactic cranial irradiation versus observation in patients with extensive-disease small-cell lung cancer: a multicentre, randomised, open-label, phase 3 trial. *Lancet Oncol* (2017) 18:663–71. doi:10.1016/S1470-2045(17)30230-9
13. Slotman B, Faivre-Finn C, Kramer G, Rankin E, Snee M, Hatton M, et al. Prophylactic cranial irradiation in extensive small-cell lung cancer. *N Engl J Med* (2007) 357:664–72. doi:10.1056/NEJMoa071780
14. Varlotta JM, Fakiris A, Flickinger JC, Medford-Davis L, Liss A, Shelkey J, et al. Matched-pair and propensity score comparisons of outcomes of patients with clinical stage I non-small cell lung cancer treated with resection of stereotactic radiosurgery. *Cancer* (2013) 119:2683–91. doi:10.1002/cncr.28100
15. Verstegen NE, Oosterhuis JW, Palma DA, Rodrigues G, Lagerwaard FJ, van der Elst A, et al. Stage I-II non-small cell lung cancer treated using either stereotactic ablative radiotherapy(SABR) or lobectomy by video-assisted thoracoscopic surgery(VATS): outcomes of a propensity score-matched analysis. *Ann Oncol* (2013) 24:1543–8. doi:10.1093/annonc/mdt026
16. Chang JY, Senan S, Paul M, Mehran RJ, Louie AV, Balter P, et al. Stereotactic ablative radiotherapy versus lobectomy for operable stage I non-small cell lung cancer: a pooled analysis of two randomized trials. *Lancet Oncol* (2015) 16:630–7. doi:10.1016/S1470-2045(15)70168-3
17. Yan TD, Cao C, D'Amico TA, Demmy TL, He J, Hansen H, et al. Video-assisted thoracoscopic surgery lobectomy at 20 years: a consensus statement. *Eur J Cardiothorac Surg* (2014) 45:633–9. doi:10.1093/ejcts/ezt463
18. Cerfolio RJ, Ghanim AE, Dylewski M, Veronesi G, Spaggiari L, Park BJ. The long-term survival of robotic lobectomy for non-small cell lung cancer: a multi-institutional study. *J Thorac Cardiovasc Surg* (2018) 155:778–86. doi:10.1016/j.jtcvs.2017.09.016
19. Veronesi G, Park B, Cerfolio R, Dylewski M, Toker A, Fontaine JP, et al. Robotic resection of stage III lung cancer: an international retrospective study. *Eur J Cardiothorac Surg* (2018). doi:10.1093/ejcts/ezy166
20. Antonia SJ, Villegas A, Daniel D, Vicente D, Murakami S, Hui R, et al. Durvalumab after chemoradiotherapy in stage III non-small-cell lung cancer. *N Engl J Med* (2017) 377:1919–29. doi:10.1056/NEJMoa1709937

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Varlotta and Veronesi. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.