

May 22nd, 4:30 PM - 6:00 PM

Surveillance of Smokeless Tobacco Products Sold in Massachusetts (1997-2010)

Doris Cullen

Massachusetts Department of Public Health

Kevin J. Kane

University of Massachusetts Medical School

Lois Keithly

Massachusetts Department of Public Health

See next page for additional authors

Follow this and additional works at: http://escholarship.umassmed.edu/cts_retreat



Part of the [Community Health and Preventive Medicine Commons](#)

Cullen, Doris; Kane, Kevin J.; Keithly, Lois; Land, Thomas; Paskowsky, Mark; and Li, Wenjun, "Surveillance of Smokeless Tobacco Products Sold in Massachusetts (1997-2010)" (2012). *UMass Center for Clinical and Translational Science Research Retreat*. 11. http://escholarship.umassmed.edu/cts_retreat/2012/posters/11

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.

Presenter Information

Doris Cullen, Kevin J. Kane, Lois Keithly, Thomas Land, Mark Paskowsky, and Wenjun Li

Creative Commons License

This work is licensed under a [Creative Commons Attribution-Noncommercial-Share Alike 3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/).

SURVEILLANCE OF SMOKELESS TOBACCO PRODUCTS SOLD IN MASSACHUSETTS (1997-2010)

Doris Cullen¹, MA; Kevin Kane², MS; Lois Keithly¹ PhD; Thomas Land¹ PhD; Mark Paskowsky¹, MS; Wenjun Li², PhD

¹Massachusetts Department of Public Health, Tobacco Control Program; ²University of Massachusetts Medical School

Contact: Kevin Kane, 508-856-3973, Kevin.Kane1@umassmed.edu

Background

Moist snuff and snus are smokeless tobacco products that are marketed toward smokers for use at times smoking is prohibited. Little information is available about the nicotine content and design features of these products, particularly snus.

Methods

Data on free nicotine levels and design features (pH, total nicotine content, tobacco leaf cut and flavor) of smokeless tobacco products were obtained from manufactures' annual reports to Massachusetts Department of Public Health between 1997 and 2010. Descriptive statistics were provided overall, and by manufacturer and product type.

Results

Mean level of free nicotine in moist snuff remained relatively constant within the range of 3 to 5 mg/g between 1997 and 2010, although this varies by manufacturer, brand, and design characteristics. Mean free nicotine was higher in snus than in moist snuff and increased over time. Average free nicotine content of Swedish Match snus increased sharply since 2003 and reached >6 mg/g in 2010 while that of American brands of snus decreased from 2.9 mg/g in 2001 to 1.7 mg/g in 2010. Swedish Match snus had significantly higher pH than American brands of snus and moist snuff, and experienced a sharp increase from 7.3 in 2003 to 8.3 in 2010. Wide variations in nicotine and pH levels were present among manufacturers. Free nicotine level was associated with pH level of both snus and moist snuff products, and with tobacco leaf-cut for moist snuff. The number of sub-brands of both snus and moist snuff products increased during the study period.

Conclusion

There was notable increasing trend in free nicotine concentration in Swedish Match snus that are marketed to US consumers. The increase in number sub-brands suggests greater marketing efforts by the manufacturers in recent years. Continued surveillance of smokeless tobacco products in MA is an important component of the State's control effort.