

12-28-2011

Table II: Baseline Values of Primary and Secondary Outcomes for Placebo and Botox Groups

Michael K. Flynn

University of Massachusetts Medical School

Follow this and additional works at: http://escholarship.umassmed.edu/obgyn_pp



Part of the [Obstetrics and Gynecology Commons](#)

Repository Citation

Flynn, Michael K., "Table II: Baseline Values of Primary and Secondary Outcomes for Placebo and Botox Groups" (2011). *Obstetrics and Gynecology Publications and Presentations*. 53.

http://escholarship.umassmed.edu/obgyn_pp/53

Table II: Baseline of Placebo and Botox for Primary and Secondary Outcomes (mean [SD])

Table II	Baseline Placebo (N=11)	Baseline Botox (N=20)	p
Age (yrs)	67.45 [12.85]	65.15 [10.83]	0.599
Parity	2.63 [2.64]	1.8 [1.40]	0.156
BMI (kg/m ²)	29.55 [5.76]	34.38 [7.57]	0.113
Incontinence episodes/day	7.99 [3.12]	7.87 [3.75]	0.886
IIQ-7 score	68.39 [28.67]	68.74 [24.99]	0.972
UDI-6 score	61.10 [20.74]	51.66 [20.16]	0.227
# pads/day	5.30 [2.86]	4.81 [3.40]	0.687
24 pad weight (gms)	379.81 [230.67]	563.60 [494.05]	0.256
Voids per 24hr	9.98 [1.54]	11.04 [3.23]	0.317
Diurnal voiding frequency (voids / 8hrs while awake)	4.73 [1.72]	4.81 [1.48]	0.897
Nocturia	1.95 [0.90]	1.87 [1.03]	0.834
Maximum cystometric capacity (cc)	257.37 [161.38]	249.85 [133.77]	0.890
Peak Flow (cc/sec)	13.82 [9.98]	17.60 [6.58]	0.213
Detrusor pressure @ peak Flow (cm H2O)	18.45 [13.73]	17.35 [11.23]	0.810
Volume 1st uninhibited contraction (cc)	117.9 [138.65]	157.89 [107.11]	0.395
Post void residual (cc)	19.73 [21.44]	20.00 [29.02]	0.978