

May 22nd, 4:30 PM - 6:00 PM

Use of whole plant *Artemisia annua* L. as an antimalarial therapy

Elfawal Mostafa

University of Massachusetts - Amherst

Melissa J. Towler

Worcester Polytechnic Institute

Ricardo T. Gazzinelli

University of Massachusetts Medical School

See next page for additional authors

Follow this and additional works at: http://escholarship.umassmed.edu/cts_retreat

 Part of the [Immunology and Infectious Disease Commons](#), [Parasitic Diseases Commons](#), and the [Plant Sciences Commons](#)

Mostafa, Elfawal; Towler, Melissa J.; Gazzinelli, Ricardo T.; Golenbock, Douglas T.; Weathers, Pamela J.; and Rich, Stephen M., "Use of whole plant *Artemisia annua* L. as an antimalarial therapy" (2012). *UMass Center for Clinical and Translational Science Research Retreat*. 46.

http://escholarship.umassmed.edu/cts_retreat/2012/posters/46

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.

Presenter Information

Elfawal Mostafa, Melissa J. Towler, Ricardo T. Gazzinelli, Douglas T. Golenbock, Pamela J. Weathers, and Stephen M. Rich

Creative Commons License

This work is licensed under a [Creative Commons Attribution-Noncommercial-Share Alike 3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/).

USE OF WHOLE PLANT *ARTEMISIA ANNUA* L. AS AN ANTIMALARIAL THERAPY

Elfawal Mostafa¹, Melissa J. Towler², Ricardo Gazzinelli³, Douglas Golenbock³, Pamela J. Weathers², Stephen M. Rich¹

¹Dept. Plant, Soil and Insect Sciences, University of Massachusetts, Amherst, MA 01002; ²Dept. Biology and Biotechnology, Worcester Polytechnic Institute, Worcester MA 01609; ³Infectious Diseases and Immunology, University of Massachusetts Medical School, Worcester, MA

Contact: Mostafa Elfawal, elfawal@psis.umass.edu

Abstract

Anti-malarial drugs are primary weapons for reducing *Plasmodium* transmission in human populations. Successful drugs have been highly efficacious and inexpensive to synthetically manufacture. Emergence of resistant parasites reduces the lifespan of each drug that is developed and deployed. Currently, the most effective anti-malarial is artemisinin (AN), which is extracted from the leaves of *Artemisia annua*. Because of its poor pharmacokinetic properties and prudent efforts to curtail emergence of resistance, AN is prescribed only in combination with other anti-malarials composing an Artemisinin Combination Therapy (ACT). Low yield in the plant and the added cost of secondary anti-malarials in the ACT, make AN in the developing world a costly treatment. Here we show that dried leaves of *A. annua* administered orally are more effective at killing malaria parasites than a comparable dose of purified drug in a rodent malaria model (*P. chabaudi*). A single dose of whole plant (WP) *A. annua* containing 24 mg/kg AN clears 99% of parasites, where a comparable dose of pure drug has half that effect. This is consistent with findings that blood levels of AN are 40 times greater in mice receiving WP versus those given pure drug. We hypothesize that in addition to increasing bioavailability of AN, administration of WP alone may constitute a combination therapy because it contains other anti-malarial compounds that have been shown to synergize with AN. Inexpensive, efficacious, and resilient treatment for malaria based upon WP *A. annua* that can be grown and processed locally would be an effective addition to the global effort to reduce malaria morbidity and mortality.