

May 20th, 12:30 PM

Acute Symptomatic Influenza A Virus (IAV) Infection in Humans Leads to Expansion of Highly Diverse CD8 T Cell Repertoires Crossreactive with Persistent Epstein Barr Virus (EBV)


Anna Gil
University of Massachusetts Medical School

Rabinarayan Misha
University of Massachusetts Medical School

Nuray Aslan
University of Massachusetts Medical School

See next page for additional authors

Follow this and additional works at: http://escholarship.umassmed.edu/cts_retreat

 Part of the [Immunology of Infectious Disease Commons](#), [Immunopathology Commons](#), [Infectious Disease Commons](#), [Pathology Commons](#), [Translational Medical Research Commons](#), and the [Virus Diseases Commons](#)

Gil, Anna; Misha, Rabinarayan; Aslan, Nuray; and Selin, Liisa K., "Acute Symptomatic Influenza A Virus (IAV) Infection in Humans Leads to Expansion of Highly Diverse CD8 T Cell Repertoires Crossreactive with Persistent Epstein Barr Virus (EBV)" (2014). *UMass Center for Clinical and Translational Science Research Retreat*. 18.
http://escholarship.umassmed.edu/cts_retreat/2014/posters/18

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.

Presenter Information

Anna Gil, Rabinarayan Misha, Nuray Aslan, and Liisa K. Selin

Comments

Abstract of poster presented at the 2014 UMass Center for Clinical and Translational Science Research Retreat, held on May 20, 2014 at the University of Massachusetts Medical School, Worcester, Mass.

Creative Commons License

This work is licensed under a [Creative Commons Attribution-Noncommercial-Share Alike 3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/).

Acute symptomatic Influenza A virus (IAV) infection in humans leads to expansion of highly diverse CD8 T cell repertoires crossreactive with persistent Epstein Barr virus (EBV)

Anna Gil, Rabinarayan Mishra, Nuray Aslan, and Liisa K. Selin
University of Massachusetts Medical School, Department of Pathology, Worcester, MA, USA, 01605.

Contact Information:

Anna Gil
Department of Pathology
Selin Lab
Phone: 508.856.3321
Email: Anna.Gil@umassmed.edu

The competence of T cell responses predominantly depends on how efficient T cell receptors (TCRs) are at recognizing antigenic epitopes. We show here that during acute severely symptomatic IAV infection there was an expansion of IAV-M1/EBV-BRLF1 and IAV-M1/EBV-BMLF1 double-tetramer+ cells directly ex-vivo in 5 HLA-A2+ patients. We questioned whether this expansion specific to these two different crossreactive responses would lead to alterations in the TCR repertoire of the IAV-M1₅₈, EBV-BRLF1₁₀₉ and -BMLF1₂₈₀ from before, during and following acute IAV infection. Using staining with Vb MAb we found that T cell responses generated to these epitopes became surprisingly more polyclonal, with the sharing of Vb between M1, BMLF1 and BRLF1 populations which is not seen in healthy donors and which decreased 2 months later consistent with crossreactive expansion. Furthermore, by using single-cell analysis of TCR α and TCR β repertoire of tetramer sorted IAV-M1 cells we showed dramatic changes in specific clonotype usage and in JA and JB family usage during acute IAV infection compared to before infection. In summary, these changes in TCR repertoire during acute symptomatic IAV infection suggest that during severe infection there is a preferential expansion of highly diverse crossreactive responses between IAV and the persistent virus, EBV, which leads to permanent changes in TCR repertoires to both of these two viruses. (NIHAI49320).