

May 20th, 12:30 PM

# Associations of Adipose Tissue Architecture, Adipokines and Inflammatory Markers with Body Mass Index and Gestational Weight Gain in Non- diabetic Pregnancies


Tiffany A. Moore Simas  
*University of Massachusetts Medical School*

Crina Boeras  
*University of Massachusetts Medical School*

Ningning Zhang  
*University of Massachusetts Medical School*

*See next page for additional authors*

Follow this and additional works at: [http://escholarship.umassmed.edu/cts\\_retreat](http://escholarship.umassmed.edu/cts_retreat)

 Part of the [Cellular and Molecular Physiology Commons](#), [Endocrinology Commons](#), [Endocrinology, Diabetes, and Metabolism Commons](#), [Maternal and Child Health Commons](#), [Obstetrics and Gynecology Commons](#), [Reproductive and Urinary Physiology Commons](#), [Translational Medical Research Commons](#), and the [Women's Health Commons](#)

---

Moore Simas, Tiffany A.; Boeras, Crina; Zhang, Ningning; Gaelikman, Olga; Leung, Katherine G.; Wedick, Nicole M.; Robidoux, Laura; Puleo, Jodi Adams; Lee, Mary M.; Corvera, Silvia; and Rosal, Milagros C., "Associations of Adipose Tissue Architecture, Adipokines and Inflammatory Markers with Body Mass Index and Gestational Weight Gain in Non-diabetic Pregnancies" (2014). *UMass Center for Clinical and Translational Science Research Retreat*. 70.  
[http://escholarship.umassmed.edu/cts\\_retreat/2014/posters/70](http://escholarship.umassmed.edu/cts_retreat/2014/posters/70)

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact [Lisa.Palmer@umassmed.edu](mailto:Lisa.Palmer@umassmed.edu).

---

**Presenter Information**

Tiffany A. Moore Simas, Crina Boeras, Ningning Zhang, Olga Gaelikman, Katherine G. Leung, Nicole M. Wedick, Laura Robidoux, Jodi Adams Puleo, Mary M. Lee, Silvia Corvera, and Milagros C. Rosal

**Comments**

Abstract of poster presented at the 2014 UMass Center for Clinical and Translational Science Research Retreat, held on May 20, 2014 at the University of Massachusetts Medical School, Worcester, Mass.

**Creative Commons License**

This work is licensed under a [Creative Commons Attribution-Noncommercial-Share Alike 3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/).

## Associations of Adipose Tissue Architecture, Adipokines & Inflammatory Markers with Body Mass Index and Gestational Weight Gain in Non-diabetic Pregnancies

Tiffany Moore Simas MD, MPH, MEd

UMass Medical School/UMMHC

508-334-6678

[TiffanyA.MooreSimas@UMassMemorial.org](mailto:TiffanyA.MooreSimas@UMassMemorial.org)

Tiffany A. Moore Simas, MD, MPH, MEd<sup>1a</sup>; Crina Boeras, MD<sup>1a</sup>; NingNing Zhang, PhD<sup>1b</sup>; Olga Gealikman, PhD<sup>1c</sup>; Kathy Leung, MPH<sup>1a</sup>; Nicole Wedick, PhD<sup>1d</sup> Laura Robidoux, BS<sup>1a</sup>; Jodi Adams Puleo, MA<sup>1a</sup>; Mary Lee, MD<sup>1b</sup>; Silvia Corvera, PhD<sup>1c</sup>; Milagros Rosal, PhD<sup>1d</sup>

<sup>1</sup>University of Massachusetts Medical School

<sup>a</sup> Obstetrics & Gynecology, <sup>b</sup> Pediatrics, <sup>c</sup> Program in Molecular Medicine, <sup>d</sup> Medicine - Division of Preventive & Behavioral Medicine

**Background:** Some pregnancy weight gain is stored as adipose tissue (AT). Human AT depots vary in their capacity for expansion. Data suggests that subcutaneous (SQ) is adapted for healthy lipid storage. Conversely visceral (V) accumulation is associated with inflammation, obesity-related co-morbidities and Type 2 diabetes (T2DM) risk. We investigated SQ and VAT histologic architecture along with insulin, adipokines and inflammatory markers in relationship to prepregnancy BMI and gestational weight gain (GWG).

**Methods:** Subset of non-diabetic singleton gravidas from the Pregnancy & Postpartum Observational Dietary Study (PPODS), undergoing Cesareans and consenting to SQ & VAT biopsies were included. Average adipocyte size assessed in 10 sections/depot/subject. Maternal and cord blood insulin, adiponectin, leptin, PAI-1, CRP, TNF $\alpha$ , IL1b, IL6 and IL8 evaluated using Luminex MAGPIX, laser based fluorescent analytical test instrumentation with MILLIPLEX<sup>®</sup> multi-analyte panels. GWG determined by difference in pre-pregnancy and last prenatal visit weight.

**Results:** Of 110 subjects enrolled, 19 (17.3%) delivered by Cesarean with 14 consenting to AT sampling, and 7 (50%) having both SQ and VAT available for analysis. These 7 had mean pre-pregnancy BMI  $27.8 \pm 5.6$  kg/m<sup>2</sup> and GWG  $50.0 \pm 25.7$  lb (range 19-83) with delivery age  $39.2 \pm 0.7$  wks. Mean SQ and VAT adipocyte sizes were  $2892 \pm 716$  pixels<sup>2</sup> (range 1866-3775) and  $2427 \pm 641$  pixels<sup>2</sup> (range 1416-3397) respectively ( $p=0.310$ ); neither were statistically correlated with BMI or GWG. Pre-pregnancy BMI statistically correlated with maternal serum insulin ( $0.786$ ,  $p=0.036$ ) at delivery and cord blood leptin ( $0.886$ ,  $p=0.019$ ); GWG statistically correlated only with cord blood adiponectin ( $-0.900$ ,  $p=0.037$ ).

**Conclusions:** In a small sample of normoglycemic pregnancies undergoing Cesareans and AT sampling, adipocyte size was no different in SQ versus visceral depots, and did not correlate with BMI or GWG. Surprisingly, pre-pregnancy BMI but not GWG correlated with maternal serum insulin at delivery, suggesting that pre-pregnancy weight status may be associated with glycemic control at pregnancy end.