

May 20th, 12:30 PM

Analysis of A Novel Nonsense Mutation of Androgen Receptor Gene in Castration-Resistant Prostate Cancer

Dong Han

University of Massachusetts Boston

Kevin Valencia

University of Massachusetts Boston

Shuai Gao

University of Massachusetts Boston

See next page for additional authors

Follow this and additional works at: http://escholarship.umassmed.edu/cts_retreat



Part of the [Cancer Biology Commons](#), [Male Urogenital Diseases Commons](#), and the [Neoplasms Commons](#)

Han, Dong; Valencia, Kevin; Gao, Shuai; and Cai, Changmeng, "Analysis of A Novel Nonsense Mutation of Androgen Receptor Gene in Castration-Resistant Prostate Cancer" (2016). *UMass Center for Clinical and Translational Science Research Retreat*. 34.
http://escholarship.umassmed.edu/cts_retreat/2016/posters/34

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact Lisa.Palmer@umassmed.edu.

Presenter Information

Dong Han, Kevin Valencia, Shuai Gao, and Changmeng Cai

Keywords

prostate cancer, androgen deprivation therapy, castration-resistance, AR-Q784sc

Creative Commons License

This work is licensed under a [Creative Commons Attribution-Noncommercial-Share Alike 3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/).

Analysis of A Novel Nonsense Mutation of Androgen Receptor Gene in Castration-Resistant Prostate Cancer

Dong Han, PhD¹, Kevin Valencia¹, Shuai Gao, PhD¹, and Changmeng Cai, PhD¹

¹Center for Personalized Cancer Therapy, University of Massachusetts Boston

BACKGROUND: Prostate cancer (PCa) is the second leading cause of cancer mortality in American men. The standard treatment for PCa is androgen deprivation therapy (ADT) that blocks transcriptional activity of androgen receptor, but ADT invariably leads to the development of castration-resistant form of PCa (CRPC) with restored activity of AR. CRPC can be further treated with more intensive ADTs, including CYP17-inhibitors to block intratumoral androgen synthesis and more potent AR antagonist (enzalutamide). Most CRPC patients still relapse after a year of treatment and AR activity appears to be restored again. By analyzing the tumor mRNA from a CRPC patient biopsy who had developed resistance to CYP17-inhibitor treatment, we identified a novel nonsense AR mutation on ligand binding domain (Q784sc), which presumably produces a C-terminal truncated form of AR protein that lacks ligand binding domain (LBD) and may mimic certain AR splice variants that also lack LBD. We thus hypothesized that AR-Q784sc mutant may gain the androgen-independent activity or may enhance the transcriptional activity of full-length AR under low androgen environment through dimerization with full-length AR.

METHOD: We utilized luciferase reporter assays to assess the activity of AR-Q784sc in absence or presence of androgens, and with/out full-length AR. We also examined the protein stability and cellular localization of AR-Q784sc using immunoblotting and immunofluorescence. Moreover, stable cell lines that overexpress AR and/or AR-Q784sc were generated to assess the transcription activity on endogenous target genes and on PCa cell growth.

CONCLUSION: AR-Q784sc mutant produces a LBD truncated AR protein that does not have any transcriptional activity by it alone. However, AR-Q784sc can significantly enhance transcriptional activity of full-length AR through dimerization, indicating that the more intensive ADTs may allow CRPC cells to select for LBD truncated form of AR to further enhance the full-length AR activity under low androgen environment.

Dong Han, Ph.D.

Postdoctoral Research Fellow

Center for Personalized Cancer Therapy

Room 4610, Integrated Sciences Complex

University of Massachusetts Boston

100 Morrissey Blvd

Boston, MA 02125

Email: Dong.Han@umb.edu

Lab: 617-287-3447

Cell: 617-893-3515