

May 16th, 1:45 PM

# Altered Baroreflex-Mediated Cardiovascular Responses to Acute Hypotension in Heart Failure Patients Compared to Healthy Adults

Rachel C. Drew

*University of Massachusetts Boston*

Chelsea N. Lindblad

*Penn State College of Medicine*

Soraya M. Samii

*Penn State College of Medicine*

*See next page for additional authors*

Follow this and additional works at: [http://escholarship.umassmed.edu/cts\\_retreat](http://escholarship.umassmed.edu/cts_retreat)



Part of the [Cardiology Commons](#), [Cardiovascular Diseases Commons](#), and the [Translational Medical Research Commons](#)

---

Drew, Rachel C.; Lindblad, Chelsea N.; Samii, Soraya M.; Blaha, Cheryl A.; White, Michael J.; and Sinoway, Lawrence I., "Altered Baroreflex-Mediated Cardiovascular Responses to Acute Hypotension in Heart Failure Patients Compared to Healthy Adults" (2017). *UMass Center for Clinical and Translational Science Research Retreat*. 20.  
[http://escholarship.umassmed.edu/cts\\_retreat/2017/posters/20](http://escholarship.umassmed.edu/cts_retreat/2017/posters/20)

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in UMass Center for Clinical and Translational Science Research Retreat by an authorized administrator of eScholarship@UMMS. For more information, please contact [Lisa.Palmer@umassmed.edu](mailto:Lisa.Palmer@umassmed.edu).

---

**Presenter Information**

Rachel C. Drew, Chelsea N. Lindblad, Soraya M. Samii, Cheryl A. Blaha, Michael J. White, and Lawrence I. Sinoway

**Keywords**

heart failure, acute hypotension

**Creative Commons License**

This work is licensed under a [Creative Commons Attribution-Noncommercial-Share Alike 3.0 License](https://creativecommons.org/licenses/by-nc-sa/3.0/).

## ALTERED BAROREFLEX-MEDIATED CARDIOVASCULAR RESPONSES TO ACUTE HYPOTENSION IN HEART FAILURE PATIENTS COMPARED TO HEALTHY ADULTS

Rachel C. Drew, PhD<sup>1,2</sup>, Chelsea N. Lindblad, MS<sup>3</sup>, Soraya M. Samii, MD, PhD<sup>2</sup>, Cheryl A. Blaha, RN<sup>2</sup>, Michael J. White, PhD<sup>4</sup>, Lawrence I. Sinoway, MD<sup>2</sup>

<sup>1</sup>Department of Exercise and Health Sciences, University of Massachusetts Boston; <sup>2</sup>Penn State Heart and Vascular Institute, Penn State College of Medicine, Hershey, PA; <sup>3</sup>Penn State College of Medicine, Hershey, PA; <sup>4</sup>School of Sport, Exercise and Rehabilitation Sciences, University of Birmingham, Birmingham, UK

Patients with heart failure (HF) exhibit baroreflex dysfunction, which is associated with increased morbidity and mortality. Orthostatic hypotension, a decrease in blood pressure (BP) upon standing, is a condition that often occurs in HF, and may be linked with altered baroreflex responsiveness in this population. However, data on baroreflex-mediated cardiovascular responses to acute hypotension in HF patients are limited. Therefore, 8 HF patients (7 men; mean±SEM 65±3y; ejection fraction 30.5±3.1%) and 7 healthy control (CON) adults (6 men; 65±2y) underwent 7.5 minutes of unilateral lower-limb ischemia via inflation of a thigh cuff on one leg to non-pharmacologically induce acute hypotension upon cuff deflation. Beat-to-beat systolic BP, diastolic BP, and mean arterial BP (MAP; photoplethysmographic finger cuff) and heart rate (HR; electrocardiogram) were recorded continuously before, during, and after cuff inflation. Statistical analysis involved independent-samples *t*-tests. Baseline values were not different between groups (systolic BP: 128±8 vs. 128±4mmHg; diastolic BP: 73±3 vs. 82±5mmHg; MAP: 90±3 vs. 97±4mmHg; HR: 62±2 vs. 56±2b.min<sup>-1</sup> for HF and CON, respectively; *P*>0.05). The magnitude of the induced decrease in MAP was similar in both groups (HF -11±1 vs. CON -12±2mmHg; *P*>0.05). However, the time-to-peak MAP decrease was significantly longer in HF compared to CON (HF 11±2 vs. CON 6±1s; *P*<0.05). The magnitude of the resultant HR increase was not different between groups (HF 2±1 vs. CON 4±1mmHg; *P*>0.05). However, the time-to-peak HR increase was longer in HF compared to CON (HF 9±1 vs. CON 6±1s; *P*<0.05). In conclusion, these results demonstrate that in response to acute hypotension, HF patients exhibit prolonged hypotension and a delayed tachycardic response. These findings suggest that baroreflex-mediated cardiovascular responses to acute hypotension are altered in HF compared to CON, which may be linked to the higher occurrence of orthostatic hypotension experienced in this population.

### **Contact:**

Rachel Drew

University of Massachusetts Boston

[rachel.drew@umb.edu](mailto:rachel.drew@umb.edu)