

7-4-2011

## NSAIDs and atrial fibrillation

Jerry H. Gurwitz

*University of Massachusetts Medical School*

Follow this and additional works at: [https://escholarship.umassmed.edu/meyers\\_pp](https://escholarship.umassmed.edu/meyers_pp)



Part of the [Health Services Research Commons](#), and the [Primary Care Commons](#)

---

### Repository Citation

Gurwitz, Jerry H., "NSAIDs and atrial fibrillation" (2011). *Meyers Primary Care Institute Publications and Presentations*. 533.  
[https://escholarship.umassmed.edu/meyers\\_pp/533](https://escholarship.umassmed.edu/meyers_pp/533)

This material is brought to you by eScholarship@UMMS. It has been accepted for inclusion in Meyers Primary Care Institute Publications and Presentations by an authorized administrator of eScholarship@UMMS. For more information, please contact [Lisa.Palmer@umassmed.edu](mailto:Lisa.Palmer@umassmed.edu).

## EDITORIALS

## NSAIDs and atrial fibrillation

The risk is unproved, but NSAIDs should be used with caution in high risk patients anyway

Jerry H Gurwitz *professor of medicine and executive director*

Meyers Primary Care Institute, a joint endeavour of Fallon Clinic, Fallon Community Health Plan and University of Massachusetts Medical School, Worcester, MA 01609, USA

More than two million Americans and more than four million people in the European Union have paroxysmal or persistent atrial fibrillation.<sup>1</sup> Its prevalence increases dramatically with advancing age, rising from 0.1% in adults younger than 55 years to 9.0% in those aged 80 or more.<sup>2</sup> Atrial fibrillation is associated with an increased long term risk of stroke, heart failure, and death. The healthcare costs related to this condition are substantial, largely as a result of hospital admissions, consultations, diagnostic and therapeutic procedures, and drug treatments. Many patients with atrial fibrillation need lifelong treatment with oral anticoagulants for stroke prevention, which requires careful dosing and laboratory monitoring; safety concerns about the risk of bleeding persist for these patients even under the most ideal systems of care. For these reasons, any opportunity to reduce the risk of atrial fibrillation, particularly in older adults, would be welcome. In the linked case-control study (doi:10.1136/bmj.d3450), Schmidt and colleagues describe an association between the use of non-selective non-steroidal anti-inflammatory drugs (NSAIDs) or selective cyclo-oxygenase-2 (COX 2) inhibitors and atrial fibrillation or flutter.<sup>3</sup>

In 2008, an expert panel convened by the National Heart, Lung, and Blood Institute highlighted that although treatments for atrial fibrillation have been studied extensively, prevention has received relatively little attention.<sup>4</sup> Modifiable risk factors include hypertension, diabetes, obesity, and smoking.<sup>5</sup> Predisposing clinical conditions include heart failure, myocardial infarction, valvular heart disease, thyroid disease, and sleep disordered breathing. These risk factors and predisposing conditions probably interact with various non-modifiable risk factors including age, sex, race, and genetic factors.

The identification of drug related precipitants of atrial fibrillation adds an interesting new dimension to its prevention.<sup>6</sup> Using administrative data for a Danish population from 1999 to 2008, Schmidt and colleagues examined the risk of atrial fibrillation or flutter associated with the use of non-selective NSAIDs or selective COX 2 inhibitors.<sup>3</sup> After adjustment for age, sex, and selected risk factors for atrial fibrillation, they found a significant increase in the risk of atrial fibrillation or flutter with current drug use compared with no use (non-selective NSAIDs: adjusted

odds ratio 1.17, 95% confidence interval 1.10 to 1.24; COX 2 inhibitors: 1.27, 1.20 to 1.34). Risks were greatest for new users compared with non-users (non-selective NSAIDs: 1.46, 1.33 to 1.62; COX 2 inhibitors: 1.71, 1.56 to 1.88), but much less strong for long term users compared with non-users (non-selective NSAIDs: 1.05, 0.98 to 1.13; COX 2 inhibitors: 1.10, 1.03 to 1.18).

An association between use of NSAIDs and atrial fibrillation has important clinical and public health implications because of the high prevalence of use of these agents, particularly among older adults, and the increasing prevalence of atrial fibrillation with advancing age. The validity of the study's findings must be carefully considered, however, because case-control studies are susceptible to unmeasured confounders, potentially limiting the inferences that can be drawn from the results.<sup>7</sup> Adjusting for covariates modestly reduced the strength of the associations between the use of non-selective NSAIDs and COX 2 inhibitors and atrial fibrillation or flutter. However, Schmidt and colleagues lacked data on several important clinical measures, such as body mass index. Obesity is strongly associated with osteoarthritis,<sup>8,9</sup> one of the most common indications for treatment with NSAIDs. This unmeasured confounding variable could have affected the study's findings, because obesity is an established risk factor for atrial fibrillation. Schmidt and colleagues' efforts to estimate the effects of a hypothetical confounder only partially allay these concerns about unmeasured confounding.

The current study is not the first case-control study to report an association between the use of NSAIDs and atrial fibrillation. A study that used data from the United Kingdom General Practice Research Database found that current use of NSAIDs was associated with an increased risk of atrial fibrillation, yet the results of that study differed from the current findings in that risk was greatest for long term users rather than for new users.<sup>10</sup> In both studies, the dose-response relation was not consistent, which makes the association even more tenuous.<sup>7</sup>

The association between NSAIDs and atrial fibrillation does not imply a cause and effect relation. One proposed explanation for the association is: "the existence of an underlying inflammatory condition, increasing the risk of AF [atrial

fibrillation] on the one hand and prompting the use of NSAIDs on the other.”<sup>10</sup> Perhaps a more likely scenario is that these agents trigger or exacerbate clinical conditions, such as heart failure and hypertension, that lie along the causal pathway of atrial fibrillation.

What should clinicians do in practice in the light of current evidence? With uncertainty regarding a plausible biological mechanism, the susceptibility of case-control studies to unmeasured confounders, and inconsistent results in the two studies performed to date, a cautious approach seems warranted in applying the study’s results to the care of patients. However, NSAIDs (non-selective NSAIDs and COX 2 inhibitors) should continue to be used very cautiously in older patients with a history of hypertension or heart failure, who are already at high risk for adverse effects of these drugs, regardless of whether an association between NSAIDs and atrial fibrillation actually exists.

Competing interests: All authors have completed the Unified Competing Interest form at [www.icmje.org/coi\\_disclosure.pdf](http://www.icmje.org/coi_disclosure.pdf) (available on request from the corresponding author) and declare: no support from any organisation for the submitted work; no financial relationships with any organisations that might have an interest in the submitted work in the previous three years; no other relationships or activities that could appear to have influenced the submitted work.

Provenance and peer review: Commissioned; not externally peer reviewed.

- 1 Fuster V, Rydén LE, Cannom DS, Crijns HJ, Curtis AB, Ellenbogen KA, et al. 2011 ACCF/AHA/HRS focused updates incorporated into the ACC/AHA/ESC 2006 guidelines for the management of patients with atrial fibrillation: a report of the American College of Cardiology Foundation/American Heart Association Task Force on practice guidelines. *Circulation* 2011;123:e269-367.
- 2 Go AS, Hylek EM, Phillips KA, Chang Y, Henault LE, Selby JV, et al. Prevalence of diagnosed atrial fibrillation in adults. National implications for rhythm management and stroke prevention: the AnTicoagulation and Risk Factors in Atrial Fibrillation (ATRIA) study. *JAMA* 2001;285:2370-5.
- 3 Schmidt M, Christiansen CF, Mehnert F, Rothman KJ, Sørensen HT. Non-steroidal anti-inflammatory drug use and risk of atrial fibrillation or flutter: population based case-control study. *BMJ* 2011;343:d3450.
- 4 Benjamin EJ, Chen P-S, Bild DE, Mascette AM, Albert CM, Alonso A, et al. Prevention of atrial fibrillation. Report from a National Heart, Lung, and Blood Institute workshop. *Circulation* 2009;119:606-18.
- 5 Huxley RR, Lopez FL, Folsom AR, Agarwal SK, Loefer LR, Soliman EZ, et al. Absolute and attributable risks of atrial fibrillation in relation to optimal and borderline risk factors: The Atherosclerosis Risk in Communities (ARIC) study. *Circulation* 2011;123:1501-8.
- 6 Van der Hooft CS, Heeringa J, van Herpen G, Kors JA, Kingma JH, Stricker BHC. Drug-induced atrial fibrillation. *J Am Coll Cardiol* 2004;44:2117-24.
- 7 Levine M, Walter S, Lee H, Holbrook A, Moyer V. Users’ guides to the medical literature. IV. How to use an article about harm. *JAMA* 1994;271:1615-9.
- 8 Felson DT, Anderson JJ, Naimark A, Walker AM, Meenan RF. Obesity and knee osteoarthritis. The Framingham Study. *Ann Intern Med* 1988;109:18-24.
- 9 Holliday KL, McWilliams DF, Maciewicz RA, Muir KR, Zhang W, Doherty M. Lifetime body mass index, other anthropometric measures of obesity and risk of knee or hip osteoarthritis in the GOAL case-control study. *Osteoarthritis Cartilage* 2011;19:37-43.
- 10 De Caterina R, Ruigómez A, García Rodríguez LA. Long-term use of anti-inflammatory drugs and risk of atrial fibrillation. *Arch Intern Med* 2010;170:1450-5.

Cite this as: *BMJ* 2011;343:d2495